

2015 ANNUAL MONITORING REPORT

SFM Indicators and High Conservation Value Forests



Port Hawkesbury Paper - Woodlands

EXECUTIVE SUMMARY



The 2015 Annual Monitoring Report provides a summary of Port Hawkesbury Paper's safety, environmental, and forest management progress in the Woodlands Unit. Since 2002, Port Hawkesbury Paper (PHP) has been monitoring and reporting on a suite of sustainable forest management indicators to measure its progress towards achieving targets regarding social, economic, and environmental forest values. Long-term monitoring of these values allows the public to better understand PHP's forest management activities, and the goals and objectives we set to ensure our forest management is having a positive impact. This is an important element of continual improvement that PHP strives for every day.

This report also summarizes the effectiveness monitoring program for High Conservation Value Forests (HCVF). These values were first identified in 2008 for Forest Stewardship Council[®] (FSC[®]) certification and updated in March 2015 to include new knowledge and information. Annual monitoring is conducted to assess the effectiveness of the measures used to maintain or enhance the identified values.

Contents

EXECUTIVE SUMMARY	2
CONTENTS	3
ABOUT THE WOODLANDS UNIT	5
Background	5
Crown Land Forest Management Area	6
KEY COMMITMENTS TO SAFETY	9
Woodlands Safety Results	9
KEY COMMITMENTS TO THE ENVIRONMENT	10
Woodlands Environment Results	10
Crown Contractor Audit Results – Winter 2015	12
Crown Contractor Audit Results – Summer 2015	13
Crown Contractor Audit Results – Fall 2015	14
Private Supplier Audit Results – 2015	15
SUSTAINABLE FOREST MANAGEMENT INDICATORS	16
Indicators of Sustainable Forest Management	16
Summary of SFM Indicators	43
High Conservation Value Forest Effectiveness Monitoring Program	45
HCV – American Marten Habitat	47
HCV – Mainland Moose Habitat	49
HCV – Canada Lynx Habitat	51
HCV – Wood Turtle Habitat	53
HCV – Bicknell's Thrush Habitat	55
HCV – Rusty Blackbird Habitat	58
HCV – Roseate Tern Habitat	60
HCV – Olive-Sided Flycatcher Habitat	61
HCV – Eastern Whip-Poor-Will Habitat	62
HCV – Eastern Wood Peewee Habitat	63
HCV – Canada Warbler Habitat	65
HCV – Chimney Swift Habitat	66
HCV – Common Nighthawk Habitat	67
HCV – Wood Thrush Habitat	68
HCV – Black-foam Lichen Habitat	70
HCV – Little Brown Myotis Habitat	71

HCV – Tri-colored Bat Habitat	72
HCV – Northern Myotis Habitat	73
HCV – New Jersey Rush Habitat	74
HCV – Boreal Felt Lichen Occurrences	75
HCV – Vole Ears Lichen Occurrences	77
HCV – Blue Felt Lichen Occurrences	78
HCV – Eastern White Cedar	79
HCV – Black Ash	80
HCV – Frosted Glass Whiskers Habitat	81
HCV – Cold Water Refugia Sub-watersheds	82
HCV – International Bird Areas	84
HCV – Red Spruce	85
HCV – Protected Areas	87
HCV – Special Management Zone Adjacent to Protected Area Boundaries	
HCV – Significant, Old or Unique Forests	95
HCV – Old Forest	96
HCV – Poorly Represented Ecosystems	97
HCV – Connectivity Management Zones	98
HCV – Margaree & St. Mary's River Watershed	99
HCV – Water Supply Intake Areas	101
HCV – Steep Slopes	102
HCV – Viewshed Areas	104
HCV – Traditional Cultural Identity	

ABOUT THE WOODLANDS UNIT



BACKGROUND

With a dedicated staff of 25 people, the company's Woodlands Unit currently manages 30% of the Crown land in Nova Scotia, which represents 58% (roughly 523,000 hectares) of the Crown land in the seven eastern counties. As a result of 45 years of silviculture activities on these lands, the forest will increase in economic activity over the next 20 years.

Our wood supply primarily comes from the seven eastern counties of Nova Scotia with additional wood purchased in central Nova Scotia. The Woodlands Unit provides silviculture services and information on sustainable forest management practices to private woodland owners. Additionally, we provide training on best management practices for Crown and private contractors and operators. In addition to acquiring softwood pulpwood from the managed forest, Port Hawkesbury Paper also manages its forest lands to produce high quality softwood and hardwood logs for sale to local sawmills and buyers.

As the largest Crown license holder in eastern Nova Scotia, we believe that good business includes strong community support and involvement, environmental awareness, continued growth in forest management and contribution to the Nova Scotia economy. The public use of Crown lands for recreation, accessibility, hunting and fishing illustrates the wide variety of values held by the general public. To achieve sustainable forest management, the needs of all stakeholders must be assessed and managed appropriately.

Forest management certification is one of many tools to support the sustainability of Port Hawkesbury Paper. Certification is a voluntary process by which planning, procedures, systems and performance of on-the-ground forestry operations are audited by a qualified and independent third party against a predetermined standard. Forest operations found to be in conformance with the given standard are issued a certificate. Port Hawkesbury Paper supports the mutual recognition of credible forest certification systems that take into account national and regional characteristics such as natural conditions, forest ownership structures and legislation.


Forest operations at Port Hawkesbury Paper are carefully planned to deliver a valuable, sustainable resource that satisfies economic, social, and environmental benefits. More specifically, Port Hawkesbury Paper works diligently to ensure sustainable harvests, increased forest productivity, and protection for wildlife, water, and recreational resources. With the use of high-end computer mapping and software systems, we supervise our operations to strict standards to ensure we continually meet or exceed our expectations for a healthy productive forest for the future.

CROWN LAND FOREST MANAGEMENT AREA

PHP's Defined Forest Area (DFA) is located in the seven eastern counties of Nova Scotia. The geographic extent of the DFA is shown in Figure 1. The company manages approximately 523,000 hectares of Crown lands through a license agreement with the provincial government within the DFA. The land inventory managed by PHP is broken down into four main components (Table 1).

In addition to acquiring wood from PHP company managed lands, the company harvests wood from private woodland owners through short-term stumpage leases. Private wood is also procured from private suppliers that operate on private woodlands located in central and eastern Nova Scotia. Private wood is purchased at roadside and the company provides competitive pricing. In addition, the company provides silviculture services and training in sustainable forest management practices to encourage good stewardship practices.

The public use of Crown lands for recreation, accessibility, hunting and fishing, to name a few, illustrates the wide variety of values held by the general public. Tourism plays an important role in the regional economy; as a result, unique challenges in meeting the needs of all stakeholders must be assessed and managed appropriately. The NSDNR has implemented an integrated resource management (IRM) land use approach for the management of Crown lands.



Crown Lands Licensed to Port Hawkesbury Paper under the Forest Utilization License Agreement

The Crown land-base was assembled using photo-interpreted forest inventory flown in 2008 and 2009 as a base. Historic treatment GIS data were incorporated from PHP and government databases to update the spatial boundaries and attributes of the forest inventory. Wildlife habitat, ecosystem data, special management layers, and hydrology and roads layers were compared, agreed upon and amalgamated where appropriate to create the most recent and accurate dataset possible.

As land-base layers are overlaid, attributes are coded to allow for partitioning of results based on forest and non-forest values. The total land area includes all area, crown wilderness area and non-forested land are removed to create the forested land-base. After removing permanent exclusions (off limits to forest management prescriptions), the remainder is the working land-base which contributes to wood supply. The working land-base is largely occupied by special management lands, which dictate treatment prescription details. The below table summarizes the land-base net down.


Landbase Category					Area(ha)
1. Total Land Area					699,090
1.a Protected Area Land	108,532				
2. Land Outside Protection					590,558
2.a Non-Forested Land	98,782				
3. Forested Landbase (FLB)					491,776
3.a Inoperable/Subjective Removals		38,064	38,064		
3.b Land Use Restrictions (IRM C3)		545	512		
3.c 12 Percent Areas		77,943	65,076		
3.d.1 DNR Lynx Habitat Buffers		24,523	18,836		
3.d.2 DNR Moose Habitat Buffers		10,980	8,314		
3.d.3 Coastal Plains Flora Buffers		106	77		
3.e DNR Old Growth Policy		19,585	7,213		
3.f Aboriginal Offered Lands		0	0		
3.g Other Regional Harvest Exclusions		18,581	7,759		
3 FLB Exclusions Sub-Total			145,850		
4. Working Landbase (WLB)					341,097
4.a Watercourse Buffers (20m)		36,980	15,822	15,822	
4.b Marten Patches		21,804	9,786	9,255	
4.c Deer Wintering Areas		18,082	10,904	10,436	
4.d Mainland Moose Areas (Softwood)		90,454	66,869	63,518	
4.e IRM - C2 Areas		247,010	149,922	112,265	
4 Working Landbase (WLB)				211,296	
5. WLB No Restrictions					134,630

KEY COMMITMENTS TO SAFETY



WOODLANDS SAFETY RESULTS

The Woodlands safety results show a strong trend towards zero lost time accidents and medical aids since 2007. However, having just one accident or medical aid is not acceptable. PHP continues to promote employee and worker safety through effective training programs, monitoring, safety meetings, and other communication means to strive towards our objective of zero safety incidents year after year.



A lost time accident occurs when an employee or worker is injured on the job which results in lost work time.

A medical aid occurs when an employee or worker is injured on the job and requires medical aid, but does not result in lost work time.

KEY COMMITMENTS TO THE ENVIRONMENT



managing our forest areas.

WOODLANDS ENVIRONMENT RESULTS


Monitoring and reporting on Woodlands environmental performance is an integral part of achieving responsible forest management across the working landscape. Harvest contractors working on Crown land on behalf of the company are audited by PHP three times a year - Winter, Summer and Fall. Compliance and performance is checked against a range of items related to layout compliance, operational safety and environmental compliance, and job quality. Contractors must obtain a certain percentage or higher in each category to pass an audit. Deficiencies are addressed through communication and/or re-training.

With a total of 42 audits completed, harvest contractors continue to achieve a very high level of overall compliance and performance as shown in the 2015 results. This is due in large part to the long-standing working relationship between the company and its Crown land harvest contractors. Working together to monitor performance, share information, and strive to continually improve has resulted in strong on-the-ground results.

PHP also audits its private suppliers. A total of 30 suppliers were audited in 2015. Using the previous quarter's deliveries, wood suppliers are randomly chosen to be audited by a PHP Area Supervisor. Private supplier audits are performed on active jobs when possible. However, auditing a completed job may be necessary with smaller suppliers.

Below are summaries of PHP's Crown and private supplier audit program for 2015. Areas of deficiencies are highlighted in orange. If the deficiencies are consistently on-going or deemed to be of significant concern, communications are made to suppliers to improve performance.

CROWN CONTRACTOR AUDIT RESULTS – WINTER 2015









1 WEEKLY INSPECTIONS COMPLETED ACCURATELY

LAYOUT COMPLIANCE

- 2 Followed Cutting Boundaries
- 3 Wildlife clumps (as per instructions)
- 4 Wildlife corridors (50M min)
- 5 Properly buffered watercourses and wetlands
- 6 Residual Trees retained (10/Ha)
- OPERATION COMPLIANCE
- 7 First Aid Supplies
- 8 First Aid trained personnel (Copies to be made available)
- 9 Personal Protective Equipment
- Fire Protection Equipment
- 10 Crew
- 11 -Machines
- 12 Remote Location Emergency Plan
- 13 Proper Warning Signs Posted on 2-way Public Traffic Road
- 14 System to check on Employees who Work Alone
- 15 Current Documentation
- 16 Lock Out Tag Out Policy in place
 - Fuel & Oil Storage:
- 17 · Spill Kit
- 18 · Pumps (able to be locked for transport or off duty.)
- 19 Trailer Permits if not floated.
- 20 · Central collection spot for Hazardous Material. (2.04)
- 21. WHMIS and TDG trained personnel (Copies to be made available)
- 22. Waste oil disposal system in place
- 23 Tanks Properly Labeled / Placarded to TDG and WHIMS Regulations 24 Storage tanks located not closer than 100m from any watercourse
- Water Quality:
- 25 Bridges used and Erosion controlled on approaches to stream crossing
- 26 Temporary bridges removed, water courses cleared of debris
- 27 No evidence of siltation
- 28 Machine Rutting: Within Guidelines (or as permitted by Supervisor) 29 Ground Disturbance: Within Guidelines
- 30 Safety Meeting Minutes
- 31 EMS Training New Employees
- 32 Biodegradable Chain Oil used
- Total


JOB QUALITY

ALL JOBS

Housekeeping:


- 33 -Garbage & Litter collected to be discarded
- 34 No Discarded Parts/Tires
- 35 -Disposed of Hazardous Materials
- 36 Road drains and culverts cleared of debris
- 37 Unmerchantable hardwood trees protected
- 38 Damage To Leave Trees Acceptable
- 39 Plantations
- 40 -Spacing
- Regeneration 41 .
- **NON CLEARCUT TREATMENTS HARVEST**
- 42 Tree Spacing
- 43 Basal Area
- 44 Trail spacing
- 45 Trail width 46 No Damage To Leave Trees

CROWN CONTRACTOR AUDIT RESULTS – SUMMER 2015









1 WEEKLY INSPECTIONS COMPLETED ACCURATELY

LAYOUT COMPLIANCE

2 Followed Cutting Boundaries

- 3 Wildlife clumps (as per instructions)
- 4 Wildlife corridors (50M min)
- 5 Properly buffered watercourses and wetlands
- 6 Residual Trees retained (10/Ha)
- OPERATION COMPLIANCE

OPERATION COMPLIANCE

Water Quality:


- Bridges used and Erosion controlled on approaches to stream crossing 7.
- 8. Temporary bridges removed, water courses cleared of debris
- 9 No evidence of siltation
- 10 Machine Rutting: Within Guidelines (or as permitted by Supervisor)
- 11 Ground Disturbance: Within Guidelines

Total


JOB QUALITY

- Housekeeping:
- 12 . Garbage & Litter collected to be discarded No Discarded Parts/Tires .
- 13 14 • Disposed of Hazardous Materials
- 15 Road drains and culverts cleared of debris
- 16 Unmerchantable hardwood trees protected
- 17 Damage To Leave Trees Acceptable
- Plantations (N/A for 2015)
- 18 -
- Spacing 19 ·
- Regeneration 20 .
- UTILIZATION <= 2 M3/HA NON CLEARCUT TREATMENTS HARVEST 21 UTILIZATION
- 22 Tree Spacing
- 23 Basal Area
- 24 Trail spacing 25 Trail width
- 26 No Damage To Leave Trees









1 WEEKLY INSPECTIONS COMPLETED ACCURATELY

LAYOUT COMPLIANCE

2 Followed Cutting Boundaries

- 3 Wildlife clumps (as per instructions)
- 4 Wildlife corridors (50M min)
- 5 Properly buffered watercourses and wetlands
- 6 Residual Trees retained (10/Ha)
 - **OPERATION COMPLIANCE OPERATION COMPLIANCE**

Water Quality:

- Bridges used and Erosion controlled on approaches to stream crossing
- 7. 8. Temporary bridges removed, water courses cleared of debris
- 9 No evidence of siltation

- 10 Machine Rutting: Within Guidelines (or as permitted by Supervisor)
- 11 Ground Disturbance: Within Guidelines

Total

JOB QUALITY

Housekeeping:

- 12 . Garbage & Litter collected to be discarded
- 13 No Discarded Parts/Tires .
- Disposed of Hazardous Materials 14 -
- 15 Road drains and culverts cleared of debris
- 16 Unmerchantable hardwood trees protected
- 17 Damage To Leave Trees Acceptable
- 18 -Plantations
- Spacing 19 -
- Regeneration 20 -
- 21 UTILIZATION < = 2 M3/HA
- **NON CLEARCUT TREATMENTS HARVEST**
- 22 Tree Spacing
- 23 Basal Area
- 24 Trail spacing
- 25 Trail width
- 26 No Damage To Leave Trees

PRIVATE SUPPLIER AUDIT RESULTS – 2015









Legal Requirements

- Properly buffered watercourses and wetlands.
 Wildlife clumps left on site.
- 3. Coarse woody debris left on site.
- No construction debris/slash in stream.
 No silt source from road entering stream.
- 6. There is no evidence of un-cleaned oil spills over 100 litres. Personal protective equipment
- 7. 8. First Aid Kit
- 9. Training records shown for First Aid, WHMIS, and TDG (as required)- within 1 month
- 10. Fire protection equipment as required for crew and machines
- 11. Fuel tanks properly labeled/placarded/stored/secured to TDG and WHMIS regs. 12. Lock Out Tag Out in place

- Operations Management Plan 14. Operations Management Plan
- 15. Property and cut boundaries flagged. Roads and Landings
- 16. Take off ditches or cross culverts present and functional
- 18. No blockage of natural drainage

19. Haul roads ditched and crowned.

- d graveled wh
- 21. No ditches running into stream.
- 23. Proper size culvert or bridge used.
- 26. Landing location minimizes risk of stream siltation.

Operating Practices

- 27. Forwarder trails on driest locations. 28. Forwarder approach to roadside chosen to minimize damage.
- 30. Supplier has demonstrated efforts to minimize rutting.
- 31. Portable bridge used.
- 32. Immature stands are preserved. 33. Harvested merchantable trees have been fully utilized.
- 34. Wood is piled outside of the special management zone.
- 36. Boundary lines kept clear of brush. 13. Fuel storage is more than 30 metres from stream.
- Equipment
- 38. Fuel and oil leaks are not present on machinery.
- 39. Spill kit available on job.
- Housekeeping 40. Garbage is properly contained and disposed of.
- 41. Used oil is properly disposed of.
- Best Management Practices Other 42. Conservation of known critical wildlife habitat elements, biodiversity & species at risk
- 43. SMPs followed in known Forests with Exceptional Conservation Value
- 44. SMPs followed for known invasive exotic plants and animals
- 45. Known characteristics of special sites preserved
- 46. Harvest residues (slash, limbs, tops) adequately distributed/utilized

SUSTAINABLE FOREST MANAGEMENT INDICATORS



The mission of the Woodlands Unit is to provide a reliable, cost effective and high quality supply of wood through the implementation of Sustainable Forest Management.



The vision of Port Hawkesbury Paper LP Woodlands Unit is "that the forest resources, for which we have responsibility, will sustain healthy ecosystems and natural biodiversity, provide a continuous and expanding supply of valuable wood and conserve the forest characteristics of value to society, wildlife and the environment."



Through the Port Hawkesbury Paper Sustainable Forest Management Policy, the Woodlands Unit implements its Mission and Vision for Sustainable Forest Management (SFM) through 15 Guiding Principles of SFM, 6 Guiding Principles of Wood Procurement, and 16 Standard Practices for SFM.

INDICATORS OF SUSTAINABLE FOREST MANAGEMENT

For over a decade, PHP has been monitoring and reporting on a variety of sustainable forest management (SFM) indicators. To measure sustainable management over time for a range of forest values, indicators were developed to monitor progress in the maintenance or enhancement of those values.

The Woodlands monitoring program for SFM indicators consists of internal assessments and audit programs. Results from these programs are analysed and summarized on an annual basis to determine if targets are being achieved or have failed to meet set targets. Accordingly, this identifies management actions that must be adjusted to achieve desired outcomes.

Local-level SFM indicators were developed according to the Canadian Council of Forest Ministers' criteria for sustainable forest management. These criteria are:

- Conservation of Biological Diversity
- Forest Ecosystem Condition and Productivity
- Conservation of Soil and Water Resources
- Forest Ecosystem Contributions to Global Ecological Cycles
- Multiple Benefits to Society
- Accepting Society's Responsibility for Sustainable Development

CRITERION 1 - CONSERVATION OF BIOLOGICAL DIVERSITY

Conserve biological diversity by maintaining integrity, function, and diversity of living organisms and the complexes of which they are part.

Indicator 1.1 - Species Diversity - Significant Species

OBJECTIVE	Managing and mitigating effects on known occurrences of endangered and threatened species.		
INDICATOR	Annual review of NSDNR's Significant Species and Habitats Database and other species status lists.		
Habitats Datab	ual review of NSDNR's Significant Species an ase, and other species status lists, and propriate management activities where	ıd	VARIANCE None allowed
2015 Update	Bald Eagle Barred Owl Little Brown Bat Moose	P to be Habita pecies ndbase	e used in forest management t database maintained by NSDNR habitats potentially affected by e. The significant species identified ing:

Osprey White-tailed Deer Wood Turtle 42 ha 21,859 ha 419 ha

The 2015 data are used in operational planning and is reviewed by NSDNR during the harvest approval process. Other species status and appropriate management strategies have been incorporated into PHP's High Conservation Value Forest (HCVF) Assessment Report.




Moose (Mainland population) - Endangered

"The native population of moose in Nova Scotia is limited to approximately 1000 individuals in isolated sub-populations across the mainland. The population has declined by at least 20% over the past 30 years with much greater reductions in distribution and population size over more than 200 years, despite extensive hunting closures since the 1930's. The decline is not well understood but involves a complex of threats including: over harvesting, illegal hunting, climate change, parasitic brainworm, increased road access to moose habitat, spread of white-tailed deer, very high levels of cadmium. deficiencies in cobalt and possibly an unknown viral disease.


Moose on Cape Breton Island are not risk as they are abundant and the result of a re-introduction of moose from Alberta in the 1940's."

Source: http://novascotia.ca/natr/wildlife/biodiversity/species-list.asp

indicator 1.2 - Genetic Diversity - Connectivity Management Zones		
OBJECTIVE	To maintain landscape level Connectivity Management Zones (CMZs) according to company connectivity guidelines.	
INDICATOR	Percent of CMZs managed according to connectivity guidelines.	
	pliance level of 100% of the total number of the 100 m solid cover with minimum 30%	VARIANCE 10%
2015 Update All 46 CMZs assessed for 100 m solid cover with minimum 30% crown closure met the connectivity guidelines for 100% compliance.		



Indicator 1.2 - Genetic Diversity - Connectivity Management Zones



Indicator 1.3 - Protected Areas - Protected Area Strategy

OBJECTIVE	To identify and maintain areas reserved from harvest under a protected areas strategy on Crown lands.		
INDICATOR	Proportion of area reserved from harvest under a protected area strategy by EPU.		
TARGET Maintain 12% a protected a	6 of total area reserved from harvest under +/- 1% area strategy.		
2015 Update	EcoregionPercent Protect1 - Cape Breton Taiga2 - Cape Breton Highlands3 - Uplands4 - Eastern5 - Northumberland8 - Atlantic Coastal	ted 72% 59% 25% 19% 19% 42%	



Source: NS Environment, French River Wilderness Area

Indicator 1.4 - Protected Areas - Old Forest

OBJECTIVE	To maintain old forest conditions throughout the landscape.		
INDICATOR Percent of DFA by EPU protected for old forest values.			
TARGET Maintain 8% d	TARGETVARIANCEMaintain 8% of forest areas in old forest condition.+/- 1%		
2015 Update			




Old Forest Area, Guysborough County, Andrea Doucette, PHP

CRITERION 2 - FOREST ECOSYSTEM CONDITION AND PRODUCTIVITY

Conserve forest ecosystem condition and productivity by maintaining the health, vitality, and rates of biological production.

Indicator 2.1 - Forest Ecosystem Resilience - Natural Regeneration

OBJECTIVE	To promote Acadian forest characteristics t	hrough the use of natural regeneration.
INDICATOR	Proportion of appropriate natural regenera	tion in company's reforestation program.
	enerate with appropriate species 50% of eforestation area.	VARIANCE +/- 10%
2015 Update	In 2015, 60% of the total reforestation program was naturally regenerated.	






Red Spruce Natural Regeneration, Derek Geldart, PHP

OBJECTIVE	Reduce clearcut area by applying alternative harvest treatments in appropriate ecoregions.	
INDICATOR	Proportion of total (softwood and hardwood) area harvested using unevenaged, thinning, shelterwood, selection cut and/or partial cut techniques by EPU.	
ecoregions to	clearcut treatments in appropriate represent 40% of total harvest by 2015 Ital harvest by 2025.	VARIANCE +/- 5 Year Period
2015 Update	In 2015, the percent of total harvest representing non-clearcut treatments was 43%.	

Indicator 2.2 - Forest Ecosystem Resilience - Harvest Treatments






Red Spruce Shelterwood, Matthew McKenna, PHP

Indicator 2.3 - Forest Ecosystem Productivity - Forest Health


OBJECTIVE	To minimize fire, insect and disease occurrence across the forest landscape.	
INDICATOR Area (by ha) of forest killed by fire, insect and disease.		
TARGET Less than 500 disease.	ha of forest killed by fire, insect and	VARIANCE + 1000 ha

2015	There was no evidence or recorded data by NS Department of Natural Resources for
Update	total forest killed by insect or disease in 2015. There was approximately 86 hectares
	killed by fire on Crown lands in the 7 eastern counties where PHP operates.



Indicator 2.4 - Forest Ecosystem Productivity - Budworm Hazard

OBJECTIVE	To minimize budworm hazard on the Cape Breton Highlands.	
INDICATOR	Area (by ha) killed by budworm outbreak on the Cape Breton Highlands.	
TARGET To have zero h outbreak.	nectares of forest killed by a budworm	VARIANCE + 800 ha
2015 Update	In 2015, 0 ha of forest in Cape Breton Highlands was killed by a budworm outbreak.	



CRITERION 3 - CONSERVATION OF SOIL AND WATER RESOURCES


Conserve soil and water resources by maintaining their quantity and quality in forest ecosystems.

Indicator 3.1	- Soil	Protection	- Steep	Slopes
---------------	--------	------------	---------	--------

OBJECTIVE	To avoid regular harvesting in identified steep slope areas.		
INDICATOR	DICATOR Area (by ha) of regular harvest in steep slope areas.		
TARGET Maintain no r 30% average	egular harvest in areas with greater than slope.	VARIANCE + 20 ha	
2015 Update	A total of 16 hectares was harvested in areas with greater than 30% average slope. NOTE: This indicator is based on spatial data that identifies slopes > than 30% average using contour data. It is not based on the actual % slope for any given area as could be determined on-the-ground. Therefore, to calculate the results for the indicator, a GIS exercise is done which overlaps the steep slope data with completed harvest jobs to determine non-conformances. Most often, the areas showing as harvested are slivers due to inaccuracies in the data.		



MacKenzie Mountain - CB, Matthew McKenna, PHP



Indicator 3.2 - Water Protection - Watersheds

OBJECTIVE	To protect hydrological functions in all watersheds.			
INDICATOR	FOR Proportion of identified watershed area (that is managed by PHP) in a closed forest condition.			
	TARGETVARIANCEEach watershed shall have 80% of its area (that is managed by PHP) in a closed forest condition 5%			
2015 PHP has identified 17 watersheds throughout its management area that are Update monitored specifically for closed forest condition (> 10 years of age and/or > 2 meters in height). Watershed size ranges from the smallest at 9 hectares to the largest at 51,293 hectares. In 2015, all 17 watersheds had 80% or more of its area in a closed forest condition.				

Watershed Name (and total hectares managed by PHP)	% Closed Forest 2015	% Closed Forest 2014	% Closed Forest 2013	% Closed Forest 2012	% Closed Forest 2011	% Closed Forest 2010
Antigonish Municipal Watershed (647 ha)	100%	100%	100%	100%	100%	100%
Guvsborough 1 Municipal Watershed (2515 ha)	91%	96%	100%	100%	92%	93%
Guvsborough 2 Municipal Watershed (9 ha)	100%	100%	100%	100%	100%	100%
Inverness Municipal Watershed (125 ha)	92%	92%	95%	95%	97%	97%
Pictou Municipal Watershed (40 ha)	100%	100%	90%	88%	100%	100%
Victoria Municipal Watershed (962 ha)	98%	98%	96%	98%	98%	98%
Baddeck River Watershed (15545 ha)	96%	94%	95%	99%	93%	93%
East River Watershed (9468 ha)	93%	93%	95%	94%	89%	89%
Grand River Watershed (5662 ha)	89%	85%	82%	85%	82%	82%
Liscomb River Watershed (12760 ha)	91%	90%	91%	96%	90%	90%
Margaree River Watershed (29118 ha)	88%	89%	98%	100%	91%	92%


Middle River Watershed (15757 ha)	90%	87%	94%	99%	92%	92%
Mira River Watershed (13337 ha)	91%	92%	100%	100%	92%	93%
New Harbour River Watershed (452 ha)	94%	99%	98%	98%	99%	99%
North River Watershed (16108 ha)	86%	83%	92%	96%	79%	84%
River Inhabitant Watershed (4922 ha)	93%	93%	96%	96%	94%	94%
St. Marv's River Watershed (51293 ha)	92%	93%	93%	96%	90%	90%




St. Mary's River Watershed, Andrea Doucette, PHP

Indicator 3.3 - Water Protection - Riparian Zone Management

OBJECTIVE	To protect and maintain all riparian functions.		
INDICATOR Number of riparian zone non-conformance incidents related to Wildlife & Watercourse Protection Regulations.			
TARGET	TARGET VARIANCE		
To have zero r	To have zero non-conformance incidents. None allowed		
2015 There were no infractions of the Wildlife Habitat & Watercourse Protection Update Regulations in 2015.			




The Wildlife Habitat and Watercourse Protection Regulations can be found under Section 40 of the Forests Act. They were developed for application by people working in forestry and are applicable to watercourses and marshes, which include wetlands, lakes, ponds, rivers, streams, creek, estuary, or salt-water body that contains water for at least part of the year. The below image illustrates how special management zones must be established around watercourses and marshes when conducting forestry operations.



Indicator 3.4 - Water Protection - Roads and Stream Crossings

OBJECTIVE	To reduce negative impacts on water quality resulting from road construction.			
INDICATOR	Number of road construction and stream crossing incidents (new and upgrades) according to company guidelines.			
TARGET To have zero r	TARGETVARIANCETo have zero non-conformance incidents.None allowed			
2015 Update				






New bridge installation, Paul MacDonald, PHP

<u>CRITERION 4 - FOREST ECOSYSTEM CONTRIBUTIONS TO GLOBAL</u> <u>ECOLOGICAL CYCLES</u>

Maintain forest conditions and management activities that contribute to the health of global ecological cycles.

Indicator 4.1 - Forest Carbon - Harvest Volume

OBJECTIVE	To reduce carbon emissions.		
INDICATOR Period average volume per hectare harvested.			
TARGET Increase the a next 25 years	average harvest volume by 20% within the	VARIANCE +/- 5 Year Period	
2015 The average volume per hectare harvested was 71 m ³ /ha. This is based on all Update treatments excluding commercial thinnings.			



OBJECTIVE	To contribute to total carbon storage through maintenance of above-ground carbon pool.			
INDICATOR	Total growing stock of both merchantable and non-merchantable species on forest lands.			
TARGET	VARIANCE			
Total growing	otal growing stock of 21,221,500 m ³ +/- 1,000,000 m ³			
2015 Update	The total growing stock for softwood is estimated to be 17,895,038 m ³ and the total growing stock for hardwood is estimated to be 15,019,044 m ³ .			

Indicator 4.2 - Forest Carbon - Total Growing Stock

Indicator 4.3 - Forest Land - Road Construction

OBJECTIVE	To minimize amount of deforested land.			
INDICATOR	INDICATOR Width of permanently disturbed area due to road construction.			
TARGET Reduce avera by 10%.	Reduce average road width of newly constructed roads 5% +/-			
2015 Update	The average road width of newly constructed roads in 2015 was 8.5 meters.			







Photo: Lefort Road Bridge, Cape Breton, PHP

CRITERION 5 - MULTIPLE BENEFITS TO SOCIETY

Sustain flows of forest benefits for current and future generations by providing multiple goods and services.

Indicator 5.1 - Timber and Non-timber Benefits - Hardwood Management

OBJECTIVE	To increase the future value of the hardwood resource.		
INDICATOR Area (by ha) of hardwood management.			
	TARGETVARIANCEManage 2,800 hectares of hardwood in the first five year period of the 2015 Long-Term Plan.+/- 500 ha		
2015 Update			






Single Tree Selection, Obidary Road, Antigonish

Indicator 5.2 - Communities and Sustainability - Harvest Level

OBJECTIVE	To continue to harvest at a sustainable rate.	
INDICATOR Annual harvest level.		
TARGET Harvest 400,0	100 tonnes of softwood per year.	VARIANCE -10%
2015 Update	In 2015, the volume amount harvested wa harvest level).	as 300,102 m ³ of softwood (80% of annual






Softwood clearcut harvest, Liscomb, Matthew McKenna, PHP

Indicator 5.3 - Communities and Sustainability - Third Party Requests

OBJECTIVE	Where appropriate, provide economical, recreational and cultural opportunities to the general public.		
INDICATOR	INDICATOR Number of reasonable third party requests approved.		
TARGET Approve all each year.	ve all reasonable third party requests received 10 requests		
2015 Update	A total of 15 third party requests were received and all were approved.		

Indicator 5.4 - Fair Distribution of Benefits and Costs - Sales to Other Mills


OBJECTIVE To ensure fair distribution of forest resources.		
INDICATOR	Proportion harvest volume sold to other bu	ıyers.
TARGET Sell at least buyers.	40% of annual harvest volume to other	VARIANCE +/- 5 Year Period
2015 Update		



CRITERION 6 - ACCEPTING SOCIETY'S RESPONSIBILITY FOR SUSTAINABLE


Soceity's responsibility for sustainable forest management requires that fair, equitable and effective forest management decisions are made.

OBJECTIVE	BJECTIVE To provide opportunities to better understand, recognize and respect local Mi'kmaw and Treat Rights.	
INDICATOR	INDICATOR Number of opportunities to meet with Mi'kmaw community representatives.	
	mum of six opportunities to meet with viduals annually.	VARIANCE - 1 Meeting
2015 In 2015, the company met on 7 separate occasions with Mi'kmaw communities a Update individuals.		ccasions with Mi'kmaw communities and



Indicator 6.2 - Aboriginal and Treaty Rights - First Nation Agreements

OBJECTIVE To build capacity within Mi'kmaq communities to provide increased emplois opportunities for Mi'kmaw individuals.		unities to provide increased employment
INDICATOR	Volume harvested under agreements with	Mi'kmaq communities.
	the softwood and hardwood volume nder First Nation agreements to 60,000	VARIANCE - 5,000 tonnes
2015 In 2015, the total volume harvested was 21,436 tonnes. Update		1,436 tonnes.




Indicator 6.3 - Respect Aboriginal Traditions - FAC Participation

OBJECTIVE	To ensure meaningful Mi'kmaw participation in the Forest Advisory Committee (FAC).		
INDICATOR	NDICATOR Number of regular FAC meetings attended by a Mi'kmaw representative or desig		
TARGET To engage Mi ^r annually.	Wi'kmaq participation in FAC meetings - 2 meetings		
2015 Update	The Mi'kmaq representative on the company's Forest Advisory Committee did not attend meetings in 2015 (two meetings and one field tour).		

Indicator 6.4 - Public Participation - Opportunities for Participation


OBJECTIVE To engage the public in sustainable forest management planning.		nanagement planning.
INDICATOR	Number of FAC meetings and general publi participation.	c opportunities/avenues for public
	mum of six FAC meetings annually, public and/or regional forest tours.	VARIANCE None

2015 Update	In 2015, there were 10 public outreach events held by PHP. These included FAC meetings,



Indicator 6.5 - Decision-Making - Education and Extension

OBJECTIVE	To advance sustainable forest management principles through commitments to research and extension.		
INDICATOR	DICATOR Level of investment and contribution to education and extension initiatives.		
TARGET		VARIANCE	
The company will provide \$0.03 of direct and/or in-kind +/- 0.01 contributions to education and extension initiatives for every m ³ harvested within the defined forest area.		+/- \$0.01	
2015 In 2015, \$0.81 for every m ³ harvested was contributed to education and extension Update initiatives.			



SUMMARY OF SFM INDICATORS

Since 2002, the Woodlands Unit has developed, monitored and reported on a suite of SFM indicators. During that time, indicators have been revised or removed to be replaced with better indicators based on a new understanding of forest dynamics. PHP has made significant progress in achieving several of our targets and others are on their way. To improve their conditions over time, management decisions and activities will be implemented to result in indicators trending towards their desired targets.

Target Achieved within Acceptable Variance

Target On-going

Target Not Achieved within Acceptable Variance

1.1 Annual review of NSDNR's significant species and habitats database, and other species status lists.	
1.2 Percent of CMZs meeting the 60% closed forest condition guideline.	
1.3 Proportion of area reserved from harvest under a protected areas strategy by EPU.	
1.4 Percent of defined forest area by EPU protected for old forest values.	
2.1 Proportion of natural regeneration in reforestation program.	
2.2 Proportion of total (softwood and hardwood) area harvested using unevenaged, thinning, shelterwood, selection cut and/or partial cut techniques by EPU.	
2.3 Area of forest disturbed by fire, insect and disease.	
2.4 Area (by ha) affected by budworm outbreak on the Cape Breton Highlands.	
3.1 Area (by ha) of regular harvest in steep slope areas.	
3.2 Proportion of identified watershed area (that is managed by PHP) in closed forest condition.	
3.3 Number of riparian zone non-conformance incidents.	
3.4 Number of road construction and stream crossing incidents according to company guidelines.	

4.1 Yearly average volume per hectare harvested.		
4.2 Total growing stock of both merchantable and non-merchantable species on forest lands.		
4.3 Width of permanently disturbed area due to road construction.	•	
5.1 Area (by ha) of hardwood management.		
5.2 Annual harvest level.		
5.3 Number of reasonable 3rd party requests approved.		
5.4 Proportion harvest volume sold to other mills.		
6.1 Number of opportunities to meet with Mi'kmaw community representatives.		
6.2 Volume harvested under agreements with Mi'kmaq communities.		
6.3 Number of regular FAC meetings attended by Mi'kmaw representative or designate.		
6.4. Number of FAC meetings and general public opportunities/ avenues for public participation.		
6.5 Level of investment and contribution to education and extension initiatives.		

HIGH CONSERVATION VALUE FOREST EFFECTIVENESS MONITORING PROGRAM

Introduction

This HCVF Effectiveness Monitoring Program was developed to fulfill the requirements of Principle 9 of the FSC Maritimes Standard. To meet Principle 9 of the standard, forest managers must complete an assessment of their forest lands to identify high conservation values. There are six distinct categories that give an area critical conservation significance. FSC Canada defines an HCVF as:

High Conservation Value Forests are those that that possess one or more of the following attributes:

a) Forest areas containing globally, regionally or nationally significant:

i) Concentrations of biodiversity values (e.g., endemism, endangered species, refugia); and/or

ii) Large landscape level forests, contained within, or containing the management unit, where viable populations of most (if not all) naturally occurring species exist in natural patterns of distribution and abundance.

b) Forest areas that are in or contain rare, threatened or endangered ecosystems.

c) Forest areas that provide basic services of nature in critical situations (e.g., watershed protection, erosion control).

d) Forest areas fundamental to meeting basic needs of local communities (e.g., subsistence, health) and/or critical to local communities" traditional cultural identity (areas of cultural, ecological, economic or religious significance identified in cooperation with such local communities).

Once HCVF's are identified on the land-base, the forest manager must decide how these areas will be managed to maintain or enhance the values that are present. Where values exist, monitoring is needed to show that the prescribed management is effective. PHP's effectiveness monitoring program identifies two stages of monitoring for several HCVF's.

The first level of monitoring is for the basic operational procedures or special management practices that have been identified for the value (e.g. buffer zones, maintenance of special habitat characteristics, protection). This level of monitoring is typically done on an annual basis. It is also important at this stage of monitoring to ensure that PHP is aware of and implementing the best management approach, prescriptions, and/or special management practices as defined by an outside organization. Therefore, PHP will also contact known experts and/or organizations to gather any new available information regarding management or to verify that its current management approach is the best at that time. All HCV's have an identified operational monitoring protocol that is implemented on an annual basis.

The second level of monitoring, if applicable, is strategic monitoring to determine if the HCV attribute(s) are being maintained on the landscape. For example, for a species at risk such as Boreal Felt Lichen, it is important to determine that the identified forest habitat is still suitable and that the species is still present in the habitat. Contrary to operational monitoring, not all HCV's require a strategic level of monitoring. For example, the HCV of old forest are legally protected and therefore, not available for active forest management. Therefore, the attribute of maintaining old forests on the landscape is automatically preserved. Alternatively, strategic monitoring is important for species at risk

when the objective is to maintain areas of good forest habitat for a threatened species, and to ensure that the species is still present in this habitat.

For strategic monitoring, PHP recognizes that there is a required level of involvement by government agencies and/or other organizations for the monitoring of species populations and health. It is PHP's intention to collaborate with these agencies to collect the necessary information.

HCV – AMERICAN MARTEN HABITAT

HCV ATTRIBUTE	Species at Risk – Ha	bitat and Population
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintain and/or enhance American Marten habitat in home range patches	
INDICATOR	Reserve stand structure as required within harvest areas located within the American Marten Habitat Management Zone	
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Habitat management requirements are implemented through the DNR approval process for Crown lands. Monitor implementation of stand structure reserve using TFM. Verify annually that special management practices are still current and/or make operational changes as needed.
DATA SOURCES		COST AND DIFFICULTY
The Forest Manag DNR field audits	ger (TFM); PHP &	Low to Moderate - Dependant on PHP's required level of involvement
	LONG-TERM STR	ATEGIC MONITORING PROGRAM
MANAGEMENT OBJECTIVE	American Marten population recovery	
INDICATOR	Population estimate	es / use within the Marten Habitat Management Zone
MONITORING/REPORTING FREQUENCY Every 3 years. Baseline year is 2014.		MONITORING STRATEGY DNR is responsible for population inventory and studying habitat use. PHP is responsible for obtaining this data.
DATA SOURCES		COST AND DIFFICULTY
American Marten Recovery Team DNR Biologist Peter Austin-Smith		Low to High - Dependant on PHP's required level of involvement
FOREST MANAGEMENT PRESCRIPTION		
These prescriptions are applied throughout the Cape Breton Highlands:		
 12-14 standing and live mature trees per ha must be left evenly spaced throughout the harvest site; 		
- These are in addition to all other requirements of the Wildlife Habitat and Watercourse		

- Large yellow birch trees should be left standing where possible;

- Special management practices for commercial thinning operations in marten patches;

- Harvest sites should maintain at least 100 m3 of coarse woody debris/ha and mean maximum diameter of downed logs should exceed 22 cm.

There are also 30 home range patches established within the Marten Habitat Management Zone. These patches may 'migrate' within the zone, but must be a minimum 500 ha in size, circular in shape, and contain a minimum 60% marten habitat as described in the marten recovery strategy.

2015 MONITORING UPDATE

- 1. All harvest treatments applied throughout the Cape Breton Highlands included the above management prescriptions as required and approved by DNR regional staff.
- 2. A total of 250 hectares (0.3%) was treated in 2015 inside the Marten Habitat Management Zone as per DNR's approval process.
- 3. The American Marten Recovery Strategy (2007) estimates that the Marten population is less than 50 individuals. A re-introduction program began in 2007, which brought 130 individuals into Cape Breton from New Brunswick. A total of 35 individuals were collared, but their movements were lost approximately 6 months after release. DNR does have pictures, have live-trapped, recorded tracks in snow and have received reports/sightings of marten in the Cape Breton Highlands (Peter Austin-Smith, pers. comm., 2013). A goal of the Marten Recovery Team is to have >= 30 marten in Cape Breton by 2010, >= 100 by 2030 and >= 350 by 2040.
- 4. Some work happening related to habitat issues for both American Marten and Canada Lynx (joint recovery team). Still being developed by DNR Species at Risk group. Looking to get more funding to do habitat modeling and management issues in CB Highlands. Nothing approved yet; still in discussion phase.

SUPPORTING DOCUMENTS/REFERENCES

Marten Special Management Practices, NSDNR July 2012; Proposed Marten Recovery Strategy, NSDNR May 2006; Status Report on American Marten, F. Scott June 2001; Weaseling their Way Back into Cape Breton? Assessing the Feasibility of Augmenting the Cape Breton Island Marten Population Through Habitat Suitability and Individual-based Modeling, Rebecca Jepessen, Acadian University Thesis, 2010.

HCV – MAINLAND MOOSE HABITAT

HCV ATTRIBUTE	Species at Risk – Ha	bitat and Population
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT	Maintain and/or enhance Mainland Moose habitat	
OBJECTIVE		
INDICATOR	Reserve stand structure as required within harvest areas located within the five Significant Mainland Moose Population Concentration areas mapped by NSDNR	
MONITORING/RE	PORTING	MONITORING STRATEGY
FREQUENCY Annual		Habitat management requirements are implemented through the DNR approval process for Crown lands. Monitor implementation of stand structure reserve using TFM. Verify annually that special management practices are still current and/or make operational changes as needed.
DATA SOURCES		COST AND DIFFICULTY
The Forest Manag DNR field audits	ger (TFM); PHP &	Low to Moderate - Dependent on PHP's required level of involvement
	LONG-TERM STR	ATEGIC MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Mainland Moose po	pulation recovery
INDICATOR	Population estimate	es / use of population concentration areas
MONITORING/RE	PORTING	MONITORING STRATEGY
FREQUENCY Every 3 years. Baseline year is 2014.		DNR is responsible for population inventory and studying habitat use. PHP is responsible for obtaining this data.
DATA SOURCES		COST AND DIFFICULTY
Mainland Moose Recovery Team DNR Biologist Mark Pulsifer		Low to High - Dependent on PHP's required level of involvement
FOREST MANAGEMENT PRESCRIPTION		
 Moose shelter patches (within 250 metres of the edge of any forest harvest (partial or clearcut) a minimum of two closed canopy coniferous stands > 3 hectares in area) 		

- Moose retention patches (Smaller coniferous must also be retained within each harvest area to provide temporary shelter and concealment)

- Moose buffers (Forested buffers should be retained around and or near open wetlands, watercourses, and waterbodies)

- Roads and access points (Development of roads and improved trails should be avoided where extended extraction trails can be used as an alternative)

- Coarse woody debris (leave tree tops and substantial amounts of woody debris on extraction trails to discourage access)

- Decommission roads to reduce human access

2015 MONITORING UPDATE

- 1. All harvest treatments within the mainland moose concentration areas included the above management prescriptions. Currently, there are concerns within DNR about the special management practices for mainland moose, so future harvest treatments in the moose concentration areas are assessed and approved individually by DNR for specific habitat requirements.
- 2. The Mainland Moose Recovery Plan (2007) estimates approximately 1000-1200 individuals on mainland Nova Scotia. This is the most current information available on mainland moose population numbers.
- 3. No changes have been made to the special management practices for mainland moose as issued by DNR; however, there are current discussions within the department about modifications to the SMPs. A guidance document was provided to PHP on April 13, 2014 which outlined guidance regarding moose retention patches for a specified time period (January 20 to July 18, 2014). The guidance document is now obsolete.
- 4. Mersey Tobeatic Research Institute had been contracted to develop a Mainland Moose Recovery Action Plan, which will outline planned activities for moose habitat monitoring and management, and population recovery. This Plan was delivered to the Department of Natural Resources in spring 2014. PHP has requested a copy from the Department once it becomes publicly available.

SUPPORTING DOCUMENTS/REFERENCES

Mainland Moose Special Management Practices, NSDNR July 2012; Recovery Plan for Mainland Moose in Nova Scotia, March 2007

HCV – CANADA LYNX HABITAT

HCV ATTRIBUTE	Species at Risk – Ha	bitat and Population
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT	Maintain and/or enhance Canada Lynx habitat	
OBJECTIVE		
INDICATOR	Reserve stand structure in lynx bog buffers within harvest areas located throughout the Cape Breton Lynx Range	
MONITORING/RE	PORTING	MONITORING STRATEGY
FREQUENCY Annual		Habitat management requirements are implemented through the DNR approval process for Crown lands. Monitor implementation of stand structure reserve using TFM. Verify annually that special management practices are still current and/or make operational changes as needed.
DATA SOURCES		COST AND DIFFICULTY
The Forest Manag DNR field audits	ger (TFM); PHP &	Low to Moderate - Dependant on PHP's required level of involvement
LONG-TERM STRATEGIC MONITORING PROGRAM		
MANAGEMENT	Canada Lynx popula	tion recovery
OBJECTIVE		
INDICATOR	Population estimate	es / use of treed bog leave areas
MONITORING/RE	PORTING	MONITORING STRATEGY
FREQUENCY Every 3 years. Baseline year is 2014.		DNR is responsible for population inventory and studying habitat use. A joint project between DNR and Acadian University is assessing the efficacy of the 100 meter treed bog buffers. The project began in January 2013 and is expected to end in 2014. PHP is responsible for obtaining the results of this work.
DATA SOURCES		COST AND DIFFICULTY
Canada Lynx Recovery Team DNR Biologist Peter Austin-Smith		Low to High - Dependant on PHP's required level of involvement

FOREST MANAGEMENT PRESCRIPTION

- To supplement red squirrel habitat, which is an important food source for lynx, a wider buffer strip of 100 m of unharvested forest should be left around all treed bogs in the Cape Breton lynx range.

- Where possible, decommission secondary, non-main trunk forest access roads following harvest.

- Plan access roads to have dead ends.

- Plan harvesting to allow decommissioning of sectors of road networks.

- Where possible, narrow and orient road right-of-ways to create shade conditions to reduce snow compaction, thereby reducing ease of travel for coyotes.

- Maintain a continuous supply of >50ha patches of mid-regeneration (15-35-year old) conifer dominated habitat that supports high densities of snowshoe hare over each lynx management unit.

- Create a landscape that will maintain a continuous presence of a mosaic of successional stages, especially mid-regeneration patches that will support resident lynx.

- Employ silvicultural techniques that create, maintain, or prolong use of stands by high populations of snowshoe hares.

- Retain coarse woody debris for denning sites.

2015 MONITORING UPDATE

- 1. All harvest treatments within the Cape Breton lynx range where treed bogs were identified, a 100 meter strip of unharvested forest was left as approved by DNR regional staff.
- 2. The Canada Lynx Recovery Strategy (2005) estimates approximately 50-500 individuals in the Cape Breton lynx range. This is the most current information available on Canada lynx population numbers.
- 3. No changes have been made to the special management practices for Canada Lynx as issued by DNR.
- 5. Some work happening related to habitat issues for both American Marten and Canada Lynx (joint recovery team). Still being developed by DNR Species at Risk group. Looking to get more funding to do habitat modeling and management issues in CB Highlands. Nothing approved yet; still in discussion phase.
- 4. DNR and Acadia University are using transect sampling and back-tracking to record/measure habitat use and presence by Canada lynx, snowshoe hare and red squirrel within 100 meter treed bog buffers and adjacent stands. The results of this work are not yet available.

SUPPORTING DOCUMENTS/REFERENCES

Canada Lynx Special Management Practices NSDNR July 2012; Lynx Recovery Strategy Feb 2007; Endangered Canada Lynx Proposed Project: Assessing the interim 100 metre buffers around highland bogs, DNR 2014

HCV – WOOD TURTLE HABITAT

HCV ATTRIBUTE		
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintain and/or enhance Wood Turtle habitat	
INDICATOR	Implementation of temporal and spatial special management practices for wood turtles	
MONITORING/RE FREQUENCY Annual	PORTING	MONITORING STRATEGY Habitat management requirements are implemented through the DNR approval process for Crown lands. Monitor implementation of temporal and spatial requirements using TFM. Verify annually that special management practices are still current and/or make operational changes as needed.
DATA SOURCES		COST AND DIFFICULTY
The Forest Manag DNR field audits	ger (TFM); PHP &	Low to Moderate - Dependant on PHP's required level of involvement
	LONG-TERM STR	ATEGIC MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Wood Turtle population recovery	
INDICATOR	Population estimates	
MONITORING/REPORTING FREQUENCY Every 3 years. Baseline year is 2014.		MONITORING STRATEGY DNR is responsible for population inventory and studying habitat use. PHP is responsible for obtaining this data.
DATA SOURCES		COST AND DIFFICULTY
Wood Turtle Recovery Team DNR Biologist Mark Pulsifer		Low to High - Dependant on PHP's required level of involvement
FOREST MANAGEMENT PRESCRIPTION		
- Adjust the timing and location of motorized vehicle use for forest management activities to when Wood Turtles are inactive or less likely to be occupying terrestrial babitat (Nov – March)		

- Adjust the timing and location of motorized vehicle use for forest management activities to when Wood Turtles are inactive or less likely to be occupying terrestrial habitat (Nov – March)

- Use temporary bridge crossings for perennial streams to avoid altering stream bank, creating erosion and sedimentation, damaging stream bed, and impacting overwintering turtles.

- Forest management roads and landings should not be constructed parallel to watercourses within 200 m of watercourses where wood turtles occur.

- Special management practices for overwintering, nesting, and basking (see DNR Special Management Practices 2012).

2015 MONITORING UPDATE

- 1. All harvest treatments where wood turtles are presumed to be have the above management prescriptions implemented as approved by DNR regional staff.
- 2. The population of wood turtles in PHP's operating area is estimated to be approximately 3,500 individuals (M. Pulsifer, pers. comm., 2013)
- 3. No changes have been made to the special management practices for wood turtle as issued by DNR.
- 4. Monitoring for new locations has not been a funding priority for DNR, and outside funding has been significantly reduced. A graduate student working on overwintering habitat and communal distribution has just finished his MSc at Acadia. There is nothing significantly different with DNR's understanding of wood turtle distribution within the watershed.
- 5. DNR is not receiving reports of dead turtles that can be linked directly to the forest industry.

SUPPORTING DOCUMENTS/REFERENCES

Wood Turtle Special Management Practices NSDNR July 2012; Protecting and Conserving Wood Turtles: A Stewardship Plan for NS, 2003

HCV – BICKNELL'S THRUSH HABITAT

HCV ATTRIBUTE	Species at Risk – Habita	at and Population
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintain and/or enhance Bicknell's Thrush habitat	
INDICATOR	Implementation of temporal and spatial special management practices for Bicknell's Thrush	
MONITORING/RE	PORTING FREQUENCY	MONITORING STRATEGY
Annual		Identify planned pre-commercial thinning activities in Bicknell's Thrush habitat in the Highlands, so field surveys by Bird Studies Canada can first be conducted to identify presence/absence of the bird during their breeding/nesting season (May, June, July). Monitor implementation of leave patches in thinned/cleared areas using a GIS overlay. Verify annually that special management practices are still current and/or make operational changes as needed.
DATA SOURCES		COST AND DIFFICULTY
The Forest Manager (TFM); PHP and Bird Studies Canada field audits		Low – Bird Studies Canada has consistently completed Bicknell's Thrush surveys each spring if PHP has pre-commercial thinning activities planned for that summer.
	LONG-TERM STRATE	GIC MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Bicknell's Thrush popul	ation recovery
INDICATOR	Population estimates	
MONITORING/REPORTING FREQUENCY MONITORING STRATEGY		
Every 3 years. Baseline year is 2014.		Bird Studies Canada annually monitors high elevation bird species through the High Elevation Landbird Program. Since 2002, the Bicknell's Thrush has been monitored in the Cape Breton Highlands to gather critical information about population status and habitat use. Approximately 20 to 30 routes are monitored each June with the continued goal of monitoring long-term trends of the Bicknell's Thrush.

DATA SOURCES	COST AND DIFFICULTY
- Bird Studies Canada - Becky Stewart/Holly Lightfoot - Cape Breton Highlands National Park – Matt Smith	Low – Bird Studies Canada has consistently taken the lead on Bicknell's Thrush habitat and population research.
 International Bicknell's Thrush Conservation Group (<u>http://www.bicknellsthrush.org/</u>) 	
 High Elevation Landbird Report: 10- year Summary, March 2012 	
FOREST MANAGEMENT PRESCRIPTION	

- Industrial forest stands that support Bicknell's Thrush should remain un-thinned until the trees are no longer at a successional stage that is suitable for nesting, as determined by further research.

- If clearing, construction and/or thinning in Bicknell's Thrush breeding habitat cannot be avoided, activities should be performed outside of the bird breeding season, before June 1st and after July 31st, to prevent the direct destruction of nests, eggs, nestlings, fledglings or adult birds.

- When forest clearing and thinning in Bicknell's Thrush breeding habitat cannot be avoided, patches of intact forest should be left whenever possible. These patches should:

- cover at least one quarter hectare;
- be located 20 to 50 metres from the uncut or unthinned edge; and
- contain intact undisturbed underbrush.

2015 MONITORING UPDATE

- 1. Three Bicknell's Thrush surveys by Bird Studies Canada were conducted in unthinned BITH habitat areas in the CB Highlands working forest in 2015. BITH was not observed to be present in the surveys.
- 2. The current population estimate for the Bicknell's Thrush in Canada is between 40,570 and 49,258 birds, and it was previously estimated that approximately 1,200 breed in NB and NS (HELP Report, March 2012).
- 3. No changes have been made to the special management practices for Bicknell's Thrush as issued by Bird Studies Canada.
- 4. In March 2012, Bird Studies Canada released a 10-year summary report of their High Elevation Landbird Program. The results for Bicknell's Thrush monitoring found that the sampling intensity was not enough to detect statistically significant trends in population and habitat use.
- In 2012-13, Bird Studies Canada refined HELP, using a Generalized Random Tessalation Stratified sampling design to randomly select routes and increase sampling intensity in Cape Breton, thus enabling us to meet international, national and regional information needs (HELP Report, March 2012).

6. In March 2014, Bird Studies Canada released <u>High Elevation Landbird Program: Annual</u> <u>Report for Cape Breton Highlands National 2013-2014</u>, which is a report of their last two years work on monitoring Bicknell's Thrush in the Cape Breton Highlands. The report states:

"With similar sampling intensity as last year, detections of Bicknell's Thrush in 2013 were up slightly on Cape Breton. Routes near Lake of Islands and Tipover Lake in the center of the National Park were especially productive. However, Bicknell's Thrush was absent again this year from Benji's Lake Trail where they had been consistently detected over the past 10 years. Bicknell's Thrush continue to be detected on Money Point, which although is designated an Important Bird Area by Bird Life International, is not afforded any formal protection. Of particular note, Bicknell's Thrush was detected on twice as many routes on Cape Breton compared to northern New Brunswick, despite more routes (49 in total) surveyed in New Brunswick. Furthermore, slightly more individual thrush was detected on Cape Breton (26) than New Brunswick (20).

The combination of Port Hawkesbury Paper "year since last cut" data and Bicknell's Thrush potential habitat data proved successful in predicting areas where Bicknell's Thrush were present in 2013. These results suggest this is a viable approach and new routes will be established using this combination of data for the spring 2014 survey season allowing for broader scale coverage in Cape Breton's industrial forest. Furthermore, adding "year since last cut" as an important variable will help to develop predictive models for other areas in the Maritimes."

SUPPORTING DOCUMENTS/REFERENCES

Conserving the Bicknell's Thrush: Stewardship and Management Practices for High Elevation Forest, 2009; High Elevation Landbird Program: 10-year Report, March 2012

High Elevation Landbird Program: Annual Report for Cape Breton Highlands National 2013-2014

HCV – RUSTY BLACKBIRD HABITAT

HCV ATTRIBUTE	Species at Risk – Ha	bitat and Population	
	OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintain and/or enhance Rusty Blackbird habitat		
INDICATOR	Reserve stand structure in Rusty Blackbird habitat		
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor implementation of reserve stand structure using field audits. Verify annually that special management practices are still current and/or make operational changes as needed.	
DATA SOURCES The Forest Manager (TFM); PHP field audits		COST AND DIFFICULTY Low – PHP currently monitors for riparian buffer management on its operational field audits	
LONG-TERM STRATEGIC MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Rusty Blackbird population recovery		
INDICATOR	Population estimate	25	
MONITORING/REPORTING FREQUENCY Every 3 years. Baseline year is 2014.		MONITORING STRATEGY DNR is responsible for population inventory and studying habitat use. PHP is responsible for obtaining this data.	
DATA SOURCES NSDNR		COST AND DIFFICULTY Low to High - Dependent on PHP's required level of involvement	
FOREST MANAGEMENT PRESCRIPTION			
- PHP implements the Wildlife Habitat and Watercourse Protection Regulations, which is			

- PHP implements the Wildlife Habitat and Watercourse Protection Regulations, which is deemed sufficient for Rusty Blackbirds since they tend to occupy forests near the edges of wetlands, bogs, rivers and streams.

- PHP also establishes 100 meter buffers around all treed bogs in Cape Breton for Canada Lynx habitat management, which is also presumed to be beneficial for Rusty Blackbird.

2015 MONITORING UPDATE

- There were no infractions regarding implementation of Wildlife Habitat and Watercourse Protection Regulations in 2015.

- The population of Rusty Blackbird in Nova Scotia is currently unknown.

SUPPORTING DOCUMENTS/REFERENCES

COSEWIC assessment and status report on the Rusty Blackbird *Euphagus carolinus*in Canada (2006)

HCV – ROSEATE TERN HABITAT

HCV ATTRIBUTE	Species at Risk – Habitat		
	OPERATION	AL MONITORING PROGRAM	
MANAGEMENT OBJECTIVE	Maintain Roseate Tern Habitat		
INDICATOR	Reserve stand struct	ture in Roseate Tern habitat	
MONITORING/RE	PORTING	MONITORING STRATEGY	
FREQUENCY Annual		Maintain a 200-meter buffer zone along the coast at Fisherman's Harbour. Within this buffer zone, no management will occur.	
DATA SOURCES		COST AND DIFFICULTY	
The Forest Manager (TFM)		Low – PHP does not conduct forest management activities within the 200-meter buffer zone.	
FOREST MANAGE	FOREST MANAGEMENT PRESCRIPTION		
 PHP does not conduct forest management activities within the 200-meter buffer zone at Fisherman's Harbour. 			
- Other critical habitat for the Roseate Tern is located on offshore islands.			
2015 MONITORING UPDATE			
A GIS overlay using completed harvest treatment data from 2015 shows that there have been no forest management activities within the 200-meter buffer zone at Fisherman's Harbour.			
SUPPORTING DOCUMENTS/REFERENCES			
Roseate Tern Rec	Roseate Tern Recovery Strategy 2006		

HCV – OLIVE-SIDED FLYCATCHER HABITAT

HCV ATTRIBUTE	Species at Risk – Ha	bitat
	OPERATION	AL MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Olive-sided Flycatcher Habitat	
INDICATOR	Reserve stand structure in Olive-sided flycatcher habitat	
MONITORING/REI FREQUENCY Annual	PORTING	MONITORING STRATEGY PHP is currently participating in a working committee with Bird Studies Canada on a Habitat Stewardship Program project called 'Forest Birds at Risk'. The purpose of the project is to "provide guidance in the collection of habitat information and forest management practices that could potentially enhance conservation efforts for these species, and to evaluate their value in the Maritime context." Since this project is currently underway, special management practices have not yet been defined for the above bird species.
DATA SOURCES		COST AND DIFFICULTY
ACCDC		Low – PHP does not yet implement SMP's

FOREST MANAGEMENT PRESCRIPTION

The habitat characteristics of olive-sided flycatcher are minimally impacted by forestry activities due to where they typically inhabit. PHP leaves snags throughout its operations and the presence of tall trees can be found in several PHP silviculture treatments (e.g. single selection, group selection, partial cuts, shelterwoods, patch cuts, red spruce management). PHP also provides habitat features such as forest edges, openings, and clearcuts. Therefore, specific special management practices are deemed to be not necessary at this time and PHP believes there is adequate habitat across the forest management area. However, if special management practices are developed by government or other agencies, they will be implemented as applicable to forest management. If an active nest is located during regular operational activities, the activity will be stopped and the local DNR Wildlife Biologist will be notified so appropriate measures can be implemented.

2015 MONITORING UPDATE

The 'Forest Birds at Risk' project with Bird Studies Canada is still active, so SMP's have not yet been finalized for this species.

SUPPORTING DOCUMENTS/REFERENCES

HCV ATTRIBUTE	Species at Risk – Ha	bitat
	OPERATION	AL MONITORING PROGRAM
MANAGEMENT OBJECTIVE INDICATOR	Eastern Whip-poor-will Habitat Reserve stand structure in Eastern Whip-poor-will habitat	
MONITORING/RE FREQUENCY Annual	PORTING	MONITORING STRATEGY PHP is currently participating in a working committee with Bird Studies Canada on a Habitat Stewardship Program project called 'Forest Birds at Risk'. The purpose of the project is to "provide guidance in the collection of habitat information and forest management practices that could potentially enhance conservation efforts for these species, and to evaluate their value in the Maritime context." Since this project is currently underway, special management practices have not yet been defined for the above bird species.
DATA SOURCES ACCDC		COST AND DIFFICULTY Low – PHP does not yet implement SMP's

HCV - EASTERN WHIP-POOR-WILL HABITAT

FOREST MANAGEMENT PRESCRIPTION

The habitat characteristics of whip-poor-will are minimally impacted by forestry activities due to where they typically inhabit. PHP creates forest edges and openings through active management, as well as even-aged stands that can contain well-spaced trees. Therefore, precautionary specific special management practices are deemed to be not necessary at this time and PHP believes there is adequate habitat across the forest management area. However, if special management practices are developed by government or other agencies, they will be implemented as applicable to forest management. If an active nest is located during regular operational activities, the activity will be stopped and the local DNR Wildlife Biologist will be notified so appropriate measures can be implemented.

2015 MONITORING UPDATE

The 'Forest Birds at Risk' project with Bird Studies Canada is still active, so SMP's have not yet been finalized for this species.

SUPPORTING DOCUMENTS/REFERENCES

HCV ATTRIBUTE	Species at Risk – Habitat	
	OPERATION	AL MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Eastern Wood Peewee Habitat	
INDICATOR	Reserve stand structure in Eastern wood peewee habitat	
MONITORING/RE FREQUENCY Annual	PORTING	MONITORING STRATEGY PHP is currently participating in a working committee with Bird Studies Canada on a Habitat Stewardship Program project called 'Forest Birds at Risk'. The purpose of the project is to "provide guidance in the collection of habitat information and forest management practices that could potentially enhance conservation efforts for these species, and to evaluate their value in the Maritime context." Since this project is currently underway, special management practices have not yet been defined for the above bird species.
DATA SOURCES ACCDC		COST AND DIFFICULTY Low – PHP does not yet implement SMP's

FOREST MANAGEMENT PRESCRIPTION

The Eastern wood peewee can be impacted by forest management activities since this species of bird prefers mature and intermediate age stands of deciduous and mixed forests. However, PHP manages the forest management area by creating a range of age classes through forest modeling, long-term planning, and operational planning. Also, PHP manages deciduous and mixed forest stands with a variety of harvest treatments that can still maintain adequate forest structure (e.g. single selection, group selection, partial cuts, shelterwoods, patch cuts). The above figure shows the Maritime Breeding Bird Atlas data for the species. In eastern Nova Scotia where PHP operates, the breeding evidence shows a variety of results with the most common type being 'possible' evidence.

Forestry practices that maintain large tracts of intermediate aged forest with closed canopy and limited clear cuts (less than 10 ha) along with thinning to remove mature trees and largediameter woody growth should provide adequate habitat for Eastern Wood-Pewees (Stauffer and Best 1980, Crawford et al. 1981). Therefore, precautionary specific special management practices are deemed to be not necessary at this time and PHP believes there is adequate habitat across the forest management area. However, if special management practices are developed by government or other agencies, they will be implemented as applicable to forest management. If an active nest is located during regular operational activities, the activity will be stopped and the local DNR Wildlife Biologist will be notified so appropriate measures can be implemented.

2015 MONITORING UPDATE

The 'Forest Birds at Risk' project with Bird Studies Canada is still active, so SMP's have not yet been finalized for this species.

SUPPORTING DOCUMENTS/REFERENCES

HCV – CANADA WARBLER HABITAT

HCV ATTRIBUTE	Species at Risk – Ha	bitat	
	OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Canada Warbler Hal	pitat	
INDICATOR	Reserve stand structure in Canada warbler habitat		
MONITORING/RE FREQUENCY Annual	PORTING	MONITORING STRATEGY PHP is currently participating in a working committee with Bird Studies Canada on a Habitat Stewardship Program project called 'Forest Birds at Risk'. The purpose of the project is to "provide guidance in the collection of habitat information and forest management practices that could potentially enhance conservation efforts for these species, and to evaluate their value in the Maritime context." Since this project is currently underway, special management practices have not yet been defined for the above bird species.	
DATA SOURCES		COST AND DIFFICULTY Low – PHP does not yet implement SMP's	
		Low – The does not yet implement Sivir S	

FOREST MANAGEMENT PRESCRIPTION

The habitat characteristics of Canada warbler are minimally impacted by forestry activities due to where they typically inhabit. PHP creates regenerating stand structures and forest edge through active management, which is preferred by this species, but also avoid steep slope areas, ravines, swamps, and bogs. The provision of stumps and coarse woody debris left by PHP is also believed to create understory conditions preferred by the Canada warbler. Furthermore, PHP does not contribute to habitat loss by converting swamp forests to agricultural land. Therefore, precautionary special management practices are deemed to be not necessary at this time and PHP believes there is adequate habitat across the forest management area. However, if special management practices are developed by government or other agencies, they will be implemented as applicable to forest management. If an active nest is located during regular operational activities, the activity will be stopped and the local DNR Wildlife Biologist will be notified so appropriate measures can be implemented.

2015 MONITORING UPDATE

The 'Forest Birds at Risk' project with Bird Studies Canada is still active, so SMP's have not yet been finalized for this species.

SUPPORTING DOCUMENTS/REFERENCES

HCV – CHIMNEY SWIFT HABITAT

HCV ATTRIBUTE	Species at Risk – Ha	bitat
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Chimney Swift Habitat	
INDICATOR	Reserve stand structure in Chimney swift habitat	
MONITORING/REI FREQUENCY Annual	PORTING	MONITORING STRATEGY PHP is currently participating in a working committee with Bird Studies Canada on a Habitat Stewardship Program project called 'Forest Birds at Risk'. The purpose of the project is to "provide guidance in the collection of habitat information and forest management practices that could potentially enhance conservation efforts for these species, and to evaluate their value in the Maritime context." Since this project is currently underway, special management practices have not yet been
DATA SOURCES		defined for the above bird species.
ACCDC		Low – PHP does not yet implement SMP's

FOREST MANAGEMENT PRESCRIPTION

The Chimney swift can be impacted by forest management activities since this species of bird may nest in wooded areas with large diameter trees. Currently, there are no special management practices identified for forest managers regarding Chimney swift habitat. However, since the feeding and nesting habitat relies heavily on urban and suburban areas where there is an abundance of chimneys for nesting, PHP believes it currently has a low impact on Chimney swift populations.

2015 MONITORING UPDATE

The 'Forest Birds at Risk' project with Bird Studies Canada is still active, so SMP's have not yet been finalized for this species.

SUPPORTING DOCUMENTS/REFERENCES

HCV – COMMON NIGHTHAWK HAB	ITAT
----------------------------	------

HCV ATTRIBUTE	Species at Risk – Ha	bitat
	OPERATION	AL MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Common Nighthawk Habitat	
INDICATOR	Reserve stand structure in Common nighthawk habitat	
MONITORING/REF FREQUENCY Annual	PORTING	MONITORING STRATEGY PHP is currently participating in a working committee with Bird Studies Canada on a Habitat Stewardship Program project called 'Forest Birds at Risk'. The purpose of the project is to "provide guidance in the collection of habitat information and forest management practices that could potentially enhance conservation efforts for these species, and to evaluate their value in the Maritime context." Since this project is currently underway, special management practices have not yet been defined for the above bird species.
DATA SOURCES		COST AND DIFFICULTY
ACCDC		Low – PHP does not yet implement SMP's

FOREST MANAGEMENT PRESCRIPTION

The Common Nighthawk prefers some habitats where PHP does not operate such as rocky areas, sandy areas, and wetlands. However, they do prefer open wooded areas, which PHP does create through its forest management (e.g. clearcuts, partial cuts, shelterwoods, selection cuts). The Common Nighthawk Recovery Strategys lists a variety of threats including changes in natural processes, climate and natural disasters, accidental mortality, pollution, exotic or invasive species, and habitat loss or degradation. Types of habitat loss include change in roof construction and materials, residential and commercial development, agriculture, and logging and wood harvesting. It is currently unknown if logging and wood harvesting causes a significant severity to populations with a low causal certainty that there is a high degree of evidence linked to the threat of logging.

2015 MONITORING UPDATE

The 'Forest Birds at Risk' project with Bird Studies Canada is still active, so SMP's have not yet been finalized for this species.

SUPPORTING DOCUMENTS/REFERENCES

HCV – WOOD THRUSH HABITAT

OPERATIONAL MONITORING PROGRAM MANAGEMENT OBJECTIVE Wood Thrush Habitat INDICATOR Reserve stand structure in Wood thrush habitat MONITORING/REPORTING FREQUENCY Annual MONITORING STRATEGY PHP is currently participating in a working committee with Bird Studies Canada on a	HCV ATTRIBUTE	Species at Risk – Ha	bitat
OBJECTIVE Wood Thrush Habitat INDICATOR Reserve stand structure in Wood thrush habitat MONITORING/REPORTING MONITORING STRATEGY FREQUENCY PHP is currently participating in a working committee with Bird Studies Canada on a	OPERATIONAL MONITORING PROGRAM		
MONITORING/REPORTINGMONITORING STRATEGYFREQUENCYPHP is currently participating in a working committee with Bird Studies Canada on a		Wood Thrush Habitat	
FREQUENCYPHP is currently participating in a working committee with Bird Studies Canada on a	INDICATOR	Reserve stand struct	ture in Wood thrush habitat
Habitat Stewardship Program project called 'Forest Birds at Risk'. The purpose of the project is to "provide guidance in the collection of habitat information and forest management practices that could potentially enhance conservation efforts for these species, and to evaluate their value in the Maritime context." Since this project is currently underway, special management practices have not yet been defined for the above bird species.	FREQUENCY	PORTING	PHP is currently participating in a working committee with Bird Studies Canada on a Habitat Stewardship Program project called 'Forest Birds at Risk'. The purpose of the project is to "provide guidance in the collection of habitat information and forest management practices that could potentially enhance conservation efforts for these species, and to evaluate their value in the Maritime context." Since this project is currently underway, special management practices have not yet been
DATA SOURCES COST AND DIFFICULTY	DATA SOURCES		COST AND DIFFICULTY
ACCDC Low – PHP does not yet implement SMP's	ACCDC		Low – PHP does not yet implement SMP's

FOREST MANAGEMENT PRESCRIPTION

Currently, there are no required management practices for Wood Thrush in Nova Scotia or Canada. Regardless, given the preferred breeding habitat of Wood Thrush in mature deciduous and mixed-wood forests, PHP believes its uneven-aged and mixedwood forest management techniques in these forest types do not greatly impact the breeding requirements of the Wood Thrush. The COSEWIC 2012 report supports this notion by stating that "the species is relatively tolerant of forest management activities that are conducted on a small spatial scale (i.e. single-tree, group selection cuts, uneven-age forest management, selective removal of mature trees). The report further states that Sugar Maple and American Beech are preferred species for nesting. PHP manages tolerant Sugar Maple stands using only single or group selection depending on tree quality. American Beech is present throughout the forest management area and pure stands are not managed, but if found dispersed throughout a hardwood stand, it is managed as necessary to meet the forest management prescription. Additionally, PHP does not apply herbicides in its forest management area, which allows for the continued natural growth of deciduous trees and shrubs in forest stands.

2015 MONITORING UPDATE

The 'Forest Birds at Risk' project with Bird Studies Canada is still active, so SMP's have not yet been finalized for this species.

SUPPORTING DOCUMENTS/REFERENCES

HCV – BLACK-FOAM LICHEN HABITAT

HCV ATTRIBUTE	Species at Risk – Habitat		
	OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Black-foam Lichen Habitat		
INDICATOR	Reserve stand structure in Black-foam lichen habitat		
MONITORING/RE FREQUENCY Annual	PORTING	MONITORING STRATEGY Confirm with NS Department of Natural Resources, NS Environment, Atlantic Canada Conservation Data Centre, and Mersey Tobeatic Research Institute if any new locations of black- foam lichen have been discovered on PHP's Crown lease.	
DATA SOURCES ACCDC		COST AND DIFFICULTY Low – PHP does not yet implement SMP's	
FOREST MANAGEMENT PRESCRIPTION			

Although there are no special management practices developed by government or other responsible agencies, PHP will implement the same management practice applied to boreal felt lichen and blue felt lichen. Known and confirmed locations of black-foam lichen will have a 100 meter no harvest buffer around the site.

2015 MONITORING UPDATE

No new locations of black-foam lichen have been found on PHP's Crown lease. Currently, there is still only one known location of this lichen in the seven eastern counties where PHP operates and it is in the Cape Breton Highlands National Park. An expert lichenologist also verified that this lichen is still most commonly found in southwest Nova Scotia.

SUPPORTING DOCUMENTS/REFERENCES

NS Department of Natural Resources, NS Environment, ACCDC, MTRI

HCV – LITTLE BROWN MYOTIS HABITAT

HCV ATTRIBUTE	Species at Risk – Habitat	
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Little Brown Myotis Habitat	
INDICATOR	Reserve stand structure in Little brown myotis habitat	
MONITORING/RE FREQUENCY Annual	PORTING	MONITORING STRATEGY Monitor Environment Canada's work on the development of beneficial management practices for the forest industry. Monitor Crown contractor audits to verify that unmerchantable trees, such as snags, wolf trees, and cavity trees, are being retained on site.
DATA SOURCES ACCDC		COST AND DIFFICULTY Low – PHP does not yet implement SMP's

FOREST MANAGEMENT PRESCRIPTION

Currently in Nova Scotia, there are no best forest management practices required for bats. Regarding wolf trees which are important for roosting, the NS Forest Wildlife Guidelines of 1988, which is now a Crown land policy, recommends that snags, wolf trees, and cavity trees be left on harvest sites as much as possible. Most often, wolf trees are so large and difficult to harvest because of many branches, and have low economic value, that PHP leaves on site. PHP is currently managing the forest in a variety of ways that benefit bat habitat needs, based on a 2006 report called "Forest Management & Bats" by Bat Conservation International which lists a variety of forest management activities that can support bat habitat needs.

2015 MONITORING UPDATE

Currently, there are no beneficial management practices developed for the forest industry. A large colony of approximately 300 females was recently found (July 2016) in Nova Scotia. Due to the highly sensitive nature of bat populations, its location was not made publicly available. However, it was confirmed to PHP by a DNR management executive that the colony was not found on PHP's Crown lease. The 2015 Crown operations audits show that unmerchantable trees were left on harvest sites.

SUPPORTING DOCUMENTS/REFERENCES

NS Department of Natural Resources, Environment Canada

HCV – TRI-COLORED BAT HABITAT

HCV ATTRIBUTE	Species at Risk – Habitat	
	OPERATION	AL MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Tri-colored Bat Habitat	
INDICATOR	Reserve stand structure in Tri-colored bat habitat	
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor Environment Canada's work on the development of beneficial management practices for the forest industry. Monitor Crown contractor audits to verify that unmerchantable trees, such as snags, wolf trees, and cavity trees, are being retained on site.
DATA SOURCES ACCDC		COST AND DIFFICULTY Low – PHP does not yet implement SMP's
FOREST MANAGEMENT PRESCRIPTION		

FOREST MANAGEMENT PRESCRIPTION

Currently in Nova Scotia, there are no best forest management practices required for bats. Regarding wolf trees which are important for roosting, the NS Forest Wildlife Guidelines of 1988, which is now a Crown land policy, recommends that snags, wolf trees, and cavity trees be left on harvest sites as much as possible. Most often, wolf trees are so large and difficult to harvest because of many branches, and have low economic value, that PHP leaves on site. PHP is currently managing the forest in a variety of ways that benefit bat habitat needs, based on a 2006 report called "Forest Management & Bats" by Bat Conservation International which lists a variety of forest management activities that can support bat habitat needs.

2015 MONITORING UPDATE

Currently, there are no beneficial management practices developed for the forest industry. The 2015 Crown operations audits show that unmerchantable trees were left on harvest sites.

SUPPORTING DOCUMENTS/REFERENCES

NS Department of Natural Resources, Environment Canada

HCV – NORTHERN MYOTIS HABITAT

HCV ATTRIBUTE	Species at Risk – Habitat	
	OPERATION	AL MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Northern Myotis Habitat	
INDICATOR	Reserve stand structure in Northern myotis habitat	
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor Environment Canada's work on the development of beneficial management practices for the forest industry. Monitor Crown contractor audits to verify that unmerchantable trees, such as snags, wolf trees, and cavity trees, are being retained on site.
DATA SOURCES		COST AND DIFFICULTY
ACCDC Low – PHP does not yet implement SMP's FOREST MANAGEMENT PRESCRIPTION		

Currently in Nova Scotia, there are no best forest management practices required for bats. Regarding wolf trees which are important for roosting, the NS Forest Wildlife Guidelines of 1988, which is now a Crown land policy, recommends that snags, wolf trees, and cavity trees be left on harvest sites as much as possible. Most often, wolf trees are so large and difficult to harvest because of many branches, and have low economic value, that PHP leaves on site. PHP is currently managing the forest in a variety of ways that benefit bat habitat needs, based on a 2006 report called "Forest Management & Bats" by Bat Conservation International which lists a variety of forest management activities that can support bat habitat needs.

2015 MONITORING UPDATE

Currently, there are no beneficial management practices developed for the forest industry. The 2015 Crown operations audits show that unmerchantable trees were left on harvest sites.

SUPPORTING DOCUMENTS/REFERENCES

NS Department of Natural Resources, Environment Canada

HCV – NEW JERSEY RUSH HABITAT

HCV ATTRIBUTE	Species at Risk – Ha	bitat	
	OPERATION	AL MONITORING PROGRAM	
MANAGEMENT OBJECTIVE	Maintain New Jersey Rush Habitat		
INDICATOR	Administratively protect New Jersey Rush habitat identified in NSDNR's Significant Habitat database and Atlantic Canada Conservation Data Centre database		
MONITORING/RE	PORTING	MONITORING STRATEGY	
FREQUENCY Annual		Monitor annual harvest operations to ensure New Jersey Rush habitat is administratively protected from all forest management activities.	
DATA SOURCES		COST AND DIFFICULTY	
The Forest Manager (TFM)		Low – PHP does not conduct forest management activities within New Jersey Rush habitat	
FOREST MANAGE	FOREST MANAGEMENT PRESCRIPTION		
 PHP does not conduct forest management activities within New Jersey Rush habitat identified in NSDNR's Significant Habitat database and Atlantic Canada Conservation Data Centre database 			
2015 MONITORING UPDATE			
A GIS overlay using completed harvest treatment data from 2015 shows that there have been no forest management activities in identified New Jersey Rush habitat.			
SUPPORTING DOCUMENTS/REFERENCES			
Recovery Strategy and Management Plan for Multiple Species of Atlantic Coastal Plain Flora 2010			

HCV – BOREAL FELT LICHEN OCCURRENCES

HCV ATTRIBUTE	Species at Risk – Ha	abitat and Population
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT	Protect identified locations of Boreal Felt Lichen	
OBJECTIVE		
INDICATOR	Administratively protect identified locations of Boreal Felt Lichen by establishing 100-meter buffer around site	
MONITORING/RE	PORTING	MONITORING STRATEGY
FREQUENCY Annual		Monitor annual harvest operations to identify areas needing Boreal Felt Lichen presence/absence surveys prior to active operations. Locations of Boreal Felt Lichen are buffered by 100 meters and excluded from forest management activities.
DATA SOURCES		COST AND DIFFICULTY
The Forest Manag Felt Lichen Poten		Moderate – PHP financially contributes annually to Boreal Felt Lichen surveys. Surveys are conducted by an expert lichenologist.
	LONG-TERM STR	ATEGIC MONITORING PROGRAM
MANAGEMENT	Boreal Felt Lichen population recovery	
OBJECTIVE		
INDICATOR	Population estimates	
MONITORING/RE	PORTING	MONITORING STRATEGY
FREQUENCY Every 3 years. Baseline year is 2014.		DNR is responsible for population inventory and studying habitat use. PHP is responsible for obtaining this data.
DATA SOURCES		COST AND DIFFICULTY
NSDNR; NSDOE		Low to High - Dependent on PHP's required level of involvement
FOREST MANAGEMENT PRESCRIPTION		
- In planned harvest operations containing potential Boreal Felt Lichen habitat, a field survey is conducted to determine if Boreal Felt Lichen is present on site. If found, a 100 meter no harvest buffer is left around the site.		
2015 MONITORING UPDATE		
- In 2015, there were 13 planned harvest sites surveyed where Boreal Felt Lichen was		

identified. A 100 meter no harvest buffer was left around each lichen location.

- Since 2008, PHP has worked with the Mersey Tobeatic Research Institute to conduct Boreal Felt Lichen surveys. Prior to these surveys, there were three known locations of Boreal Felt Lichen in Nova Scotia. Since PHP and MTRI's surveys began, the number of known locations has increased to 170 in PHP's forest management area, with 123 of those locations being found in Richmond County, Cape Breton.

- NSDNR recently published a paper entitled "Forest Harvesting Impacts on Mortality of an Endangered Lichen at the Landscape and Stand Scales". This paper supports the 100 meter protection buffer around known Boreal Felt Lichen sites to maintain the micro-climate around the site. NSDNR is also currently working on a habitat supply research paper. NSDNR will also be working on improving the predicted habitat model for Boreal Felt Lichen. NSDNR will also be monitoring how different buffer widths affect microclimate using iButtons (micro-climate data loggers). Currently, there are about 24 iButtons in the field which are being used to assess the variation between and within stands. This information will help determine the sample size needed for a future buffer width study.

SUPPORTING DOCUMENTS/REFERENCES

Boreal Felt Lichen Recovery Strategy; Boreal Felt Lichen Recovery Team

HCV – VOLE EARS LICHEN OCCURRENCES

HCV ATTRIBUTE	Species at Risk – Habitat and Population	
	OPERATION	AL MONITORING PROGRAM
MANAGEMENT	Protect identified lo	ocations of Vole Ears Lichen
OBJECTIVE		
INDICATOR	Administratively protect identified locations of Vole Ears Lichen according to SMP	
MONITORING/RE	PORTING	MONITORING STRATEGY
FREQUENCY Annual		Spatial data of known vole ears lichen has been provided to PHP by the NS Department of Environment. There are no known locations of vole ears lichen in the 7 eastern counties where PHP operates.
DATA SOURCES COST AND DIFFICULTY		COST AND DIFFICULTY
The Forest Manager (TFM); Boreal Felt Lichen Potential Habitat Layer		Low – There are no known locations of Vole Ears Lichen in PHP's management area
FOREST MANAGE	MENT PRESCRIPTION	
 Due to the known locations and predictive habitat to exist outside of PHP's forest management area, there are no special management practices identified at this time for vole ears lichen. PHP is a participating member of the provincial Lichen Recovery Team, so should any change occur where special management practices are required on PHP's lands, they will be implemented. 		
2015 MONITORING UPDATE		
There are no existing or new locations of Vole Ears Lichen in PHP's forest management area.		
SUPPORTING DOCUMENTS/REFERENCES		

COSEWIC Assessment and Status Report, 2009; NSDNR; ACCDC 2016 Database

HCV – BLUE FELT LICHEN OCCURRENCES

HCV ATTRIBUTE	Species at Risk – Ha	abitat and Population	
	OPERATION	IAL MONITORING PROGRAM	
MANAGEMENT OBJECTIVE	Protect identified locations of Blue Felt Lichen		
INDICATOR	Administratively protect identified locations of Blue felt lichen according to SMP		
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Spatial data of known vole ears lichen has been provided to PHP by the NS Department of Environment. There are known locations of vole ears lichen in the 7 eastern counties where PHP operates. Of the 104 locations, 13 are located in the 7 eastern counties. Of the 13 located in the 7 eastern counties, 3 are located in PHP's forest management area and are in TFM for planning and operations.	
DATA SOURCES The Forest Manager (TFM); Boreal Felt Lichen Potential Habitat Layer		COST AND DIFFICULTY Low – A survey was conducted for the two known locations of blue felt lichen in PHP's management area.	
FOREST MANAGE	MENT PRESCRIPTION	·	
 Of the three locations known to exist on PHP's lands, one is located in a candidate protected area and is currently under a harvest moratorium. The other two locations are located in the working forest. These two locations are nearby to planned harvest operations. In the fall of 2014, PHP requested a survey of each area by an expert field lichenologist to verify if the lichen is still present. The lichen was present in both locations, so a 100 meter no harvest buffer will be maintained around each location, which is the same practice used for boreal felt lichen in Nova Scotia. 			
2015 MONITORING UPDATE			
There are no new locations of Blue Felt Lichen in PHP's forest management area.			
SUPPORTING DOG	SUPPORTING DOCUMENTS/REFERENCES		

COSEWIC Assessment and Status Report, 2009; NSDNR; ACCDC 2016 Database

HCV – EASTERN WHITE CEDAR

HCV ATTRIBUTE	Species at Risk – Ha	abitat and Population
	OPERATION	AL MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Protect identified locations of Eastern White Cedar	
INDICATOR	Protection of all kno	own locations of Eastern White Cedar
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Ensure all known locations of Eastern White Cedar in PHP's management area are protected from harvest activities.
DATA SOURCES The Forest Manager (TFM); NSDNR, NSE, ACCDC databases		COST AND DIFFICULTY Low – PHP does not conduct include the harvest of Eastern White Cedar in its management
FOREST MANAGEMENT PRESCRIPTION		
- PHP does not include the harvest of Eastern White Cedar in its forest management. Queries of the NSDNR forest inventory, as well as reviews of the rare species databases from NSDNR, NSE, and ACDCC, did not identify eastern white cedar stands for lands managed by PHP.		
2015 MONITORING UPATE		
No known stands of eastern white cedar occur within PHP's area of operation.		
SUPPORTING DOCUMENTS/REFERENCES		
A Management Plan for Native Occurrences of Eastern White Cedar in Nova Scotia, 2010		

HCV – BLACK ASH

HCV ATTRIBUTE	Species at Risk – Habitat and Population	
	OPERATION	AL MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Protect identified locations of Black Ash	
INDICATOR	Protection of all kno	own locations of Black Ash
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Ensure all known locations of Black Ash in PHP's management area are protected from harvest activities.
DATA SOURCES The Forest Manager (TFM); NSDNR, NSE, ACCDC databases		COST AND DIFFICULTY Low – PHP does not conduct include the harvest of Black Ash in its management
FOREST MANAGEMENT PRESCRIPTION - PHP does not include the harvest of Black Ash in its forest management. Queries of the NSDNR forest inventory, as well as reviews of the rare species databases from NSDNR, NSE, and ACDCC, did not identify black ash stands for lands managed by PHP.		
2015 MONITORING UPATE		
No known stands of black ash occur within PHP's area of operation.		
SUPPORTING DOCUMENTS/REFERENCES		
NSDNR, NSE, ACCDC databases		

HCV – FROSTED GLASS WHISKERS HABITAT

HCV ATTRIBUTE	Species at Risk – Ha	bitat	
	OPERATION	AL MONITORING PROGRAM	
MANAGEMENT OBJECTIVE	Maintain Frosted Glass Whiskers Habitat		
INDICATOR	Administratively protect Frosted Glass Whiskers habitat identified in NSDNR's Significant Habitat database and Atlantic Canada Conservation Data Centre database		
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor annual harvest operations to ensure Frosted Glass Whiskers habitat is administratively protected from all forest management activities.	
DATA SOURCES The Forest Manager (TFM)		COST AND DIFFICULTY Low – PHP does not conduct forest management activities within Frosted Glass Whiskers habitat	
FOREST MANAGE	FOREST MANAGEMENT PRESCRIPTION		
 PHP does not conduct forest management activities within Frosted Glass Whiskers habitat identified in NSDNR's Significant Habitat database and Atlantic Canada Conservation Data Centre database 			
2015 MONITORIN	2015 MONITORING UPDATE		
A GIS overlay using completed harvest treatment data from 2015 shows that there have been no forest management activities in identified Frosted Glass Whiskers forest stands. There have been no new locations identified for Frosted Glass Whiskers in 2015.			
SUPPORTING DOCUMENTS/REFERENCES			
Management Plan for the Frosted Glass Whiskers, Nova Scotia Population, 2011; ACCDC 2016 Database			

HCV – COLD WATER REFUGIA SUB-WATERSHEDS

HCV ATTRIBUTE	Long-term hydrolog	gic functions
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintenance of the	rmal cover for Atlantic Salmon and Brook Trout habitat
INDICATOR		50% crown closure at the stand level in cold water 12,218 hectares) with the exception of stands d firm trees.
MONITORING/RE FREQUENCY Annual	PORTING	MONITORING STRATEGY Monitor implementation of stand structure reserve using GIS overlay of completed harvest treatments with cold water refugia sub-watershed areas.
DATA SOURCES		COST AND DIFFICULTY
The Forest Manag	ger (TFM)	Low – PHP monitors this internally with resources currently available.
LONG-TERM STRATEGIC MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintenance of thermal cover for Atlantic Salmon and Brook Trout habitat	
INDICATOR	Maintenance of nor	mal long-term hydrologic functions
MONITORING/REPORTING FREQUENCY Every 3 years.		MONITORING STRATEGY PHP will explore a long-term monitoring program with partners such as NS Department of Aquaculture and Fisheries, and Fisheries and Oceans Canada, to monitor stream temperatures pre and post-harvest in a sample area of stands within cold water refugia sub-watersheds to determine effectiveness of thermal cover retention.
DATA SOURCES		COST AND DIFFICULTY
- NS Department of Aquaculture and Fisheries		Low to High – dependent on level of involvement by PHP
- Fisheries and Oceans Canada		
FOREST MANAGEMENT PRESCRIPTION		
 Cold water refugia areas are managed to maintain as much thermal cover as possible by leaving a minimum 50% crown closure wherever possible at the stand level following harvest treatments 		

- The only exception is in stands containing a high proportion of non-wind firm trees, such as balsam fir or white spruce that are vulnerable to blowdown.

- No intensive forest management will occur in these HCVF's (i.e. establishing FSC defined plantations).

2015 MONITORING UPDATE

A total of 64 hectares (0.5% of total cold water refugia area) was clearcut or overstory removal in cold water refugia areas in 2015. No other treatments were applied in cold water refugia areas. Since the stand condition was not conducive to maintaining minimum 50% crown closure, the clearcut or overstory method was applied. That is, these stands were dominated by non-wind firm trees such as fir or spruce.

SUPPORTING DOCUMENTS/REFERENCES

N/A

HCV – INTERNATIONAL BIRD AREAS

HCV ATTRIBUTE	Migratory birds habitat	
	OPERATION	AL MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Maintain and/or enhance migratory bird habitat	
INDICATOR	Reserve stand structure in Important Bird Areas (IBA's)	
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Currently, all identified IBA's in PHP's operating area are not impacted by forest management activities due to their location (i.e. off shore islands or inaccessible forest areas). Specific protocols for monitoring birds at IBAs are in development and will leverage and adapt existing monitoring programs that are directly relevant to the IBA Program (IBA Canada website). Verify annually that spatial list of IBA's is up-to-date for PHP's operating area.
DATA SOURCES The Forest Manager (TFM); IBA Canada		COST AND DIFFICULTY Low – PHP does not conduct forest management activities in IBA's, therefore, monitoring is not considered necessary.

FOREST MANAGEMENT PRESCRIPTION

- The Scaterie Island IBA site is located within the boundaries of a legally-designated protected wilderness area. No harvesting is permitted to occur within this site.

- Coastal IBA sites are not impacted by PHP's forest management activities, therefore, no special management practices are required.

- IBA sites Cape North and Central Cape Breton Highlands have been addressed in Category 1, Question 1 for Bicknell's thrush. Additionally, the Cape North IBA site contains significant concentrations of Boreal owl. For this HCVF, no harvesting currently occurs and is not expected to occur in the future. Should harvest plans be developed, a management strategy for this HCVF will be developed.

2015 MONITORING UPDATE

PHP has not conducted any forest management activities in IBA's identified within the forest management area.

SUPPORTING DOCUMENTS/REFERENCES

IBA Canada website http://www.ibacanada.ca/

HCV – RED SPRUCE

HIGH CONSERVATION VALUE – RED SPRUCE		
HCV ATTRIBUTE	Natural Red Spruce Stands	
	OPERATION	NAL MONITORING PROGRAM
MANAGEMENT OBJECTIVE	Manage red spruce stands according to PHP Work Instruction for red spruce	
INDICATOR	-	naintenance of red spruce stands to improve the ged conditions over time.
MONITORING/RE	PORTING FREQUENCY	MONITORING STRATEGY
Annual		Verify that annual harvest completions in natural red spruce stands were implemented using PHP's work instruction for red spruce management.
DATA SOURCES		COST AND DIFFICULTY
The Forest Manager (TFM)		Low – PHP has forest cover and historical data that shows natural red spruce stand locations. The PHP planner identifies these areas for management.
FOREST MANAGE	MENT PRESCRIPTION	
Red Spruce D	ominated Stands	
- Strive for tw	o to three cohort stand	structures.
- Over time, v	ve will strive to increase	the area of multiple ages in many stands.
- Promote nat	tural red spruce regener	ation
 At harvest (other than tending), trees should be large and of high value. Management (spacings, thinnings) should be carried out to help meet this objective. 		
 Retain overstory structure, including snags; future snags; other tolerant species; and residual red spruce component – both individuals when windfirm and in clumps (structure and seed). 		
Immature stands		
The option for immature red spruce stands is to partially remove the overstory in 2-3+ stages separated by a period of 10 to 20 years. The trees retained should be windfirm and quality immature trees. This helps ensure increased value of residual stand and regeneration establishment, and subsequent regeneration release(s).		

Maturing stands

As a preferred option, a modified shelterwood treatment providing increased retention will be implemented, with a plan for two ages initially, with the intent of a third as the newly regenerated stand grows into the existing overstory canopy. As possible, considering tree ages and wind firmness:

Step One: Initial shelterwood harvest is modified to include more patch retention, by doubling the present wildlife clump retention – move to 20 trees per hectare, with patches scattered throughout the treatment area (i.e. an irregular shelterwood)

Step Two (once regeneration is 60 cm tall at 5-10 yrs): Overstory harvest to release regeneration is needed (regeneration protection harvest techniques implemented). The retention includes both small patches of residuals, as well as individuals (as available, few isolated pines/hemlock/hardwoods, and snags with designated red spruce retention). Ten living trees per hectare are required. Shelterwood completed with adequate established regeneration.

Step Three (15-35 yrs): The young and immature stand is tended as it grows (space – thin).

Longer term: As trees grow into the upper canopy, some of the patches and individual trees will be harvested, excluding designated wildlife clumps and legacy trees.

At this time (in the future), three cohorts are introduced in to the stand with the intent of patterning an uneven-aged structure.

In some instances, trees in the forest stand planned for treatment are not wind-firm and excessive blowdown and significant wood losses would occur following implementing one of the treatments described above. If the stand is determined to be a high risk for blowdown, an alternative treatment may be implemented (eg. strip cuts), or it should be left to grow until maturity then harvested.

2015 MONITORING UPDATE

A GIS overlay using completed harvest treatment data from 2015 shows that approximately 67 hectares of natural red spruce stands were managed using PHP's work instruction for red spruce management.

SUPPORTING DOCUMENTS/REFERENCES

N/A

HCV – PROTECTED AREAS

MANAGEMENT	OPERATION			
MANAGEMENT		AL MONITOR	ING PROGRAM	
	Establish protected	areas (legal, p	pending, and/or a	administrative) in PHP's
OBJECTIVE	management area			
INDICATOR	Establishment of leg	al, pending, a	and/or administra	ative protected areas
MONITORING/REPORTING		MONITORI	MONITORING STRATEGY	
FREQUENCY		Continue to	o monitor provin	cial government's
Annual			•	the establishment and
		legal prote	ction of new wild	lerness areas and/or
		other decis	ions made regar	ding areas.
DATA SOURCES		COST AND	DIFFICULTY	
The Forest Manage	er (TFM); NSDNR;	Low		
NSDOE				
FOREST MANAGE	MENT PRESCRIPTION			
and are clearly mar	ked as administrative	protected ar	eas.	een delineated in TFM to occur in these areas.
Protected Area Ca	tegory		# of Sites	Total Hectares
New Provincial Prov	tected Area (pending leg	gal status)	89	98,184
Provincial Parks and			21	1,492
Provincial Nature R			7	1,868
Provincial Wilderne			19	106,526
National Migratory National Parks	Bird Sanctuaries		<u> </u>	<u> </u>
Trational Tarks	ТОТА	L HECTARE		303,332
	1011		5	000,002
Administratively P	Protected Area Categor	'y	# of Sites	Total Hectares
Old Forest Areas			N/A	84,717
PHP Protected Area			8	6,147
IBP Sites & Sites of Ecological Significance		e	12	3,107
IBP Sites & Sites of	TOTAL HECTARES			
IBP Sites & Sites of	ТОТА	L HECTARE	S	93,971
IBP Sites & Sites of	ΤΟΤΑ	L HECTARE	S	93,971

2015 MONITORING UPDATE

A GIS overlay using completed harvest treatment data from 2015 shows that there has been no forest management activities conducted in legal, pending, or administrative protected areas.

In December 2015, the provincial government completed its designation of a new suite of protected areas throughout the province. In total, over 14,000 hectares have been designated as new protected areas, which has led to 12.26% of total land protection in Nova Scotia.

SUPPORTING DOCUMENTS/REFERENCES

Our Parks and Protected Areas: A Plan for Nova Scotia, 2013; TFM Data; NSE Protected Areas

HCV – SPECIAL MANAGEMENT ZONE ADJACENT TO PROTECTED AREA BOUNDARIES

HCV Limit Protected Area Access ATTRIBUTE				
	OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Minimize road construction			
INDICATOR		ruction to reduce access points into protected areas 200 meter wide special management zone.		
		MONITORING STRATEGY Assess whether new roads have been built in the special management zone using GIS overlay.		
DATA SOURCES The Forest Manager (TFM)		COST AND DIFFICULTY Low – PHP currently monitors the special		
management zone and road construction using TFM. FOREST MANAGEMENT PRESCRIPTION				
- Minimize road construction to reduce access points into the protected area. If roads are needed, they are to build parallel to the protected area boundary to minimize access points.				
2015 MONITORING UPDATE				

A GIS overlay using completed road construction data from 2015 shows that there have been no new roads built in the special management zone adjacent to protected area boundaries.

SUPPORTING DOCUMENTS/REFERENCES

N/A

HCV ATTRIBUTE				
	OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	To maintain biodiversity values and intactness in large landscape level forests			
INDICATOR	Manage large landscape level forests with special practices in protected areas, core roadless areas, and special management areas			
		MONITORING STRATEGY Assess management activities within large landscape level forests to ensure practices comply with		
		requirements outlined for protected areas, core roadless areas, and special management areas.		
DATA SOURCES		COST AND DIFFICULTY		
The Forest Manag	ger (TFM)	Low – PHP currently monitors large landscape level forests using TFM.		
 No new roads in Core Roadless areas For HCV area outside core roadless, follow road design objectives as shown below. Road Index value at HCV level not to exceed 0.58 km/km2. If feasible and where necessary, block off access to reduce road travel. 				
N- close between resource use				
prescriptions t	 Use the provincial Forest Ecosystem Classification Guide to identify ecosite level prescriptions that: o Promote ecosite patches by combining stands through treatment 			
 Employ 'extensive' management practices that support: 				

HCV – Large Landscape Level Forests

- natural regeneration

- longer rotations with consideration of natural disturbance processes

- tree species diversity consistent with the vegetation type, while promoting those that support long-term resilience (i.e. best options for future)

- No full-tree logging

- Reduce road length by increasing average forwarding distance targets by 20% (from 250 m to 300 m)

- Bridge construction may be temporary and removed as practical

- Retain minimum 60% area in non-clearcut condition (at the HCV level). Non-clearcut defined as forest stand greater than 10 years of age.

- No FSC plantations / Intensive management

- No planting of exotic species

- Acadian Forest Restoration (considering N.S. Forest Code; FSC)

- Management will align with natural disturbance regimes

- Application of Forest Ecosystem Classification to identify appropriate treatments

- Appropriate forest covertype management: Use of hardwood management keys

- Appropriate forest covertype management: Use of mixedwood management keys

- Natural regeneration where appropriate

- Appropriate use of PHP's 12 different harvest techniques (CC, PC, SW, ST, Single, Group, Patch, CT, OR, CTR, RS, SC)

- Species at Risk Recovery Strategy/SMP Implementation

- No herbicides

- Steep Slope Exclusion

- Leave patches (e.g. active eagle/hawk nest sites, inoperable areas, vernal pools, DNR requests during approval process)

2015 MONITORING UPDATE

The below table summarizes the current status of each large landscape level forest with respect to road index and minimum non-clearcut condition. Non-clearcut condition is defined as anything greater than 10 years of age. Roads built in 2015 occurred only in the Isaacs Harbour River HCVF, Salmon Gaspereaux HCVF and Boisedale Hills HCVF, and they were all built outside the core roadless area.

			mum Allowed = 0.58 n/km2	Minimum Non- clearcut Condition = 60%
HCVF LLLF Name	Total HA	2014 Road Index	Future Road Index	2015 Non-clearcut Condition
	-			
Barren Hill	1,318	0.08 km/km2	0.20 km/km2	88%
Boisdale Hills Bornish Hill	5,630	0.40 km/km2	0.52 km/km2	98%
(fully protected)	2,106	0 km/km2	0 km/km2	100%
Country Harbour	8,202	0.03 km/km2	0.03 km/km2	100%
East Bay Hills	1,865	0.23 km/km2	0.31 km/km2	78%
French River	25,226	0 km/km2	0 km/km2	100%
Hill Lake	877	0.55 km/km2	0.65 km/km2	93%
Ingonish River	15,210	0.01 km/km2	0.01 km/km2	100%
Isaacs Harbour River	6,157	0.25 km/km2	0.42 km/km2	97%
Jim Campbells Barren (fully protected)	4,586	0.21 km/km2	0.21 km/km2	100%
Masons Mountain (fully protected)	1,022	0.06 km/km2	0.06 km/km2	100%
North River	6,328	0.20 km/km2	0.20 km/km2	100%
Oban	1,618	0.57 km/km2	0.78 km/km2	92%
Petit Lake Ruiss Noir (fully protected)	1,612	0 km/km2	0 km/km2	100%
Salmon Gaspereaux	2,357	0.30 km/km2	0.61 km/km2	93%
Upper Liscomb River	7,398	0.07 km/km2	0.07 km/km2	100%
TOTAL HECTARES	91,512		Future index may e 0.58 km/km2. Will road index to meet	need to manage

HCVF LLLF	Total	2015 Total Area	
Name	HA	Treated	Treatment Used

Barren Hill	1,318	No area treated	
Boisdale Hills	5,630	159 hectares	Partial Cut/Planting
	3,030	139 Hectales	Cut/Flanting
Bornish Hill (fully protected)	2,106	No area treated	
Country			
Harbour	8,202	No area treated	
East Bay Hills	1,865	34 hectares	Clearcut/Planting
French River	25,226	No area treated	
Hill Lake	877	28 hectares	Weeding
Ingonish River	15,210	No area treated	
Isaacs Harbour			
River	6,157	No area treated	
Jim Campbells			
Barren (fully			
protected)	4,586	No area treated	
Masons			
Mountain (fully			
protected)	1,022	No area treated	
North River	6,328	No area treated	
	,		
Oban	1,618	No area treated	
Petit Lake Ruiss			
Noir (fully			
protected)	1,612	No area treated	
Salmon			
Gaspereaux	2,357	26 hectares	Clearcut
Upper Liscomb			
River	7,398	No area treated	
TOTAL			
HECTARES	91,512		

HCVF Category 3 – Rare, Threatened or Endangered Ecosystems

		-	
HCV ATTRIBUTE			
	OPERATION	AL MONITORING PROGRAM	
MANAGEMENT OBJECTIVE	Maintain rare, threatened or endangered ecosystems		
INDICATOR	Rare, threatened or endangered ecosystems administratively protected from forest management activities		
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor rare, threatened or endangered ecosystems to ensure they are administratively protected from forest management activities. Exception applies if the mapped ecosystem type does not match on-the- ground characteristics.	
DATA SOURCES The Forest Manager (TFM); NSDOE		COST AND DIFFICULTY Low	
FOREST MANAGE			

HCV – Significant Ecosites

- All significant ecosites are administratively protected from forest management activities with the following exceptions:

- Karst conifer forest, karst hardwood forest, calcareous forest, and hemlock forest that have been previously managed will continue to be managed to maintain and restore mature climax conditions.

- Significant ecosites are identified using the provincial forest inventory data and there has been limited field verification, so there is a certain amount of ambiguity within the dataset. Since there may be data inaccuracies between the digital information versus on-the-ground characteristics, stands that do not match the inventory data are exempt from special management activities as outlined here.

2016 MONITORING UPDATE

A GIS overlay using completed harvest treatment data from 2015 and significant ecosite data shows that there was 97 hectares managed in Coastal Barrens. The significant ecosite database shows these stands to be Coastal barrens, however, they were predominately balsam fir/black spruce stands. Additionally, 43 hectares were managed in Inland Barrens, where balsam fir/black spruce stands were present and not barrens. Therefore, no special management practices or protection was implemented.

SUPPORTING DOCUMENTS/REFERENCES

Significant Ecosite data layer, NSDOE

HCV – SIGNIFICANT, OID OR UNIQUE FORESTS

HCV ATTRIBUTE	Rare, threatened or endangered ecosystems				
	OPERATIONAL MONITORING PROGRAM				
MANAGEMENT OBJECTIVE	Maintain rare, threatened or endangered ecosystems				
INDICATOR	Rare, threatened or endangered ecosystems administratively protected from forest management activities				
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor rare, threatened or endangered ecosystems to ensure they are administratively protected from forest management activities. Exception applies if the mapped ecosystem type does not match on-the-ground characteristics.			
DATA SOURCES The Forest Manager (TFM); NSDOE FOREST MANAGEMENT PRESCRIPTION		COST AND DIFFICULTY Low			

- All significant, old or unique forests (SOUF) are administratively protected from forest management activities that meet the following species composition. Exception applies if the mapped ecosystem type does not match on-the-ground characteristics.

Species composition	Stand height
70% or more spruce or red spruce	≥17m
50% or more eastern hemlock	≥15m
50% or more white pine	≥18m
70% or more climax coniferous species with the most common species	≥17m
50% or more sugar maple	≥17m
50% or more yellow birch	≥17m
70% or more climax deciduous species (tolerant hardwood)	≥17m
70% or more climax coniferous or deciduous species with neither group	≥17m
30% or more red pine	≥12m
10% or more red oak	Any height
10% or more eastern white cedar	Any height

2015 MONITORING UPDATE

A GIS overlay using completed harvest treatment data from 2015 shows that there were no SOUF stands managed.

SUPPORTING DOCUMENTS/REFERENCES

Significant, old or unique data layer, NSDOE

HCV – OLD FOREST

HCV ATTRIBUTE	Old Forest Protected Area		
	OPERATION	AL MONITORING PROGRAM	
MANAGEMENT OBJECTIVE	Establish old forest protected areas on land-base		
INDICATOR	Establishment and legal protection of old forest protected areas		
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor old forest protected areas TFM to ensure no forest management activities are conducted.	
DATA SOURCES The Forest Manager (TFM); NSDNR; NSDOE		COST AND DIFFICULTY Low	
- All identified old forest areas are legally protected by the provincial government PHP staff is aware that no forest management activities are allowed to occur in these areas			

- PHP staff is aware that no forest management activities are allowed to occur in these areas.

2015 MONITORING UPDATE

A GIS overlay using completed harvest treatment data from 2015 shows that there have been no forest management activities conducted in the old forest areas identified by the provincial government.

SUPPORTING DOCUMENTS/REFERENCES

Old forest GIS layer, NSDNR

HCV – POORLY REPRESENTED ECOSYSTEMS

HCV ATTRIBUTE					
	OPERATIONAL MONITORING PROGRAM				
MANAGEMENT OBJECTIVE	Establish protection of poorly represented ecosystems on land-base				
INDICATOR	Establishment and administrative protection of poorly represented ecosystems				
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor poorly represented ecosystems in TFM to ensure no forest management activities are conducted.			
DATA SOURCES The Forest Manager (TFM)		COST AND DIFFICULTY Low			
FOREST MANAGEMENT PRESCRIPTION					
 All identified poorly represented ecosystems are administratively protected by PHP. PHP staff is aware that no forest management activities are allowed to occur in these areas. 					

Poorly Represented Ecosystem	Total Hectares
Masons Mountain	197
Jim Cambells Barren	2,844
Boisdale Hills	1,727
Country Harbour	829
North River	27
Oban	170
Hill Lake	113
Salmon Gaspereaux	240

TOTAL HECTARES 6,147

2015 MONITORING UPDATE

There have been no management activities in the above PHP administratively protected areas.

SUPPORTING DOCUMENTS/REFERENCES

PHP Gap Analysis Report


HCV ATTRIBUTE	Continuous Canopy cover			
	OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Maintain continuous canopy cover between protected areas and old forest areas			
INDICATOR	Maintain a 100 meter wide continuous canopy cover (minimum 30%) corridor within the 500 meter wide Connectivity Management Zone (CMZ)			
MONITORING/RE	PORTING FREQUENCY	MONITORING STRATEGY		
Annual		Monitor 100 meters within the CMZ to ensure a continuous canopy cover and CMZ's are not severed across their width.		
DATA SOURCES		COST AND DIFFICULTY		
The Forest Manager (TFM)		Low		
FOREST MANAGEMENT PRESCRIPTION				

- The Connectivity Management Zones will be managed to provide continuous canopy cover (minimum 30%) within the 500 meter wide corridors, which will include a solid 100 meter wide core zone. Although harvesting can occur within the CMZ's, these corridors will not be severed across their width.

- The 500 meter wide CMZ's are static on the landscape, but the 100 meter wide core zone can 'move' within the CMZ.

2015 MONITORING UPDATE

The Connectivity Management Zones continue to maintain a continuous canopy cover within the 100 meter wide core zone.



SUPPORTING DOCUMENTS/REFERENCES

The Forest Manager

HCV – MARGAREE & ST. MARY'S RIVER WATERSHED

HIGH CONSERVATION VALUE – MARGAREE & ST. MARY'S RIVER WATERSHED			
HCV ATTRIBUTE	Non-clearcut Condition		
OPERATIONAL MONITORING PROGRAM			
MANAGEMENTTo maintain a high level of non-clearcut condition in St. Mary's and Margaree Watersheds, and restoration management			
INDICATOR	Each watershed shall have minimum 80% of its area (that is managed by PHP) in a non-clearcut condition, and 90% of each watershed shall be managed for restoration (i.e. no more than 10% of each watershed will be established as a FSC plantation).		
Annual Ma		MONITORING STRATEGY Monitor non-clearcut condition in each watershed to ensure target of minimum 80% is met.	
DATA SOURCES The Forest Manager (TFM)		COST AND DIFFICULTY Low	
FOREST MANAGEMENT PRESCRIPTION			

- The St. Mary's and Margaree watersheds will be managed to maintain 80% or more of all lands managed by PHP in the watershed in a closed forest condition (> 12 years of age).

- Additionally, PHP will maintain at least 90% of the St. Mary's and Margaree watersheds in a natural condition for restoration, and will establish 200 m forest restoration zones (i.e. non-intensive management) along all main watercourses.

2015 MONITORING UPDATE

PHP has been monitoring the non-clearcut condition in these watersheds for several years. The below graph shows that since 2008 each watershed has maintained 80% or more of its forest area under management by PHP in a non-clearcut condition. Additionally, PHP has not yet identified areas on the land base that will be established as an FSC defined plantation (up to 10% of the total forest lands), therefore, all forest areas are currently being managed for restoration and/or maintenance of existing Acadian forest characteristics.

See Indicator 3.2 for current condition of St. Mary's and Margaree watersheds.

SUPPORTING DOCUMENTS/REFERENCES

The Forest Manager

HCVF Category 4 – Basic Services of Nature

HCV ATTRIBUTE	Water Health	
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintain water health for communities	
INDICATOR	Implement water protection measures in legally protected municipal water supply areas	
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor implementation of water protection measures.
DATA SOURCES		COST AND DIFFICULTY
The Forest Manager (TFM); NSDOE Low FOREST MANAGEMENT PRESCRIPTION		

HCV – Legally Protected Municipal Water Supply Areas

- There is no land managed by PHP within the water supply areas, since they are legally protected and therefore excluded from forest management activities.

2015 MONITORING UPDATE

A GIS overlay using completed harvest treatment data from 2015 shows that there have been no forest management activities within the water supply areas.

SUPPORTING DOCUMENTS/REFERENCES

Nova Scotia Department of Environment

HCV – WATER SUPPLY INTAKE AREAS

HCV ATTRIBUTE	Water Health	
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintain water health for communities	
INDICATOR	Implement water protection measures around water supply intake areas.	
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor implementation of water protection measures.
DATA SOURCES		COST AND DIFFICULTY
The Forest Manager (TFM); NSDOE		Low
FOREST MANAGEMENT PRESCRIPTION		
- Water protection measures include the Wildlife and Habitat Watercourse Protection Regulations.		

- Water protection measures include the Wildlife and Habitat Watercourse Protection Regulations, monitoring of % closed forest condition, steep slope management, HCVF aquatic watershed management, and rutting and ground disturbance guidelines.

2015 MONITORING UPDATE

A GIS overlay of completed harvest treatments and water supply intake areas shows no hectares were managed within the intake areas.

SUPPORTING DOCUMENTS/REFERENCES

Nova Scotia Department of Environment

HCV – STEEP SLOPES

HCV ATTRIBUTE	Soil Health; Community Health		
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT Maintain soil health and community health		th and community health	
OBJECTIVE INDICATOR	No conventional harvesting in steep slope areas (30% average slope or greater)		
MONITORING/REPORTING FREQUENCY		MONITORING STRATEGY Monitor steep slope areas and conventional harvesting activities.	
Annual DATA SOURCES The Forest Manager (TFM)		COST AND DIFFICULTY	
FOREST MANAGEMENT PRESCRIPTION			

- Conventional harvesting is not permitted in areas with 30% average slope or greater. Nonconventional harvesting such as cable logging is permitted, however, PHP is currently not using this practice.

2015 MONITORING UPDATE

A GIS overlay using completed harvest treatment data from 2015 shows that there was 16 hectares of forest management activities within steep slope areas.

NOTE: This indicator is based on spatial data that identifies slopes > than 30% average using contour data. It is not based on the actual % slope for any given area as could be determined on-the-ground. Therefore, to calculate the results for the indicator, a GIS exercise is done which overlaps the steep slope data with completed harvest jobs to determine non-conformances. Most often, the areas showing as harvested are slivers due to inaccuracies in the data.

SUPPORTING DOCUMENTS/REFERENCES

The Forest Manager

HCVF Category 5 – Basic Needs of Local Communities

HCV – Cattle Grazing on Cape Breton Highlands

HCV ATTRIBUTE	Local commu	nities	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT	Support needs of local communities		
OBJECTIVE			
INDICATOR	Cattle grazing	on the Cape Breton Highlands is allowed	
MONITORING/REPORT	ING	MONITORING STRATEGY	
FREQUENCY Annual		Monitor any issues arising from cattle grazing on Cape Breton Highlands	
DATA SOURCES		COST AND DIFFICULTY	
N/A		Low	
FOREST MANAGEMEN	FOREST MANAGEMENT PRESCRIPTION		
 Local farmers have let their cattle graze on the Cape Breton Highlands for several years during the summer/fall months. PHP does not restrict this use. 			
2015 MONITORING UP	2015 MONITORING UPDATE		
No issues have arisen in 2015 regarding cattle grazing in the Cape Breton Highlands.			
SUPPORTING DOCUMENTS/REFERENCES			
N/A			

HCV – VIEWSHED AREAS

HCV ATTRIBUTE	Local Communities	
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Minimize visual impacts to local communities from harvest activities	
INDICATOR	Implement work instruction 'Harvest View from Roadside'	
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor any issues in identified viewshed areas arising from harvest activities.
DATA SOURCES The Forest Manager (TFM); Harvest View from Roadside Work Instruction		COST AND DIFFICULTY Low
FOREST MANAGEMENT PRESCRIPTION		

- District staff is responsible for determining the visibility rating using the TFM layer 'Viewsheds'. A harvest area determined to be within the **low category** will not require any specific landscape planning beyond regular housekeeping measures and removal of unsightly damaged residual trees. - Cut blocks falling in **the medium category** on the visibility grid should be designed using the "Landscape Level" instructions in the Harvest View from Roadside Work Instruction. Blocks falling into the high visibility category will follow the "Landscape level", "Stand level" and "Road design" practices as applicable.

2015 MONITORING UPDATE

Below is a list of hectares treated within the Viewshed area. Forest stands with a visability rating of 'low' do not require any specific landscape planning beyond regular housekeeping measures. Forest stands rated as moderate or high were managed by implementing the 'Harvest View from Roadside' Work Instruction. These areas are automatically flagged in TFM during planning and are included in forest management plans that are provided operations staff.

Visibility by Treatment	Hectares Treated	
High		10
PARTIALCUT		10
Low		360
		200
CLEARCUT		110

SINGLE	9
THINNING	9
Moderate	446
CLEARCUT	178
OVERSTORY	21
PARTIALCUT	151
SHELTER	5
SINGLE	71
THINNING	21
SUPPORTING DOCUME	INTS/REFEREN

Viewshed layer in TFM; Harvest View from Roadside Work Instruction

HCVF Category 6 – Traditional Cultural Identity

HCV ATTRIBUTE	First Nations Forest Values and Uses		
	OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Minimize impacts to First Nations Forest Values and Uses		
INDICATOR	Implement work instruction 'Aboriginal Value – Suspending Operations'		
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor any issues identified during annual review of operations plans with First Nations communities. PHP also maintains a public inquiry database, which captures concerns or questions the general public may have regarding planned operations.	
DATA SOURCES The Forest Manager (TFM); Public Inquiry Database		COST AND DIFFICULTY Low	
FOREST MANAGEMENT PRESCRIPTION			

HCV – Forest Values and Uses

- If operations plans are known to affect First Nations forest values or uses through a review of annual operating plans or public inquiries, PHP will suspend all activities until a resolution is found.

2015 MONITORING UPDATE

- A review of PHP's public inquiry database shows that there was a request for firewood from Chief Marshall at Potlotek First Nation. This request was granted.

- There were no other public inquiries related to PHP's operating plans that may affect First Nations.

SUPPORTING DOCUMENTS/REFERENCES

PHP Work Instruction 'Aboriginal Values – Suspending Operations'

HCV – TRADITIONAL CULTURAL IDENTITY

HCV ATTRIBUTE	First Nations Traditional Cultural Identity		
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Minimize impacts to First Nations Traditional Cultural Identity		
INDICATOR	Successful implementation of Impact Benefit Agreement and Environmental Agreement with The Assembly of Nova Scotia Mi'kmaq Chiefs		
MONITORING/REPORTING FREQUENCY Annual		MONITORING STRATEGY Monitor successful completion and implementation of Impact Benefit Agreement and Environmental Agreement with The Assembly of Nova Scotia Mi'kmaq Chiefs	
DATA SOURCES The Forest Manager (TFM); Public		COST AND DIFFICULTY Moderate	
Inquiry Database FOREST MANAGEMENT PRESCRIPTION			
Once the Impact Benefit Agreement and Environmental Agreement are finalized, forest management prescriptions will be developed in collaboration with Nova Scotia Mi'kmaq.			

2015 MONITORING UPDATE

- PHP is currently working with the The Assembly of Nova Scotia Mi'kmaq Chiefs to finalize an Impact Benefits Agreement and Environmental Agreement, which will include provisions for the protection of Mi'kmaq Aboriginal and Treaty Rights and Archaeological and Cultural Resources. This work is currently on-hold in 2016 as other issues get resolved.