

2022



**ANNUAL MONITORING REPORT**  
**A Public Report on Safety, Environmental and**  
**Responsible Forest Management Progress**



**WOODLANDS**  
**PORT HAWKESBURY PAPER**

## Executive Summary



The 2022 Annual Monitoring Report provides a summary of Port Hawkesbury Paper’s safety, environmental, and forest management progress in the Woodlands Unit. Since 2002, Port Hawkesbury Paper (PHP) has been monitoring and reporting on a suite of sustainable forest management indicators to measure its progress towards achieving targets regarding social, economic, environmental, and cultural forest values. Long-term monitoring of these values allows the public to better understand PHP’s forest management activities, and the goals and objectives we set to ensure our forest management is having a positive impact and to implement action items in areas that we are not. This is an important element of continual improvement, which PHP strives for every day.

This report also summarizes the effectiveness monitoring program for High Conservation Value Forests (HCVF). These values were first identified in 2008 for Forest Stewardship Council® (FSC®) certification and updated in January 2018 to include new knowledge and information related to species at risk and protected areas. Annual monitoring is conducted to assess the effectiveness of the measures used to maintain or enhance the identified values.



The mark of  
responsible forestry



## Contents

Executive Summary .....	1
Contents .....	2
Key Commitments to Safety .....	7
Key Commitments to the Environment.....	8
Summary of SFM Indicators .....	40
FSC Canada Specific Monitoring Updates .....	42
High Conservation Value Forest .....	47
Effectiveness Monitoring Program.....	47
HCV – American Marten Habitat .....	49
HCV – Mainland Moose Habitat .....	52
HCV – Canada Lynx Habitat .....	55
HCV – Wood Turtle Habitat .....	58
HCV – Bicknell’s Thrush Habitat .....	61
HCV – Rusty Blackbird Habitat.....	65
HCV – Roseate Tern Habitat .....	67
HCV – Olive-Sided Flycatcher Habitat .....	69
HCV – Eastern Whip-Poor-Will Habitat .....	71
HCV – Eastern Wood Peewee Habitat.....	73
HCV – Canada Warbler Habitat .....	76
HCV – CHIMNEY SWIFT Habitat.....	79
HCV – COMMON NIGHTHAWK Habitat.....	81
HCV – WOOD THRUSH Habitat.....	84
HCV – EVENING GROSBEAK Habitat .....	86
HCV – BLACK-FOAM LICHEN Habitat .....	88
HCV – LITTLE BROWN MYOTIS Habitat.....	89
HCV – TRI-COLORED BAT Habitat .....	91

HCV – NORTHERN MYOTIS Habitat .....	92
HCV – New Jersey Rush Habitat .....	94
HCV – Boreal Felt Lichen Occurrences .....	95
HCV – Vole Ears Lichen Occurrences .....	98
HCV – Blue Felt Lichen Occurrences.....	100
HCV – Eastern White Cedar .....	101
HCV – Black Ash.....	102
HCV – Frosted Glass Whiskers Habitat .....	104
HCV – Wrinkled Shingle Lichen Habitat.....	105
HCV – Cold Water Refugia Sub-watersheds .....	107
HCV – International Bird Areas.....	108
HCV – Red Spruce .....	110
HCV – Protected Areas .....	112
HCV – Special Management Zone Adjacent to Protected Area Boundaries .....	113
HCV – Intact Forest Landscapes .....	115
HCV – Significant, Old or Unique Forests .....	123
HCV – Old Forest.....	124
HCV – Poorly Represented Ecosystems.....	125
HCV – Connectivity Management Zones.....	126
HCV – Margaree & St. Mary’s River Watershed.....	127
HCV – Water Supply Intake Areas .....	129
HCV – Steep Slopes.....	129
HCV – Viewshed Areas .....	131
HCV – Traditional Cultural Identity.....	134

## About the Woodlands Unit



2001: First forest company in Canada to achieve certification to both the Canadian Standards Association (CSA®) and Sustainable Forestry Initiative (SFI®) standards for sustainable forest management.

2008: First forest company in Maritime provinces to achieve certification to the Forest Stewardship Council® (FSC®) Maritime Standard for responsible forest management. PHP is currently certified to the FSC Canada National Standard.

2008: First forest company in Maritime provinces to achieve Chain-of-Custody certification to the FSC® standard for wood traceability.

2014: Woodlands becomes re-certified to the SFI® Forest Management and Chain-of-Custody Standards, and the PEFC™ Chain-of-Custody standard.

With a dedicated staff of 23 people, the company's Woodlands Unit currently manages 30% of the Crown land in Nova Scotia, which represents 58% (roughly 520,000 hectares) of the Crown land in the seven eastern counties. As a result of 50+ years of silviculture activities on these lands, the forest continues to see an increase in economic activity within a healthy and naturally diverse ecosystem.

Our wood supply primarily comes from the seven eastern counties of Nova Scotia with additional wood purchased in central and western Nova Scotia. The Woodlands Unit also provides silviculture services and information on sustainable forest management practices to private woodland owners. Additionally, we provide training on best management practices for Crown and private contractors and operators. In addition to acquiring softwood pulpwood from the managed forest, Port Hawkesbury Paper also manages its forest lands to produce softwood and hardwood logs, and other products, for sale to local sawmills and buyers.

As the largest Crown license holder in eastern Nova Scotia, we believe that good business includes strong Indigenous and non-Indigenous community support and involvement,

environmental awareness, continued growth in forest management and contribution to the Nova Scotia economy. The public use of Crown lands for rights holders, recreation, accessibility, hunting and fishing illustrates the wide variety of values held by the general public. To achieve sustainable forest management, the needs of all rights holders and stakeholders must be assessed and managed appropriately.

Forest management certification is one of many tools to support the sustainability of Port Hawkesbury Paper. Certification is a voluntary process by which planning, procedures, systems and performance of on-the-ground forestry operations are audited by a qualified and independent third party against a predetermined standard. Forest operations found to be in conformance with the given standard are issued a certificate. Port Hawkesbury Paper supports the mutual recognition of credible forest certification systems that take into account national and regional characteristics such as natural conditions, forest ownership structures and legislation.


Forest operations at Port Hawkesbury Paper are carefully planned to deliver a valuable, sustainable resource that satisfies economic, social, and environmental benefits. More specifically, Port Hawkesbury Paper works diligently to ensure sustainable harvests, increased forest productivity, and protection for wildlife, water, and recreational resources. With the use of high-end computer mapping and software systems, we supervise our operations to strict standards to ensure we continually meet or exceed our expectations for a healthy productive forest for the future.

#### **Port Hawkesbury Paper's Crown Land Forest Management Area**

PHP's Defined Forest Area (DFA) is located in the seven eastern counties of Nova Scotia. The geographic extent of the DFA is shown in Figure 1. The company manages approximately 520,000 hectares of Crown lands through a license agreement with the provincial government within the DFA.

In addition to acquiring wood from PHP company managed lands, the company harvests wood from private woodland owners through short-term stumpage leases. Private wood is also procured from private suppliers that operate on private woodlands located in central and eastern Nova Scotia. Private wood is purchased at roadside and the company provides competitive pricing. In addition, the company provides silviculture services and training in sustainable forest management practices to encourage good stewardship practices.

The public use of Crown lands for First Nations, recreation, accessibility, hunting and fishing, to name a few, illustrates the wide variety of values held by the general public. Tourism plays an important role in the regional economy; as a result, unique challenges in meeting the needs of all stakeholders must be assessed and managed appropriately. The Nova Scotia Department of Natural Resources & Renewables (NSDNRR) has implemented an integrated resource management (IRM) land use approach for the management of Crown lands.



#### **PHP's Crown Land-base as per the Forest Utilization License Agreement**

The Crown land-base was assembled using photo-interpreted forest inventory flown in 2008 and 2009 as a base. Historic treatment GIS data were incorporated from PHP and government databases to update the spatial boundaries and attributes of the forest inventory. Wildlife habitat, ecosystem data, special management layers, and hydrology and roads layers were compared, agreed upon and amalgamated where appropriate to create the most recent and accurate dataset possible.

As land-base layers are overlaid, attributes are coded to allow for partitioning of results based on forest and non-forest values. The total land area includes all area, crown wilderness area and non-forested land are removed to create the forested land-base. After removing permanent exclusions (off limits to forest management prescriptions such as protected areas, old growth areas, species at risk habitat buffers, and other land-use restrictions), the remainder is the working land-base which contributes to wood supply. The working land-base is largely occupied by special management lands, which dictate treatment prescription details.

## Key Commitments to Safety



Safety is a vital aspect of our operating philosophy. From production to quality assurance, cost control and environmental compliance, we focus on safety in everything we do.



The Health & Safety of employees takes precedence over all other responsibilities and activities within our Company. This is the cornerstone of our safety policy.




We believe that all accidents are preventable. Our success is measured by our safety performance relative to our goal of zero recordable injuries.



It is our objective to work toward continual improvement in health, safety and wellness aspects of our operations.

### Woodlands Safety Results

The Woodlands Safety Results show a strong trend towards zero lost time accidents and medical aids. We continue to promote employee and worker safety through effective training programs, monitoring, and communication to strive towards our objective of zero safety incidents year after year.



A Lost Time Accident occurs when an employee or worker is injured on the job which results in lost work time. There were two lost-time accidents in 2022.

A Medical Aid occurs when an employee or worker is injured on the job and requires medical aid but did not result in lost work time. No medical aids occurred in 2022.



## Key Commitments to the Environment



The graphic consists of four vertical panels, each with a circular icon at the top and text below. The panels are colored green, yellow, orange, and green from left to right.

- Panel 1 (Green):** Icon shows a recycling symbol with a globe in the center. Text: "It is the policy of Port Hawkesbury Paper to carry out operations in ways that do not endanger the environment. Sustaining a healthy environment is an integral part of all company operations."
- Panel 2 (Yellow):** Icon shows a view looking up at tall trees against a blue sky. Text: "PHP commits to continual improvement of all aspects of our sustainable forest management system for company-managed lands through experience and forest research."
- Panel 3 (Orange):** Icon shows a landscape with colorful autumn trees and mountains. Text: "Utilize long-term landscape ecosystem planning, appropriate silviculture systems, and operating practices that conserve biodiversity in managing our forest areas."
- Panel 4 (Green):** Icon shows a stack of papers with wooden blocks labeled "RULES" and "OBLIGATIONS". Text: "Meet or be better than all applicable regulations, legal obligations and other requirements to which Port Hawkesbury Paper subscribes."

Monitoring and reporting on Woodlands environmental performance is an integral part of achieving responsible forest management across the working landscape. Harvest contractors working on Crown land on behalf of the company are audited by PHP three times a year - Winter, Spring/Summer and Fall. Compliance and performance are checked against a range of items related to layout compliance, operational safety and environmental compliance, and job quality. Where deficiencies are found, an incident or violation report is written. Refresher training is provided by PHP staff if needed, and follow-up actions to address issue are completed by contractor.

In 2022, internal audits on Crown contractors were in transition from using PHP's audit checklist to DNRR's inspection checklist. A total of 12 audits were completed using PHP's audit checklist and 18 audits were completed using DNRR's inspection checklist.

With a total of 30 audits completed, Crown harvest contractors continue to achieve a high level of overall compliance and performance as shown in the 2022 results. This is due in large part to the long-standing working relationship between the company and its Crown land harvest

contractors. Working together to monitor performance, share information, and strive to continually improve has resulted in strong on-the-ground results.

PHP also audits its private suppliers. A total of 12 suppliers were audited in 2022. Using the previous quarter's deliveries, wood suppliers are randomly chosen to be audited by a PHP Area Supervisor. Private supplier audits are performed on active jobs when possible. However, auditing a completed job may be necessary with smaller suppliers.

Below are summaries of PHP's Crown and private supplier audit program for 2022. Areas of deficiencies are highlighted in blue. If the deficiencies are consistently on-going or deemed to be of significant concern, communications and/or training is provided to suppliers to improve performance.

## **Crown Contractor Audit Results – 2022 (PHP audit checklist)**

<b>LAYOUT COMPLIANCE (100% Required to Pass)</b>	
Worked within Cutting/Property Boundaries	100%
Wildlife clumps (as per instructions)	100%
Wildlife corridors (50M min)	100%
Properly buffered watercourses and wetlands	100%
<b>OPERATION SAFETY/ENVIRONMENTAL COMPLIANCE (90% Required to Pass)</b>	
Qualified Logging Professional part of operation / on site regularly (proof of training & CE hrs available)	100%
Operating Plan/Map onsite (Work Approval Instructions On Site Crown)	97%
First Aid Supplies	100%
First Aid trained personnel (Copies to be made available)	100%
Personal Protective Equipment (Refer to Safety & Environmental BMP Handbook for required PPE)	100%
Seasonal, provincially required wildfire suppression equipment (between March 15 - Oct 15)	95%
Machine: onboard suppression or (Forwarders - 5lbs or 2 x 2.5lbs / Harvesting - 20lb or 2 x 10lb)	100%
Remote Location & Emergency Response Plans in place	97%
Proper Warning Signs Posted on 2-way Public Traffic Road (Crown)	89%
System to check on Employees who Work Alone	100%
Current Documentation (MSDS Info & Labels)	97%
PHP Policies & Work Instruction Manual or Handbook on site	100%
OHS Compliant Safety System & Documentation (Safety Certificate not required for Private Suppliers)	97%
Lock Out - Tag Out Policy in place	97%
Spill Kit onsite (pads for Crown harvester + absorbent snake (sock) on Crown forwarder)	100%
Pumps (able to be locked for transport or off duty.)	100%
Trailer Permits if not floated.	96%
Central collection spot for Hazardous Material. (Crown)	100%
WHMIS and TDG trained personnel (Copies to be made available)	100%
Waste oil disposal system in place	100%
Tanks Properly Labeled / Placarded to TDG and WHIMS Regulations	92%
Storage tanks located not closer than 100m from any watercourse	100%
Drip catchers on fuel tanks (Crown)	91%
Bridges used and Erosion controlled on approaches to stream crossing	100%
Temporary bridges removed, water courses cleared of debris	100%
No evidence of siltation	100%
Machine Rutting: Within Guidelines (or as permitted by Supervisor)	100%
Ground Disturbance: Within Guidelines	100%
Safety Meeting Minutes	91%
EMS Training - New Employees	75%
<b>JOB QUALITY/HOUSEKEEPING (80% Required to Pass)</b>	
Harvested Merchantable Trees Utilized	100%
Harvest meets Mapped/Operating Plan/Prescription Requirements	100%
Residual Trees retained per Crown Land requirements	100%
Garbage & Litter collected to be discarded	100%
No Discarded Parts/Tires	100%
Disposed of Hazardous Materials	100%

\*Corrections made by contractors and/or PHP staff to restore compliance

## **Crown Contractor Audit Results – 2022 (DNRR Inspection Checklist)**

<i>Operating Plan Map &amp; Work /Approval Instructions On Site</i>	100%
<i>Work /Approval Instructions understood &amp; followed</i>	100%
<b>Conditions of IRM Approval</b>	100%
<b>Sensitive Features Present &amp; SMPs Required</b>	100%
Watercourse crossing(s)	100%
Special Management Zones	100%
Machine Exclusion Zone(s)	100%
Boundary Line(s)	94%
Sensitive Species /Habitat	100%
Corridors	100%
Recreational Trails/Portages	100%
Quality Regeneration	100%
Roadside wood piled >30m from watercourse(s)	100%
<b>Equipment condition &amp; lockout procedures</b>	
Demonstrate lock-out procedures	100%
All guards, screens, and other safety devices in working order on heavy equipment	100%
No visible signs of leaking oil/fuel	100%
Absorbent snake on all forwarders, pads on all other machines	100%
Maintenance done on level ground with minimal surface runoff and >30m from watercourses	100%
<b>Safety - First Aid</b>	
No. 2 first aid kits on site	100%
Qualified first-aiders on-site?	100%
Safety representative (known)	100%
Working alone policy documented and review of policy documented with employees during last year	100%
Remote Location Plan	100%
Safety policy documented, signed, and dated within the last year (References commitment,	100%
Copy of Occupational Health and Safety Act available at the work site	100%
MSDS available at the work site (dated within the last three years)	100%
<i>Proof of Certificate of Recognition</i>	100%
<i>Adequate emergency response procedures in place (i.e. Signs posted from public road to the work site or</i>	100%
<i>"No hunting" signs posted during hunting season (regional rules apply)</i>	100%
<b>Personal Protective Equipment</b>	
CSA approved hard hat worn or be in the machine being operated	100%
CSA grade 2 boots or equivalent	100%
Eye protection (machine maintenance)	100%
High-visibility Clothing	100%

\*Corrections made by contractors and/or PHP staff to restore compliance

Hearing protection	100%
Seat belts worn when operating equipment	100%
<b>Forest Fire Fighting Equipment - Contractor</b>	
Appropriate forest firefighting equipment during fire season (shovels, back tanks, spark arrestors, fire chainsaw /Thinning Saw Use (as per The Forest Professional)	100%
<b>Parts Trailer</b>	
Welders, generators and any other internal combustion engine vented to the outside or a documented Working carbon monoxide detector when internal combustion engine or heat source present or	100%
	100%
<b>Fuel/Oil Handling</b>	
Spill kit is on-site and well maintained	100%
Fuel/Oil handling procedures followed	100%
Containers labelled	100%
Tank capacity indicated on fuel tanks	100%
Shut-off valve on fuel tank, automatic shut-off nozzle on hose, and nozzle holding device	100%
Fuel nozzle drip catcher with drain valve to prevent overflow/spillage	100%
Tanks < 450 litres with capacity and product identification labels	100%
Tanks ≥ 450 litres with capacity and product identification labels and certification stamp	100%
Tanks ≥ 2,000 litres placarded and with manifest for transport	100%
Proof of TDG training (>2000L)	100%
All fuel trailers towed on public roads safety inspected and registered	100%
Fuel tanks & service units >30m from watercourses	100%
<b>Roads</b>	
Road surface in good condition	100%
Ditches and culverts functioning and not damaged	100%
Watercourse crossing(s) as per NSE requirements	

## Private Supplier Audit Results – 2022

<b>LAYOUT COMPLIANCE (100% Required to Pass)</b>	
Worked within Cutting/Property Boundaries	100%
Wildlife clumps (as per instructions)	100%
Wildlife corridors (50M min)	100%
Properly buffered watercourses and wetlands	100%
<b>OPERATION SAFETY/ENVIRONMENTAL COMPLIANCE (90% Required to Pass)</b>	
Qualified Logging Professional part of operation / on site regularly (proof of training & CE hrs available)	100%
Operating Plan/Map onsite (Work Approval Instructions On Site Crown)	100%
First Aid Supplies	100%
First Aid trained personnel (Copies to be made available)	100%
Personal Protective Equipment (Refer to Safety & Environmental BMP Handbook for required PPE)	100%
Seasonal, provincially required wildfire suppression equipment (between March 15 - Oct 15)	88%
Machine: onboard suppression or (Forwarders - 5lbs or 2 x 2.5lbs / Harvesting - 20lb or 2 x 10lb)	100%
Remote Location & Emergency Response Plans in place	90%
Proper Warning Signs Posted on 2-way Public Traffic Road (Crown)	N/A
System to check on Employees who Work Alone	88%
Current Documentation (MSDS Info & Labels)	88%
PHP Policies & Work Instruction Manual or Handbook onsite	100%
DHS Compliant Safety System & Documentation (Safety Certificate not required for Private Suppliers)	75%
Lock Out - Tag Out Policy in place	88%
Spill Kit onsite (pads for Crown harvester + absorbent sock (sock) on Crown forwarder)	100%
Pumps (able to be locked for transport or off duty.)	100%
Trailer Permits if not floated	100%
Central collection spot for Hazardous Material. (Crown)	100%
WHMIS and TDG trained personnel (Copies to be made available)	100%
Waste oil disposal system in place	100%
Tanks Properly Labeled / Placarded to TDG and WHMIS Regulations	100%
Storage tanks located not closer than 100m from any watercourse	100%
Drip catchers on fuel tanks (Crown)	100%
Bridges used and Erosion controlled on approaches to stream crossing	100%
Temporary bridges removed, water courses cleared of debris	100%
No evidence of siltation	75%
Machine Fitting: Within Guidelines (or as permitted by Supervisor)	100%
Ground Disturbance: Within Guidelines	100%
Safety Meeting Minutes	100%
EMS Training - New Employees	100%
<b>JOB QUALITY/HOUSEKEEPING (80% Required to Pass)</b>	
Harvested Merchantable Trees Utilized	100%
Harvest meets Mapped/Operating Plan/Prescription Requirements	100%
Residual Trees retained per Crown Land requirements	N/A
Garbage & Litter collected to be discarded	100%
No Discarded Parts/Tires	100%
Disposed of Hazardous Materials	100%
Road Drains & Culverts Cleared of Debris	100%
Unmerchantable Hardwood Trees Protected	N/A
Damage To Leave Trees Acceptable (includes plantation, regeneration, spacing trees)	N/A
Conservation of known critical wildlife habitat elements, biodiversity & species at risk	100%
SMPs followed in known Forests with Exceptional Conservation Value	100%
SMPs followed for known invasive exotic plants and animals	100%
Known characteristics of special sites preserved	100%
Work Quality Score	100%

\*Corrections made by contractors and/or PHP staff to restore compliance

## Sustainable Forest Management Indicators



The mission of the Woodlands Unit is to provide a reliable, cost effective and high quality supply of wood through the implementation of Sustainable Forest Management.



The vision of Port Hawkesbury Paper LP Woodlands Unit is “that the forest resources, for which we have responsibility, will sustain healthy ecosystems and natural biodiversity, provide a continuous and expanding supply of valuable wood and conserve the forest characteristics of value to society, wildlife and the environment.”



Through the Port Hawkesbury Paper Sustainable Forest Management Policy, the Woodlands Unit implements its Mission and Vision for Sustainable Forest Management (SFM) through 15 Guiding Principles of SFM, 6 Guiding Principles of Wood Procurement, and 16 Standard Practices for SFM.

### **Indicators of Sustainable Forest Management**

Since 2002, PHP has been monitoring and reporting on a variety of sustainable forest management (SFM) indicators. To measure sustainable management over time for a range of forest values, indicators were developed to monitor progress in the maintenance or enhancement of those values.

The Woodlands monitoring program for SFM indicators consists of internal assessments and audit programs. Results from these programs are analysed and summarized on an annual basis to determine if targets are being achieved or have failed to meet set targets. Accordingly, this identifies management actions that must be adjusted to achieve desired outcomes.

Local-level SFM indicators were developed according to the Canadian Council of Forest Ministers’ criteria for sustainable forest management. These criteria are:

- Conservation of Biological Diversity
- Forest Ecosystem Condition and Productivity
- Conservation of Soil and Water Resources
- Forest Ecosystem Contributions to Global Ecological Cycles
- Multiple Benefits to Society
- Accepting Society's Responsibility for Sustainable Development

### **CRITERION 1 - CONSERVATION OF BIOLOGICAL DIVERSITY**



**Conserve biological diversity by maintaining integrity, function, and diversity of living organisms and the complexes of which they are part.**

#### Indicator 1.1 - Species Diversity - Significant Species

OBJECTIVE	Managing and mitigating effects on known occurrences of endangered and threatened species.													
INDICATOR	Annual review of NSDNRR's Significant Species and Habitats Database and other species status lists.													
TARGET	Complete annual review of NSDNRR's Significant Species and Habitats Database, and other species status lists, and implement appropriate management activities where necessary.	VARIANCE None allowed												
2022 Update	<p>The Significant Habitat database was last updated in 2018 (confirmed by Wildlife Division in May 2023) by the provincial Department of Natural Resources &amp; Renewables and provided to PHP to be used in forest management planning activities. The 2018 Significant Habitat database maintained by NSDNRR contains 31,028 ha of significant species habitats potentially affected by forest management activities on PHP's landbase. The areas identified in the 2018 data are categorized into the following:</p> <table border="0"> <tr> <td>Deer Wintering</td> <td>19,325 ha</td> </tr> <tr> <td>Migratory Bird</td> <td>260 ha</td> </tr> <tr> <td>Moose Wintering</td> <td>5,658 ha</td> </tr> <tr> <td>Species at Risk</td> <td>3,127 ha</td> </tr> <tr> <td>Species of Concern</td> <td>1,567 ha</td> </tr> <tr> <td>Other Habitat</td> <td>1,091 ha</td> </tr> </table> <p>These data are used in operational planning and is reviewed by NSDNRR during the harvest approval process. Other species status and appropriate</p>		Deer Wintering	19,325 ha	Migratory Bird	260 ha	Moose Wintering	5,658 ha	Species at Risk	3,127 ha	Species of Concern	1,567 ha	Other Habitat	1,091 ha
Deer Wintering	19,325 ha													
Migratory Bird	260 ha													
Moose Wintering	5,658 ha													
Species at Risk	3,127 ha													
Species of Concern	1,567 ha													
Other Habitat	1,091 ha													









**Indicator 1.3 - Protected Areas - Protected Area Strategy**

OBJECTIVE	To identify and maintain areas reserved from harvest under a protected areas strategy in eastern Nova Scotia.	
INDICATOR	Proportion of area reserved from harvest under a protected area strategy.	
TARGET	Maintain 12% of total area reserved from harvest under a protected area strategy.	VARIANCE +/- 1%
2022 Update	In the 7 eastern counties where PHP operates, there is a total of approximately 209,700 ha (29%) of legally protected Crown land. On just the Crown FULA lands, approximately 82,517 hectares (16%) is legally or pending legal protection (see below map). Additionally, there is 6,147 ha administratively protected by PHP. These areas are also on Crown land and were identified as ecologically significant during the HCVF assessment process.	




**Indicator 1.4 - Protected Areas - Old Forest**

OBJECTIVE	To maintain old forest conditions throughout the landscape.	
INDICATOR Percent of forest management area protected for old forest values.		
TARGET Maintain 8% of forest areas in old forest condition.	VARIANCE +/- 1%	
2022 Update	The current total area reserved as old forest on the Crown FULA lands is 9%. In 2018, the NS Department of Natural Resources & Renewables initiated a new old-growth forest protocol for the assessment of mature climax hardwood stands greater than 11 meters height based on forest inventory data. Since that time, an additional 3,235 hectares of old-growth areas have been identified and protected on PHP's Crown license area.	



The FSC Canada National Forest Stewardship Standard requires certified companies to assess the certified land-base for the sum of protected areas and designated conservation lands. Named the Conservation Area Network, the standard requires that the network must comprise a minimum of 10% of the area of the management unit. The management unit in this case is defined as the certified land-base, which in this case, are the PHP FULA lands.

The protected, pending protected, and old growth/old forest areas identified under indicators 1.3 and 1.4 were used to assess the Conservation Area Network. Since there can be spatial overlap of old growth/old forest areas inside protected areas, the GIS data layers were unioned to remove all overlaps and the possibility of double-counting areas. The GIS analysis has identified that approximately 18% of PHP's certified land-base is considered the Conservation Area Network.






*Old Forest Area, Guysborough  
County, Andrea Doucette, PHP*

## **CRITERION 2 - FOREST ECOSYSTEM CONDITION AND PRODUCTIVITY**

Conserve forest ecosystem condition and productivity by maintaining the health, vitality, and rates of biological production.

### **Indicator 2.1 - Forest Ecosystem Resilience - Natural Regeneration**

OBJECTIVE	To promote Acadian forest characteristics through the use of natural regeneration.	
INDICATOR	Proportion of even-aged management regenerated naturally.	
TARGET	Naturally regenerate with appropriate species 50% of total even-aged management area.	VARIANCE +/- 10%
2022 Update	In 2022, 44% of the total even-aged management area was naturally regenerated.	




*Red Spruce Natural Regeneration*

**Indicator 2.2 - Forest Ecosystem Resilience - Harvest Treatments**

<b>OBJECTIVE</b>	Reduce clearcut area by applying alternative harvest treatments in appropriate ecoregions.
<b>INDICATOR</b>	Proportion of total (softwood and hardwood) area harvested using unevenaged, thinning, shelterwood, selection cut and/or partial cut techniques by EPU.
<b>TARGET</b>	<b>VARIANCE</b>
Increase non-clearcut treatments in appropriate ecoregions to represent 40% of total harvest by 2015 and 50% of total harvest by 2025.	+/- 5 Year Period
<b>2022 Update</b>	<p>PHP has been transitioning to the new provincial 'Silvicultural Guide for the Ecological Matrix', which outlines a range of harvest treatments that are different in their application and operational results from historical treatment types used in the province. Therefore, the traditional clearcut harvest system used previously by PHP is no longer used. The new Variable Retention 10% (VR10) harvest treatment would be the closest to resemble the clearcut system, however, a VR10 maintains more stand structure following harvest than the previous clearcut treatment. Therefore, it is not a straight comparison between a VR10 and a clearcut treatment.</p> <p>For the 2022 update, a summary of treatments completed is provided below as an interim update, while a new set of indicators and targets are under development. It is expected that new values, objectives, indicators and targets (VOITs) will be finalized by the end of 2023.</p> <p>Due to a focus on VR10 treatments in 2022 to aid in the transition to SGEM treatments and management of wood volume needs, larger areas were treated under the VR10 system than the traditional clearcut method. Therefore, the below graph shows an increase in average clearcut size from previous years, however, as mentioned above, a straight comparison cannot be made between VR10 and clearcut treatments.</p>

<b>SGEM Treatment</b>	<b>Total Hectare Treated</b>	<b>% Harvest</b>
Commercial Thinning	1081	29.5%
Medium Retention Irregular Shelterwood Cont Cover	32	0.9%
Medium Retention Irregular Shelterwood Gap	2	0.1%
Partial Overstory Removal Retain 1/3	7	0.2%
Partial Overstory Removal Retain 1/5	500	13.6%
Uniform Shelterwood with Reserves	18	0.5%
Variable Retention 10%	1068	29.2%
Variable Retention 20%	702	19.2%
Variable Retention 30%	254	6.9%
<b>Grand Total</b>	<b>3663</b>	





**Indicator 2.3 - Forest Ecosystem Productivity - Forest Health**

<b>OBJECTIVE</b>	To minimize fire, insect and disease occurrence across the forest landscape.	
<b>INDICATOR</b>	Area (by ha) of forest killed by fire, insect and disease.	
<b>TARGET</b>	Less than 500 ha of forest killed by fire, insect and disease.	<b>VARIANCE</b>
		+ 1000 ha
<b>2022 Update</b>	There was no evidence or recorded data by NS Department of Natural Resources & Renewables for total forest killed by fire, insect, or disease in 2022.	



### **CRITERION 3 - CONSERVATION OF SOIL AND WATER RESOURCES**


Conserve soil and water resources by maintaining their quantity and quality in forest ecosystems.

#### **Indicator 3.1 - Soil Protection - Steep Slopes**

OBJECTIVE	To avoid regular harvesting in identified steep slope areas.	
INDICATOR	Area (by ha) of regular harvest in steep slope areas.	
TARGET	Maintain no regular harvest in areas with greater than 30% average slope.	VARIANCE + 20 ha
2022 Update	A GIS exercise of overlaying steep slope areas with completed harvest areas shows 1 hectare of area. The hectare is made up of several small areas. The steep slope data are based on spatial data that identifies slopes greater than 30% average using contour data. It is not based on the actual % slope of any given area as could be determined on-the-ground. Most often, the areas showing as harvested are slivers due to inaccuracies in the data.	



*MacKenzie Mountain - CB*



### Indicator 3.2 - Water Protection - Watersheds

OBJECTIVE	To protect hydrological functions in all watersheds.	
INDICATOR	Proportion of identified watershed area (that is managed by PHP) in a closed forest condition.	
TARGET	Each watershed shall have 80% of its area (that is managed by PHP) in a closed forest condition.	VARIANCE - 5%
2022 Update	PHP has identified 15 watersheds throughout its management area that are monitored specifically for closed forest condition (> 10 years of age). In 2022, all 15 watersheds had 80% or more of its area in a closed forest condition.	


Watershed Name	% Closed Forest 2022	% Closed Forest 2021	% Closed Forest 2020	% Closed Forest 2019	% Closed Forest 2018	% Closed Forest 2017	% Closed Forest 2016	% Closed Forest 2015	% Closed Forest 2014
Antigonish Municipal (2,169 ha)	100%	100%	100%	100%	100%	100%	100%	100%	100%
Guysborough 1 Municipal (2,778 ha)	86%	82%	84%	86%	86%	86%	92%	91%	96%
Inverness Municipal (131 ha)	85%	85%	85%	86%	85%	85%	93%	92%	92%
Victoria Municipal (974 ha)	93%	95%	95%	97%	95%	97%	99%	98%	98%
Baddeck River (15,439 ha)	95%	95%	95%	94%	95%	95%	95%	96%	94%
East River (9,896 ha)	90%	89%	90%	93%	91%	95%	94%	93%	93%
Grand River (5,749 ha)	97%	95%	94%	93%	92%	90%	89%	89%	85%
Liscomb River (14,824 ha)	97%	96%	96%	96%	95%	94%	92%	91%	90%
Margaree River (35,929 ha)	93%	93%	93%	93%	90%	92%	89%	88%	89%
Middle River (20,527 ha)	90%	90%	90%	92%	93%	94%	89%	90%	87%
Mira River (13,946 ha)	93%	92%	91%	92%	92%	92%	90%	91%	92%
New Harbour River (2,101 ha)	98%	98%	98%	98%	98%	93%	95%	94%	99%
North River (15,830 ha)	90%	90%	86%	85%	88%	90%	85%	86%	83%
River Inhabitant (7,852 ha)	97%	96%	94%	96%	96%	96%	93%	90%	93%
St. Mary's River (53,442 ha)	92%	91%	91%	91%	92%	93%	92%	92%	93%




*St. Mary's River Watershed, Andrea Doucette, PHP*

### Indicator 3.3 - Water Protection - Riparian Zone Management

OBJECTIVE	To protect and maintain all riparian functions.	
INDICATOR	Number of riparian zone non-conformance incidents.	
TARGET	To have zero non-conformance incidents.	VARIANCE None allowed
2022 Update	There was one infraction of the Wildlife Habitat and Watercourse Protection Regulations in 2022. A 30-meter-long section of the watercourse contained a riparian buffer less than the required 20 meters. It was discovered by NSDNRR that it was an unintentional ribboning error, and no sediment or environmental concerns affected the brook.	



The Wildlife Habitat and Watercourse Protection Regulations can be found under Section 40 of the Forests Act. They were developed for application by people working in forestry and are applicable to watercourses and marshes, which include wetlands, lakes, ponds, rivers, streams, creek, estuary, or salt-water body that contains water for at least part of the year. The below image illustrates how special management zones must be established around watercourses and marshes when conducting forestry operations.









*New bridge installation, Paul MacDonald, PHP*


#### **CRITERION 4 - FOREST ECOSYSTEM CONTRIBUTIONS TO GLOBAL ECOLOGICAL CYCLES**

Maintain forest conditions and management activities that contribute to the health of global ecological cycles.

##### **Indicator 4.1 - Forest Carbon - Harvest Volume**

OBJECTIVE	To reduce carbon emissions.	
INDICATOR	Period average volume per hectare harvested.	
TARGET	Increase the average harvest volume by 20% within the next 25 years.	VARIANCE +/- 5 Year Period
2022 Update	The average volume per hectare harvested was 66 tonnes/ha. This is based on all treatments excluding commercial thinnings.	

### Harvest Volume per Hectare Status - 2012 to 2022




### Indicator 4.2 - Forest Carbon - Total Growing Stock

OBJECTIVE	To contribute to total carbon storage through maintenance of above-ground carbon pool.	
INDICATOR	Total growing stock of both merchantable and non-merchantable species on forest lands.	
TARGET	Total growing stock of 21,221,500 m <sup>3</sup>	VARIANCE +/- 1,000,000 m <sup>3</sup>
2022 Update	The total growing stock for softwood is estimated to be 17,895,038 m <sup>3</sup> and the total growing stock for hardwood is estimated to be 15,019,044 m <sup>3</sup> .	

### Indicator 4.3 - Forest Land - Road Construction

OBJECTIVE	To minimize amount of deforested land.	
INDICATOR	Width of permanently disturbed area due to road construction.	
TARGET	Reduce average road width of new and upgraded constructed roads by 10%.	VARIANCE 5% +/-
2022 Update	Average road width for roads measured in 2022 was 8 meters.	




### **CRITERION 5 - MULTIPLE BENEFITS TO SOCIETY**

Sustain flows of forest benefits for current and future generations by providing multiple goods and services.

#### **Indicator 5.1 - Timber and Non-timber Benefits - Hardwood Management**

OBJECTIVE	To increase the future value of the hardwood resource.	
INDICATOR	Area (by ha) of hardwood management.	
TARGET	Manage 2,800 hectares of hardwood in the first five-year period of the 2015 Long-Term Plan.	VARIANCE +/- 500 ha
2022 Update	<p>In 2022, the area of hardwood management was approximately 58 hectares. Since implementing the 2015 long-term plan, PHP has managed 2,462 ha of hardwood forests (88% of 2,800 ha target in the first 5-year period of the plan).</p> <p>Hardwood management has decreased over the last few years since the NS Department of Natural Resources &amp; Renewables initiated an old-growth assessment protocol. A potential old-growth layer of mature climax hardwood based on 11 metre height of the forest inventory is being used to identify potential old-growth stands for further assessment. Stands that are assessed and meet the new old growth definition are protected under the provincial old-forest policy.</p>	




Single Tree Selection, Obidary Road, Antigonish

#### Indicator 5.2 - Communities and Sustainability - Harvest Level

OBJECTIVE	To continue to harvest at a sustainable rate.	
INDICATOR	Annual harvest level.	
TARGET	Harvest 400,000 tonnes of softwood per year.	VARIANCE -10%
2022 Update	In 2022, the softwood volume amount harvested was 172,816 tonnes (43% of annual harvest level). This is considerably lower than previous years because PHP increased chip purchases in 2020 to support local sawmills following the closure of Northern Pulp. This resulted in less harvesting on the Crown license.	

PHP FULA Crown Land			
Year	Softwood Sustainable Harvest Level (tonnes)	PHP Actual Softwood Harvest (tonnes)	% AAC Harvested
2013	400,000	258,292	65%
2014	400,000	336,329	84%
2015	400,000	322,268	81%
2016	400,000	291,951	73%
2017	400,000	329,777	82%
2018	400,000	359,062	90%
2019	400,000	285,152	71%
2020	400,000	230,403	58%
2021	400,000	177,165	44%
2022	400,000	217,030	54%
<b>Total</b>	<b>4,000,000</b>	<b>2,807,429</b>	<b>70%</b>



**Indicator 5.3 - Communities and Sustainability - Third Party Requests**

OBJECTIVE	Where appropriate, provide economical, recreational and cultural opportunities to the general public.	
INDICATOR	Number of reasonable third party requests approved.	
TARGET	Approve all reasonable third-party requests received each year.	VARIANCE 10 requests
2022 Update	A total of 29 third party requests were received in 2022. All but two were approved by PHP. Declined requests were related to a land exchange and the development of a quarry.	



## **CRITERION 6 - ACCEPTING SOCIETY'S RESPONSIBILITY FOR SUSTAINABLE DEVELOPMENT**

Society's responsibility for sustainable forest management requires that fair, equitable and effective forest management decisions are made.

### **Indicator 6.1 - Aboriginal and Treaty Rights - Respect First Nations**

OBJECTIVE	To provide opportunities to better understand, recognize and respect local Mi'kmaw and Treat Rights.	
INDICATOR	Number of opportunities to meet with Mi'kmaw organizations and/or community representatives.	
TARGET	Ensure a minimum of six opportunities to meet with Mi'kmaw organizations and/or individuals annually.	VARIANCE - 1 Meeting
2022 Update	In 2022, the company met at least 10 times with Mi'kmaq organizations, communities, or individuals related to forest management agreements with CMM and UINR, and other initiatives related to Mi'kmaq Forestry Initiative lands and Indigenous Protected & Conservation Area lands. The Free Prior Informed Consent (FPIC) process committee met four times in 2021.	

### **Indicator 6.2 - Aboriginal and Treaty Rights - First Nation Agreements**




OBJECTIVE	To build capacity within Mi'kmaq communities to provide increased employment opportunities for Mi'kmaw individuals.	
INDICATOR	Volume harvested under agreements with Mi'kmaq communities.	
TARGET	To increase the softwood volume harvested under First Nation agreements to 30,000 tonnes.	VARIANCE - 5,000 tonnes
2022 Update	<p>In 2022, the total volume harvested by Unama'ki Institute of Natural Resources was 0 tonnes. This was due to lack of contractor capacity as well as redirection of resources on the Mi'kmaq Forestry Initiative and Indigenous Protected &amp; Conservation Area project.</p> <p>In 2022, the total volume harvested by Confederacy of Mainland Mi'kmaq was approximately 34,000 tonnes.</p>	



























## Summary of SFM Indicators

Since 2002, the Woodlands Unit has developed, monitored and reported on a suite of SFM indicators. During that time, indicators have been revised or removed to be replaced with improved indicators based on a new understanding of forest dynamics. PHP has made significant progress in achieving several of our targets and others are on their way. To improve their conditions over time, management decisions and activities will be implemented to result in indicators trending towards their desired targets.

-  Target Achieved within Acceptable Variance
-  Target On-going
-  Target Not Achieved within Acceptable Variance

1.1 Annual review of NSDNRR's significant species and habitats database, and other species status lists.	
1.2 Percent of CMZs meeting the 60% closed forest condition guideline.	
1.3 Proportion of area reserved from harvest under a protected areas strategy by EPU.	
1.4 Percent of defined forest area by EPU protected for old forest values.	
2.1 Proportion of natural regeneration in reforestation program.	
2.2 Proportion of total (softwood and hardwood) area harvested using unevenaged, thinning, shelterwood, selection cut and/or partial cut techniques by EPU.	
2.3 Area of forest disturbed by fire, insect and disease.	
2.4 Area (by ha) affected by budworm outbreak on the Cape Breton Highlands.	
3.1 Area (by ha) of regular harvest in steep slope areas.	
3.2 Proportion of identified watershed area (that is managed by PHP) in closed forest condition.	
3.3 Number of riparian zone non-conformance incidents.	

3.4 Number of road construction and stream crossing incidents according to company guidelines.	
4.1 Yearly average volume per hectare harvested.	
4.2 Total growing stock of both merchantable and non-merchantable species on forest lands.	
4.3 Width of permanently disturbed area due to road construction.	
5.1 Area (by ha) of hardwood management.	
5.2 Annual harvest level.	
5.3 Number of reasonable 3rd party requests approved.	
5.4 Proportion harvest volume sold to other mills.	
6.1 Number of opportunities to meet with Mi'kmaq community representatives.	
6.2 Volume harvested under agreements with Mi'kmaq communities.	
6.3 Level of investment and contribution to education and extension initiatives.	

# FSC Canada Specific Monitoring Updates

## Introduction

The new FSC National Forest Stewardship Standard of Canada, released in 2020, has identified specific indicators requiring monitoring and assessment as they relate to significant environmental impacts resulting from management activities. This section summarizes those requirements and PHP's monitoring results, where applicable.

FSC INDICATOR    Poor Regeneration	
<p><u>Description</u> Regeneration is when forests are restocked, either naturally or through plantings, so a new healthy forest develops over time.</p> <p><u>Monitoring Approach</u> Many areas harvested by PHP are scheduled for planting following harvest. For areas that are not, a cutover assessment is completed within two years to determine if planting is required and where natural regeneration is lacking, planting will be carried out within two years from harvest.</p> <p>Based on the vegetative community and harvest treatment employed, three options are possible:</p> <ol style="list-style-type: none"> <li>1. No regeneration treatments are required for the area. A cutover assessment is still completed.</li> <li>2. It is uncertain whether a regeneration treatment is needed. A cutover assessment is scheduled in two years from harvest.</li> <li>3. There is a need to carry out a regeneration treatment. The cutover assessment will be completed, and the treatment scheduled.</li> </ol>	
2022 Update	In 2022, there was 1,345 hectares of cutover assessment completed. A total of 1,539 hectares was planted in 2022.

FSC INDICATOR    Invasiveness or other adverse impacts associated with alien species	
<p><u>Description</u> An alien species is an insect, micro-organism or plant that is found outside of its natural range. They can affect the natural biodiversity of local ecosystems by crowding out species native to a particular area.</p> <p>PHP has planted approximately 20,000 hectares of Norway spruce since the 1970's but stopped the practice in 2014. Some of the stands making up the 20,000 hectares are now unlicensed Crown, so PHP only monitors Norway spruce stands still included in its Crown license agreement</p>	

with the province. Norway spruce is not considered to be a highly-invasive species, however, PHP continues to monitor these stands for regeneration and seeding outside the planted area.

Monitoring Approach

PHP has been monitoring planted Norway spruce stands since 2013. PHP annually selects two stands minimum for assessment that have been planted with Norway spruce to determine if the species is regenerating and seeding outside of the planted area. A Norway spruce regeneration survey tally sheet is used to tally total softwood regeneration and total Norway spruce regeneration. Plots are 1:1000<sup>th</sup> of a hectare (1.78 m) and at an intensity of 1 plot per hectare. Two plots per stand are also taken in adjacent stands to determine if any seed has dispersed outside the planted stand.

2022 Update

In 2022, two Norway spruce stands were selected for assessment. Regeneration and/or seeding inside or outside the planted stand was not found.

FSC INDICATOR Adverse effects of fertilizers

Description

Fertilizers used in forested areas can benefit tree growth, however, they can also affect soil and water quality because of chemicals found in them.

Monitoring Approach

FSC Canada states that the use of fertilizers applied directly on the forest management unit is the focus of this monitoring requirement and does not include fertilizer that may be used in the growing of nursery stock.

2022 Update

PHP does not apply fertilizers directly on the forest management unit, and therefore, no monitoring is required.

FSC INDICATOR Adverse effects of pesticides

Description

FSC Canada defines pesticides as any substance or preparation prepared or used in protecting plants or wood or other plant products or human health or livestock or biodiversity from pests; in controlling pests; or in rendering such pests harmless. (This definition includes insecticides, rodenticides, acaricides, molluscicides, larvacides, fungicides and herbicides).

Monitoring Approach

Not applicable to PHP.

2022 Update

PHP has not applied herbicides for forest management since 1997. Other pesticide use defined by FSC Canada is not applied by the company.

FSC INDICATOR Adverse effects of biological control agents	
<p><u>Description</u> FSC Canada defines biological control agents as organisms used to eliminate or regulate the population of other organisms.</p> <p><u>Monitoring Approach</u> Not applicable to PHP.</p>	
2022 Update	PHP does not use biological control agents in its forest management.

FSC INDICATOR Physical damage to soil, loss of soil nutrient and loss of productive forest area					
<p><u>Description</u> Healthy soils are a key component of responsible and sustainable forest management. Soil quality is defined as the capacity of a soil to function within an ecosystem to sustain biological productivity, maintain environmental quality, and promote plant and animal health. In addition, soil health definitions include maintaining the integrity of nutrient cycling and resilience to disturbance or stress. Tree or stand growth has often been used as an indicator of soil productivity changes.</p> <p><u>Monitoring Approach</u> PHP implements procedures and monitoring regarding ground disturbance and rutting resulting from forest management activities. Ground disturbance means any area where the surface organic horizons are completely removed or redistributed on a site (mineral soil has been exposed). Harvested sites with 10% or less of ground disturbance is considered to be harvested with due regard for soil protection (also within rutting limitations).</p> <p>PHP aims to minimize rutting by using pre-planning methods such as time of harvest (ie. season) and soil types. PHP supervisors monitor sites for excessive rutting as part of regular site visits. If rutting is deemed excessive or becomes greater than the maximum rut allowed, forwarding on that trail will cease until remediation measures are put in place or until site conditions approve (operations will be ceased).</p> <p>Ground disturbance and rutting are monitored on all Crown harvest sites using an internal audit process. Compliance with PHP's procedures are checked during seasonal audits as well as on post-harvest inspections. Crown contractors also check rutting conditions weekly and record their assessment on the contractor weekly check list.</p>					
2022 Update	<p>In 2022, 30 Crown contractors were audited by PHP operations supervisors. The overall results of these audits are shown on page 10 of this report. For ground disturbance and machine rutting guidelines specifically, there was full compliance with PHP procedures.</p> <table border="1"> <tbody> <tr> <td>Machine Rutting: Within Guidelines (or as permitted by Supervisor)</td> <td>100%</td> </tr> <tr> <td>Ground Disturbance: Within Guidelines</td> <td>100%</td> </tr> </tbody> </table>	Machine Rutting: Within Guidelines (or as permitted by Supervisor)	100%	Ground Disturbance: Within Guidelines	100%
Machine Rutting: Within Guidelines (or as permitted by Supervisor)	100%				
Ground Disturbance: Within Guidelines	100%				

FSC INDICATOR Adverse effects of increased access

Description

Forest access roads are a needed element of forest management, but their impact on the environment can be significant. Increased access can cause harm to wildlife through excessive hunting or predation, changes in wildlife habitat use due to noise, and entrance into ecologically sensitive or protected areas.

Monitoring Approach

The forest roads used by PHP to access forest management areas are owned by the provincial government. PHP does not have the mandate to close access to roads, but occasionally PHP supervisors will pull temporary bridges from roads to use in other locations. Signs are posted to warn potential access users about a crossing site. If access to an area through a road was deemed necessary to close, the provincial government would make that decision and implement their own measures to close access.

2022 Update

No forest roads were closed to public access in 2022 due to PHP or DNR activities. However, Hurricane Fiona caused significant damage to forested areas in September 2022. Due to accessibility issues, some roads were impassable until repairs and/or downed trees were removed from road surfaces.

FSC INDICATOR Site level damage of harvesting and extraction on residual trees and on environmental values

Description

Site level damage on residual trees and other environmental values from harvesting activities can affect the overall quality of the site for healthy regeneration and biological diversity.

Monitoring Approach

For several years, PHP has implemented procedures for retaining residual standing trees for ecological, aesthetic or production functions. Interim Retention Guidelines have recently been developed by the provincial government and are to be applied on Crown land. The purpose of the guidelines is to increase the amount of retention above the required wildlife clumps under the Wildlife Habitat & Watercourse Protection Regulations. The guidelines are summarized as:

1. Leave approximately 10-30% stand-retention when applying an overstory removal or seed tree harvest.
2. Leave retention trees distributed though the stand wherever possible.
3. Other priorities for retention:
  - a. Leave uncommon tree species that form a small proportion of stand (eg. Ironwood, late successional intermediate to tolerant species, large old super canopy trees)
  - b. Wildlife trees and biodiversity features
  - c. Growing stock and advanced regeneration (preferably late successional intermediate to tolerant species)
  - d. Deep rooted overstory of wind-firm trees (sugar maple, yellow birch, white pine, red oak, white ash)

- e. Shallow rooted overstory of wind-firm trees (red spruce, eastern hemlock, white spruce, red maple)
- f. Deep rooted non-late successional intermediate to tolerant species overstory (red pine, jack pine, white birch)

## 2022 Update

Internal audits completed in 2022 on Crown land operations resulted in 100% compliance to being within acceptable levels of damage to leave trees. PHP measures acceptable levels of damage or scarring as being under 4 inches squared.

## FSC INDICATOR

Damage caused by inappropriate storage or disposal of waste materials

Description

Improper disposal or inappropriate storage of waste materials can have negative consequences for soil and water quality, as well as wildlife.

Monitoring Approach

PHP implements procedures on the disposal of hazardous materials, and storage and handling of diesel fuel tanks, on all operations. Requirements on PHP operations includes:

- Training on transportation, disposal, storage and handling of hazardous materials and diesel fuel tanks.
- Storage tanks and facilities to be located not closer than 100 meters from any watercourse, pond or lake.
- Fully stocked spill kits are present on all active operations where machinery is present.
- Weekly inspections of storage tanks
- Full tanks not filled to more than 95% capacity
- Discharge hoses must be fitted with nozzles
- Pumps are well maintained and kept free of leaks
- Spills of greater than 70 litres must be reported to Department of Environment
- Garbage and other waste materials must be properly disposed of

## 2022 Update

The 2022 Crown contractor audits resulted in 100% compliance with proper disposal of hazardous materials, no discarded parts or tires, garbage and litter properly discarded, and waste oil disposal system in place. Drip catchers on fuel tanks were at 91% compliance and storage tanks not closer than 100 meters from any watercourse was at 100% compliance.

No environmental incidents were recorded in 2022 from inappropriate storage or disposal of waste materials.

# High Conservation Value Forest Effectiveness Monitoring Program

## Introduction

This HCVF Effectiveness Monitoring Program was developed to fulfill the requirements of Principle 9 of the FSC Maritimes Standard. To meet Principle 9 of the standard, forest managers must complete an assessment of their forest lands to identify high conservation values. There are six distinct categories that give an area critical conservation significance. FSC Canada defines an HCVF as:

*High Conservation Value Forests are those that possess one or more of the following attributes:*

*a) Forest areas containing globally, regionally or nationally significant:*

*i) Concentrations of biodiversity values (e.g., endemism, endangered species, refugia); and/or*

*ii) Large landscape level forests, contained within, or containing the management unit, where viable populations of most (if not all) naturally occurring species exist in natural patterns of distribution and abundance.*

*b) Forest areas that are in or contain rare, threatened or endangered ecosystems.*

*c) Forest areas that provide basic services of nature in critical situations (e.g., watershed protection, erosion control).*

*d) Forest areas fundamental to meeting basic needs of local communities (e.g., subsistence, health) and/or critical to local communities' traditional cultural identity (areas of cultural, ecological, economic or religious significance identified in cooperation with such local communities).*

Once HCVF's are identified on the land-base, the forest manager must decide how these areas will be managed to maintain or enhance the values that are present. Where values exist, monitoring is needed to show that the prescribed management is effective. PHP's effectiveness monitoring program identifies two stages of monitoring for several HCVF's.

The first level of monitoring is for the basic operational procedures or special management practices that have been identified for the value (e.g. buffer zones, maintenance of special habitat characteristics, protection). This level of monitoring is typically done on an annual basis. It is also important at this stage of monitoring to ensure that PHP is aware of and implementing the best management approach, prescriptions, and/or special management practices as defined by an outside organization. Therefore, PHP will also contact known experts and/or



organizations to gather any new available information regarding management or to verify that its current management approach is the best at that time. All HCV's have an identified operational monitoring protocol that is implemented on an annual basis.

The second level of monitoring, if applicable, is strategic monitoring to determine if the HCV attribute(s) are being maintained on the landscape. For example, for a species at risk such as Boreal Felt Lichen, it is important to determine that the identified forest habitat is still suitable and that the species is still present in the habitat. Contrary to operational monitoring, not all HCV's require a strategic level of monitoring. For example, the HCV of old forest are legally protected and therefore, not available for active forest management. Therefore, the attribute of maintaining old forests on the landscape is automatically preserved. Alternatively, strategic monitoring is important for species at risk when the objective is to maintain areas of good forest habitat for a threatened species, and to ensure that the species is still present in this habitat.

For strategic monitoring, PHP recognizes that there is a required level of involvement by government agencies and/or other organizations for the monitoring of species populations and health. It is PHP's intention to collaborate with these agencies to collect the necessary information.

## HCVF Category 1 – Biodiversity – Species at Risk

### HCV – American Marten Habitat



HCV ATTRIBUTE		Species at Risk – Habitat and Population	
<b>OPERATIONAL MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Maintain and/or enhance American Marten habitat in home range patches		
INDICATOR	Reserve stand structure as required within harvest areas located within the American Marten Habitat Management Zone		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	
		Habitat management requirements are implemented through the DNRR approval process for Crown lands. Monitor implementation of stand structure reserve using TFM. Verify annually that special management practices are still current and/or make operational changes as needed.	
DATA SOURCES	The Forest Manager (TFM); PHP & DNRR field audits	COST AND DIFFICULTY	
		Low to Moderate - Dependant on PHP's required level of involvement	
<b>LONG-TERM STRATEGIC MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	American Marten population recovery		
INDICATOR	Population estimates / use within the Marten Habitat Management Zone		
MONITORING STRATEGY			
DNRR is responsible for population inventory and studying habitat use.			
DATA SOURCES	American Marten Recovery Team DNRR Manager, Wildlife Resources	COST AND DIFFICULTY	

	Low to High - Dependant on PHP's required level of involvement
<b>FOREST MANAGEMENT PRESCRIPTION</b>	
<p>These prescriptions are applied throughout the Cape Breton Highlands:</p> <ul style="list-style-type: none"> <li>- 12-14 standing and live mature trees per ha must be left evenly spaced throughout the harvest site;</li> <li>- These are in addition to all other requirements of the Wildlife Habitat and Watercourse Protection Regulations;</li> <li>- Large yellow birch trees should be left standing where possible;</li> <li>- Special management practices for commercial thinning operations in marten patches;</li> <li>- Harvest sites should maintain at least 100 m<sup>3</sup> of coarse woody debris/ha and mean maximum diameter of downed logs should exceed 22 cm.</li> </ul> <p>There are also 30 home range patches established within the Marten Habitat Management Zone. These patches may 'migrate' within the zone, but must be a minimum 500 ha in size, circular in shape, and contain a minimum 60% marten habitat as described in the marten recovery strategy.</p>	
<b>2022 MONITORING UPDATE</b>	
<ol style="list-style-type: none"> <li>1. All harvest treatments applied throughout the Cape Breton Highlands included the above management prescriptions as required and approved by DNRR regional staff.</li> <li>2. A total of 95 hectares was thinned and 17 hectares treated with 10% variable retention. A total of 86 hectares was treated with pre-commercial thinning. These treatments were approved in the Marten Habitat Management Zone as per DNRR's approval process (see below map).</li> <li>3. The American Marten Recovery Strategy (2007) estimates that the Marten population is less than 50 individuals. A re-introduction program began in 2007, which brought 130 individuals into Cape Breton from New Brunswick. A total of 35 individuals were collared, but their movements were lost approximately 6 months after release. DNRR does have pictures, have live-trapped, recorded tracks in snow and have received reports/sightings of marten in the Cape Breton Highlands (Peter Austin-Smith, pers. comm., 2013). A goal of the Marten Recovery Team is to have <math>\geq 30</math> marten in Cape Breton by 2010, <math>\geq 100</math> by 2030 and <math>\geq 350</math> by 2040.</li> <li>4. Information provided by R. Milton, NSDNRR on May 21, 2019 states "during February and March 2018, bait and cameras were set in each of 30 patches forecast in the American Marten Special Management Practices to be available as habitat by 2019. Cameras were set for a minimum of 3 weeks to record whether American Marten visited the bait. Single and occasionally pairs of marten were recorded in 15 of the 30 patches, even though only 5 of these 'occupied' patches met desired habitat conditions described as greater than 30% softwood, greater than or equal to 6 m high and basal area greater than or equal to 18 m<sup>2</sup>/ha. Of the 30 forecasted sites from 2004, 24 will not meet desired habitat conditions by 2019. Marten not being recorded in the other 15 patches cannot be interpreted as confirming absence, but rather not recorded at this time. Cursory</li> </ol>	


examination of marten presence and patch conditions suggests flexibility in coarse habitat descriptors used in the Special Management Practices, or undescribed critical features common to documented occupied patches. This past winter, bait and cameras were established at 13 sites on the Keppoch north of the 2019 patches. Although 5 cameras still need to be retrieved, 3 of the 8 sites had marten present. Initial consolidation of American Marten records is ongoing. However, it is of interest to note that since 2010, there have been nearly 200 valid reports of either tracks, visual sightings, or camera records. Nearly 150 of these reports have occurred since 2015 which indicates the augmentation project conducted from 2007 through 2009 has been successful, at least in the immediate term, in maintaining a breeding population of American Marten on the highlands.”

5. NSDNRR is currently reviewing the draft NS recovery plan for marten to determine a definition of core habitat under the NS Endangered Species Act. Once completed, that definition will be implemented into the recovery plan.

#### SUPPORTING DOCUMENTS/REFERENCES

Marten Special Management Practices, NSDNRR July 2012; Proposed Marten Recovery Strategy, NSDNRR May 2006; Status Report on American Marten, F. Scott June 2001; Weaseling their Way Back into Cape Breton? Assessing the Feasibility of Augmenting the Cape Breton Island Marten Population Through Habitat Suitability and Individual-based Modeling, Rebecca Jepessen, Acadian University Thesis, 2010.

#### American Marten Habitat Management Zone - Cape Breton Highlands



## HCV – Mainland Moose Habitat



HCV ATTRIBUTE		Species at Risk – Habitat and Population	
<b>OPERATIONAL MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Maintain and/or enhance Mainland Moose habitat		
INDICATOR	Reserve stand structure as required within harvest areas located within the five Significant Mainland Moose Population Concentration areas mapped by NSDNRR		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	
		Habitat management requirements are implemented through the DNRR approval process for Crown lands. Monitor implementation of stand structure reserve using TFM. Verify annually that special management practices are still current and/or make operational changes as needed.	
DATA SOURCES	The Forest Manager (TFM); PHP & DNRR field audits	COST AND DIFFICULTY	
		Low to Moderate - Dependent on PHP's required level of involvement	
<b>LONG-TERM STRATEGIC MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Mainland Moose population recovery		
INDICATOR	Population estimates / use of population concentration areas		
MONITORING STRATEGY			
DNRR is responsible for population inventory and studying habitat use.			
DATA SOURCES	Mainland Moose Recovery Team	COST AND DIFFICULTY	


DNRR Biologist Lisa Doucette	Low to High - Dependent on PHP's required level of involvement
<b>FOREST MANAGEMENT PRESCRIPTION</b>	
<ul style="list-style-type: none"> <li>- Moose shelter patches (within 250 metres of the edge of any forest harvest (partial or clearcut) a minimum of two closed canopy coniferous stands &gt; 3 hectares in area)</li> <li>- The preferred conditions for shelter patches using the NSDNRR forest GIS inventory specifications are: <ul style="list-style-type: none"> <li>- FORNON = 0, and</li> <li>- 50-80% softwood, and</li> <li>- &gt;= meters height, and</li> <li>- crown closure &gt;= 60%</li> </ul> </li> <li>- Moose retention patches (Smaller coniferous must also be retained within each harvest area to provide temporary shelter and concealment)</li> <li>- Moose buffers (Forested buffers should be retained around and or near open wetlands, watercourses, and waterbodies)</li> <li>- Roads and access points (Development of roads and improved trails should be avoided where extended extraction trails can be used as an alternative)</li> <li>- Coarse woody debris (leave tree tops and substantial amounts of woody debris on extraction trails to discourage access)</li> <li>- Decommission roads to reduce human access</li> </ul>	
<b>2022 MONITORING UPDATE</b>	
<ol style="list-style-type: none"> <li>1. All harvest treatments within the mainland moose concentration areas for shelter patches and forest cover condition are in compliance with the May 2022 mainland moose SMP as required and approved by NSDNRR.</li> <li>2. The Mainland Moose Recovery Plan (2007) estimates approximately 1000-1200 individuals on mainland Nova Scotia. This is the most current information available on mainland moose population numbers.</li> <li>3. The Action Plan for the Recovery of Eastern Moose in Mainland Nova Scotia was released to the public in 2016. A total of 14 actions have been identified which are in different phases of completion. The action items related to the Mainland Moose include increased understanding of genetics, cause of death/illness, long-term monitoring, threats, poaching, translocation feasibility, review and adapt forest management practices as habitat requirements are better understood, public awareness and engagement. In May 2020, DNRR released revised special management practices as a result of the action plan.</li> <li>4. In relation to the Action Plan, PHP is a partner on a new research study to “develop tools to provide decision support in forest management planning at multiple spatial scales for moose habitat requirements”.</li> </ol>	

5. DNRR has conducted flight surveys and thermal imagery surveys in moose concentration areas but have yet to determine mainland-wide provincial estimates of population size. The government does have outside help to extrapolate the survey results to the broader area, given the use of different methods and a new thermal imagery technique. The DNRR will be appointing a new small recovery team which will assist in updating the current recovery/action plan and provide guidance related to on-going work.
6. On May 24, 2019 the Chronicle Herald newspaper published an article titled “From high overhead, a sobering look at a moose population in deep trouble”, which is a summary of survey results mentioned in the previous point. This information was obtained by the CBC to highlight the rapid decline of mainland moose population numbers.  
(<https://www.cbc.ca/news/canada/nova-scotia/mainland-moose-nova-scotia-decline-1.5148572>)

#### SUPPORTING DOCUMENTS/REFERENCES

Mainland Moose Special Management Practices, NSDNRR July 2012; Recovery Plan for Mainland Moose in Nova Scotia, March 2007; Action Plan for the Recovery of Eastern Moose in Mainland Nova Scotia 2014-2018

#### Mainland Moose Concentration, Retention and Bog Buffer Areas



## HCV – Canada Lynx Habitat



HCV ATTRIBUTE		Species at Risk – Habitat and Population	
<b>OPERATIONAL MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Maintain and/or enhance Canada Lynx habitat		
INDICATOR	Reserve stand structure in lynx bog buffers within harvest areas located throughout the Cape Breton Lynx Range		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	
		Habitat management requirements are implemented through the DNRR approval process for Crown lands. Monitor implementation of stand structure reserve using TFM. Verify annually that special management practices are still current and/or make operational changes as needed.	
DATA SOURCES	The Forest Manager (TFM); PHP & DNRR field audits	COST AND DIFFICULTY	
		Low to Moderate - Dependant on PHP's required level of involvement	
<b>LONG-TERM STRATEGIC MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Canada Lynx population recovery		
INDICATOR	Population estimates / use of treed bog leave areas		
MONITORING STRATEGY			
DNRR is responsible for population inventory and studying habitat use. A joint project between DNRR and Acadian University is assessing the efficacy of the 100-meter treed bog buffers. The project began in January 2011 and ended in 2015.			
DATA SOURCES	Canada Lynx Recovery Team	COST AND DIFFICULTY	




DNRR Biologist Peter Austin-Smith	Low to High - Dependant on PHP's required level of involvement
<b>FOREST MANAGEMENT PRESCRIPTION</b>	
<p>- To supplement red squirrel habitat, which is an important food source for lynx, a wider buffer strip of 100 m of unharvested forest should be left around all treed bogs in the Cape Breton lynx range.</p> <p>- Where possible, decommission secondary, non-main trunk forest access roads following harvest.</p> <p>- Plan access roads to have dead ends.</p> <p>- Plan harvesting to allow decommissioning of sectors of road networks.</p> <p>- Where possible, narrow and orient road right-of-ways to create shade conditions to reduce snow compaction, thereby reducing ease of travel for coyotes.</p> <p>- Maintain a continuous supply of &gt;50ha patches of mid-regeneration (15-35-year old) conifer dominated habitat that supports high densities of snowshoe hare over each lynx management unit.</p> <p>- Create a landscape that will maintain a continuous presence of a mosaic of successional stages, especially mid-regeneration patches that will support resident lynx.</p> <p>- Employ silvicultural techniques that create, maintain, or prolong use of stands by high populations of snowshoe hares.</p> <p>- Retain coarse woody debris for denning sites.</p>	
<b>2022 MONITORING UPDATE</b>	
<ol style="list-style-type: none"> <li>1. The Canada Lynx Recovery Plan is currently under review by the DNRR, Wildlife Division, to identify core habitat areas in the Cape Breton Highlands. Until this review is finalized, all planned harvest treatments within Canada lynx treed bog buffers are prohibited.</li> <li>2. The Canada Lynx Recovery Strategy (2005) estimates approximately 50-500 individuals in the Cape Breton lynx range. This is the most current information available on Canada lynx population numbers.</li> <li>3. Some work happening related to habitat issues for both American Marten and Canada Lynx (joint recovery team). Still being developed by DNRR Species at Risk group. Looking to get more funding to do habitat modeling and management issues in CB Highlands. Nothing approved yet; still in discussion phase.</li> <li>4. DNRR and Acadia University collaborated on a research study in the Cape Breton Highlands from 2013 to 2015. Some results of that work include: <ul style="list-style-type: none"> <li>• 1 km long track surveys were conducted Jan – March 2013-2015 for a total 243.9 km,</li> <li>• 87 transects were established between 350 and 500 masl</li> <li>• Transects were paired to examine use of SMP buffers with nearby treated stands</li> </ul> </li> </ol>	

- Buffers typically had “natural” stands although some buffer lengths did encounter short sections of treated areas
  - The number of lynx tracks were relatively similar between 2013 and 2014 but increased dramatically in 2015 which is believed due to the increase in recorded hare tracks beginning in 2014 and extending into 2015
  - All prey species have very similar overall patterns in terms of natural vs treated habitats and buffer vs non-buffer areas
  - Marten and coyote exhibit similar habitat use patterns to prey species while lynx exhibit higher use in natural habitat and buffer zones
  - Occupancy modelling reveals the highest probability of occupancy for lynx occurs in or near buffers in natural areas. Moving away from buffer areas, the probability of occupancy drops dramatically especially in treated stands
  - Lynx will occupy treated areas in buffers but at much lower rate
  - Conclusion is buffer zones with natural structure exhibit a much higher occupancy rate for lynx than treated areas even during an expanding population in response to an increased prey base
5. NSDNRR is currently reviewing the draft NS recovery plan for lynx to determine a definition of core habitat under the NS Endangered Species Act. Once completed, that definition will be implemented into the recovery plan.

#### SUPPORTING DOCUMENTS/REFERENCES

Canada Lynx Special Management Practices NSDNRR July 2012; Lynx Recovery Strategy Feb 2007; Endangered Canada Lynx Proposed Project: Assessing the interim 100 metre buffers around highland bogs, DNRR 2014; DNRR Wildlife Manager Randy Milton, pers. comm. 2018

#### Canada Lynx Treed Bog Buffer Areas



## HCV – Wood Turtle Habitat




HCV ATTRIBUTE		Species at Risk – Habitat and Population	
<b>OPERATIONAL MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Maintain and/or enhance Wood Turtle habitat		
INDICATOR	Implementation of temporal and spatial special management practices for wood turtles		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	
		Habitat management requirements are implemented through the DNRR approval process for Crown lands. Monitor implementation of temporal and spatial requirements using TFM. Verify annually that special management practices are still current and/or make operational changes as needed.	
DATA SOURCES	The Forest Manager (TFM); PHP & DNRR field audits	COST AND DIFFICULTY	
		Low to Moderate - Dependant on PHP's required level of involvement	
<b>LONG-TERM STRATEGIC MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Wood Turtle population recovery		
INDICATOR	Population estimates		
MONITORING STRATEGY			
DNRR is responsible for population inventory and studying habitat use.			

<p>DATA SOURCES</p> <p>Wood Turtle Recovery Team</p>	<p>COST AND DIFFICULTY</p> <p>Low to High - Dependant on PHP's required level of involvement</p>
<p><b>FOREST MANAGEMENT PRESCRIPTION</b></p>	
<ul style="list-style-type: none"> <li>- Adjust the timing and location of motorized vehicle use for forest management activities to when Wood Turtles are inactive or less likely to be occupying terrestrial habitat (Nov – March)</li> <li>- Use temporary bridge crossings for perennial streams to avoid altering stream bank, creating erosion and sedimentation, damaging stream bed, and impacting overwintering turtles.</li> <li>- Forest management roads and landings should not be constructed parallel to watercourses within 200 m of watercourses where wood turtles occur.</li> <li>- Special management practices for overwintering, nesting, and basking (see DNRR Special Management Practices 2012).</li> </ul>	
<p><b>2022 MONITORING UPDATE</b></p>	
<ol style="list-style-type: none"> <li>1. All harvest treatments within wood turtle habitat zones have the above management prescriptions implemented as approved by DNRR regional staff.</li> <li>2. The population of wood turtles in PHP's operating area is estimated to be approximately 3,500 individuals (M. Pulsifer, pers. comm., 2013)</li> <li>3. No changes have been made to the special management practices for wood turtle as issued by DNRR, however, new critical wood turtle habitat areas identified by Environment Canada in 2020 have been incorporated into the provincial wood turtle habitat layer. These new critical wood turtle areas are off-limits to all forest management activities including road building.</li> <li>4. Monitoring for new locations has not been a funding priority for DNRR, and outside funding has been significantly reduced. A graduate student working on overwintering habitat and communal distribution has just finished his MSc at Acadia. There is nothing significantly different with DNRR's understanding of wood turtle distribution within the watershed.</li> <li>5. DNRR is not receiving reports of dead turtles that can be linked directly to the forest industry.</li> <li>6. The final recovery strategy for wood turtle was expected to be released in the latter half of 2019, however it is currently not finalized as of mid-2022.</li> <li>7. "A wood turtle record of variance was approved earlier this year which has not provided much further clarity on the direction relative to a draft federal Recovery Action Plan. Currently, the federal plan is still draft and also the proposed federal critical habitat is still proposed" (T. Power, pers. Comm, July 2020)</li> <li>8. DNRR is currently reviewing the Wood Turtle SMP to ensure the federal and provincial habitat areas are appropriately managed. Until this is complete, all federal and provincial habitat areas are prohibited from all forest management activities.</li> </ol>	


## SUPPORTING DOCUMENTS/REFERENCES

Wood Turtle Special Management Practices NSDNRR July 2012; Protecting and Conserving Wood Turtles: A Stewardship Plan for NS, 2003

## Wood Turtle Special Management Areas &amp; Critical Habitat - Eastern Mainland



**Wood Turtle Special Management Areas & Critical Habitat - Cape Breton**



**HCV – Bicknell’s Thrush Habitat**



<b>HCV ATTRIBUTE</b> Species at Risk – Habitat and Population	
<b>OPERATIONAL MONITORING PROGRAM</b>	
<b>MANAGEMENT OBJECTIVE</b>	Maintain and/or enhance Bicknell’s Thrush habitat
<b>INDICATOR</b>	Implementation of temporal and spatial special management practices for Bicknell’s Thrush
<b>MONITORING/REPORTING FREQUENCY</b>	<b>MONITORING STRATEGY</b>
Annual	Identify planned pre-commercial thinning activities in Bicknell’s Thrush habitat in the

	Highlands, so field surveys by Bird Studies Canada can first be conducted to identify presence/absence of the bird during their breeding/nesting season (May, June, July). Monitor implementation of leave patches in thinned/cleared areas using a GIS overlay. Verify annually that special management practices are still current and/or make operational changes as needed.
DATA SOURCES	COST AND DIFFICULTY
The Forest Manager (TFM); PHP and Bird Studies Canada & MTRI field audits	Low – Bird Studies Canada has consistently completed Bicknell’s Thrush surveys each spring if PHP has pre-commercial thinning activities planned for that summer.
<b>LONG-TERM STRATEGIC MONITORING PROGRAM</b>	
MANAGEMENT OBJECTIVE	Bicknell’s Thrush population recovery
INDICATOR	Population estimates
MONITORING STRATEGY	
<p>Bird Studies Canada annually monitors high elevation bird species through the High Elevation Landbird Program. Since 2002, the Bicknell’s Thrush has been monitored in the Cape Breton Highlands to gather critical information about population status and habitat use. Approximately 20 to 30 routes are monitored each June with the continued goal of monitoring long-term trends of the Bicknell’s Thrush.</p>	
DATA SOURCES	COST AND DIFFICULTY
<ul style="list-style-type: none"> <li>- Bird Studies Canada - Becky Stewart/Holly Lightfoot</li> <li>- Cape Breton Highlands National Park – Matt Smith</li> <li>- International Bicknell’s Thrush Conservation Group (<a href="http://www.bicknellsthrush.org/">http://www.bicknellsthrush.org/</a>)</li> <li>- High Elevation Landbird Report: 10-year Summary, March 2012</li> </ul>	Low – Bird Studies Canada has consistently taken the lead on Bicknell’s Thrush habitat and population research.

## FOREST MANAGEMENT PRESCRIPTION

- Industrial forest stands that support Bicknell's Thrush should remain un-thinned until the trees are no longer at a successional stage that is suitable for nesting, as determined by further research.
- If clearing, construction and/or thinning in Bicknell's Thrush breeding habitat cannot be avoided, activities should be performed outside of the bird breeding season, before June 1st and after July 31st, to prevent the direct destruction of nests, eggs, nestlings, fledglings or adult birds.
- When forest clearing and thinning in Bicknell's Thrush breeding habitat cannot be avoided, patches of intact forest should be left whenever possible. These patches should:
  - cover at least one quarter hectare;
  - be located 20 to 50 metres from the uncut or unthinned edge; and
  - contain intact undisturbed underbrush.

## 2022 MONITORING UPDATE

1. A total of 244 hectares of forest stands was thinned in the BITH Critical Habitat Area as per DNRR approval. Thinning occurred only after August 10<sup>th</sup> when nests are inactive by the end of July.
2. In the summer of 2022, 20 ARUs (all from PHP) were deployed in the working forest area south of Cape Breton Highlands National Park. Bicknell's were detected at two of these sites. Sites were specifically chosen in areas where PCT was planned.
3. PCT will not be undertaken on blocks where BITH was detected, and commercial (mature tree) harvesting will not occur within 300 metres of stands with recent BITH occurrences between April 5<sup>th</sup> and August 28<sup>th</sup>. This is all captured in the *Section 16.1: Endangered Species Act Agreement on Forest Silviculture, Harvesting and Roads within Bicknell's Thrush Federal Critical Habitat* issued by the provincial government to PHP.
4. ARU data will be used to better understand habitat needs of Bicknell's Thrush, so appropriate management decisions can be made regarding forest management.
5. The current population estimate for the Bicknell's Thrush in Canada is between 40,570 and 49,258 birds, and it was previously estimated that approximately 1,200 breed in NB and NS (HELP Report, March 2012).
6. No changes have been made to the special management practices for Bicknell's Thrush as issued by Bird Studies Canada.
7. In March 2012, Bird Studies Canada released a 10-year summary report of their High Elevation Landbird Program. The results for Bicknell's Thrush monitoring found that the sampling intensity was not enough to detect statistically significant trends in population and habitat use.
8. In 2012-13, Bird Studies Canada refined HELP, using a Generalized Random Tessalation Stratified sampling design to randomly select routes and increase sampling intensity in Cape Breton, thus enabling them to meet international, national and regional information needs (HELP Report, March 2012).
9. In July 2020, the federal Bicknell's Thrush Recovery Strategy was finalized. The definition of critical habitat for bicknell's thrush is provided for Nova Scotia, which will help guide habitat management for bicknell's thrush in Nova Scotia.




10. On PHP's Crown license area, there are 66 locations in ACCDC's sensitive species dataset dated July 2020 (no change from 2019). These locations range in observation dates from 1987 to 2013 (see below map).

#### SUPPORTING DOCUMENTS/REFERENCES

Conserving the Bicknell's Thrush: Stewardship and Management Practices for High Elevation Forest, 2009; High Elevation Landbird Program: 10-year Report, March 2012

High Elevation Landbird Program: Annual Report for Cape Breton Highlands National 2013-2014

#### Bicknell's Thrush Critical Habitat Areas



## HCV – Rusty Blackbird Habitat




HCV ATTRIBUTE		Species at Risk – Habitat and Population	
<b>OPERATIONAL MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Maintain and/or enhance Rusty Blackbird habitat		
INDICATOR	Reserve stand structure in Rusty Blackbird habitat		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	
		Monitor implementation of reserve stand structure using field audits. Verify annually that special management practices are still current and/or make operational changes as needed.	
DATA SOURCES	The Forest Manager (TFM); PHP field audits	COST AND DIFFICULTY	
		Low – PHP currently monitors for riparian buffer management on its operational field audits	
<b>LONG-TERM STRATEGIC MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Rusty Blackbird population recovery		
INDICATOR	Population estimates		
MONITORING STRATEGY			
DNRR is responsible for population inventory and studying habitat use.			
DATA SOURCES	NSDNRR	COST AND DIFFICULTY	
		Low to High - Dependent on PHP's required level of involvement	

#### FOREST MANAGEMENT PRESCRIPTION

- PHP implements the Wildlife Habitat and Watercourse Protection Regulations, which is deemed sufficient for Rusty Blackbirds since they tend to occupy forests near the edges of wetlands, bogs, rivers and streams.
- PHP also establishes no harvest buffers (100+ meters wide) around all treed bogs in Cape Breton and specific bogs for Mainland Moose. These are presumed to be beneficial for Rusty Blackbird habitat since buffers larger than 75-100m around wetlands and forested wetlands are suitable or occupied by Rusty Blackbirds.

#### 2022 MONITORING UPDATE

- The population of Rusty Blackbird in Nova Scotia is currently unknown.
- On PHP's Crown license area, there are 69 locations in ACCDC's sensitive species dataset dated May 2022. These locations range in observation dates from 1987 to 2018 (see below map). Deferral of management activities in these locations during the breeding season of May to August may occur depending on data quality, year of original sighting, and input from regional NSDNRR biologists. Deferral of management activities in 2021 have occurred related to at-risk bird known locations.
- PHP is an active partner on a Habitat Stewardship Fund project titled 'Conserving Habitat for Canada Warbler, Olive-sided Flycatcher and Rusty Blackbird in Mainland Nova Scotia'. The main objectives of this project are to build partnerships with landowners and forest managers on breeding habitat conservation as well as develop and test beneficial management practices for the three species in the working forest of mainland Nova Scotia. Project work began in the fall of 2020 and will end in March 2023.
- NSDNRR is currently renewing the provincial species at risk recovery programs by developing new recovery teams for listed species. As of April 2019, a new recovery team was established for all listed birds in the province, including Rusty blackbird. These teams will "set the goals and objectives to address data gaps or threats, monitor the success of recovery and provide recovery-related advice to government to resolve management questions" ([www.novascotia.ca/news](http://www.novascotia.ca/news)).




SUPPORTING DOCUMENTS/REFERENCES

COSEWIC assessment and status report on the Rusty Blackbird *Euphagus carolinus* in Canada (2006)

HCV – Roseate Tern Habitat



HCV ATTRIBUTE		Species at Risk – Habitat
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintain Roseate Tern Habitat	
INDICATOR	Reserve stand structure in Roseate Tern habitat	

<p><b>MONITORING/REPORTING FREQUENCY</b></p> <p>Annual</p>	<p><b>MONITORING STRATEGY</b></p> <p>Maintain a 200-meter buffer zone along the coast at Fisherman’s Harbour. Within this buffer zone, no management will occur.</p>
<p><b>DATA SOURCES</b></p> <p>The Forest Manager (TFM)</p>	<p><b>COST AND DIFFICULTY</b></p> <p>Low – PHP does not conduct forest management activities within the 200-meter buffer zone.</p>
<p><b>FOREST MANAGEMENT PRESCRIPTION</b></p>	
<p>- PHP does not conduct forest management activities within the 200-meter buffer zone at Fisherman’s Harbour.</p> <p>- Other critical habitat for the Roseate Tern is located on offshore islands.</p>	
<p><b>2022 MONITORING UPDATE</b></p>	
<p>A GIS overlay using completed harvest treatment data from 2020 shows that there have been no forest management activities within the 200-meter buffer zone at Fisherman’s Harbour.</p>	
<p style="text-align: center;"><b>Roseate Tern Protected Habitat - Fishermans Harbour, Guysborough County</b></p> 	
<p><b>SUPPORTING DOCUMENTS/REFERENCES</b></p>	
<p>Roseate Tern Recovery Strategy 2006</p>	

## HCV – Olive-Sided Flycatcher Habitat



HCV ATTRIBUTE		Species at Risk – Habitat	
<b>OPERATIONAL MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Olive-sided Flycatcher Habitat		
INDICATOR	Reserve stand structure in Olive-sided flycatcher habitat		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	Monitor COSEWIC and NSDNRR's websites for recovery strategies, actions plans, and/or special management practices developed for this species.
DATA SOURCES	ACCDC	COST AND DIFFICULTY	Low – PHP does not yet implement SMP's
<b>FOREST MANAGEMENT PRESCRIPTION</b>			
<p>The habitat characteristics of olive-sided flycatcher are minimally impacted by forestry activities due to where they typically inhabit. The species prefers tall snags or residual trees for perching and foraging. PHP leaves snags throughout its operations and the presence of tall trees can be found in several PHP silviculture treatments (e.g. single selection, group selection, partial cuts, shelterwoods, patch cuts, red spruce management). PHP also provides habitat features such as forest edges, openings, and clearcuts, and spruce and fir trees are preferred nest sites. Understory thinning may also enhance foraging opportunities. Currently, no special management practices exist for this species, but once developed PHP will implement them as applicable to forest management. If an active nest is located during regular operational activities, the activity will be stopped and the local DNRR Wildlife Biologist will be notified so appropriate measures can be implemented.</p>			
<b>2022 MONITORING UPDATE</b>			
A recovery strategy for Olive-sided Flycatcher was finalized by COSEWIC in March 2016.			

Regarding critical habitat for this species, the recovery strategy states:

*“The available information is not adequate to enable the identification of critical habitat at the landscape scale for the following reasons:*

- *There is a lack of understanding and data to indicate the suitable configuration of important landscape biophysical attributes.*
- *Habitat requirements may vary across the range of the species. Management units (i.e., geographic units within which critical habitat would be managed) need to be identified in such a way to best reflect variation in habitat use.*
- *There is a lack of data related to Olive-sided Flycatcher presence and abundance in large portions of its range. Without this information any model used to predict critical habitat with current data may have a limited ability to do so in these areas.*
- *For Olive-sided Flycatcher, it is unknown whether certain habitats with specific biophysical attributes may be functionally more important than others. For example, specific habitats may have greater densities of individuals or pairs and/or result in higher reproductive success. There are few data regarding the relative importance of suitable habitat types for Olive-sided Flycatcher population numbers and indices of habitat quality.*
- *The relationships between anthropogenic disturbance and habitat quality are poorly known. A better understanding of these relationships is needed to ensure sufficient suitable habitat is available for Olive-sided Flycatcher and to identify at what scale and intensity activities would be likely to destroy the critical habitat.*


*A Schedule of Studies (Table 4) has been developed to provide the information necessary to identify the critical habitat that will be sufficient to meet the population and distribution objectives. The identification of critical habitat will be included in a revised recovery strategy or an action plan.”*

- On PHP's Crown license area, there are 225 locations in ACCDC's sensitive species dataset dated May 2022. All locations range in observation dates from 1987 to 2020 (see below map). Deferral of management activities during the breeding season of May to August may occur depending on data quality, year of original sighting, and input from regional NSDNRR biologists. Deferral of management activities in 2021 have occurred related to at-risk bird known locations.

- PHP is an active partner on a Habitat Stewardship Fund project titled 'Conserving Habitat for Canada Warbler, Olive-sided Flycatcher and Rusty Blackbird in Mainland Nova Scotia'. The main objectives of this project are to build partnerships with landowners and forest managers on breeding habitat conservation as well as develop and test beneficial management practices for the three species in the working forest of mainland Nova Scotia. Project work began in the fall of 2020 and will end in March 2023.

- NSDNRR is currently renewing the provincial species at risk recovery programs by developing new recovery teams for listed species. As of April 2019, a new recovery team was established for all listed birds in the province, including Olive-sided flycatcher. These teams will “set the goals and objectives to address data gaps or threats, monitor the success of

recovery and provide recovery-related advice to government to resolve management questions” ([www.novascotia.ca/news](http://www.novascotia.ca/news)).



SUPPORTING DOCUMENTS/REFERENCES

COSEWIC Website; ACCDC Data

HCV – Eastern Whip-Poor-Will Habitat



HCV ATTRIBUTE		Species at Risk – Habitat
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Eastern Whip-poor-will Habitat	
INDICATOR	Reserve stand structure in Eastern Whip-poor-will habitat	



<p>MONITORING/REPORTING FREQUENCY</p> <p>Annual</p>	<p>Monitor COSEWIC and NSDNRR's websites for recovery strategies, actions plans, and/or special management practices developed for this species.</p>
<p>DATA SOURCES</p> <p>ACCDC</p>	<p>COST AND DIFFICULTY</p> <p>Low – PHP does not yet implement SMP's</p>
<p>FOREST MANAGEMENT PRESCRIPTION</p>	
<p>The habitat characteristics of whip-poor-will are minimally impacted by forestry activities due to where they typically inhabit. PHP creates forest edges and openings through active management, as well as even-aged stands that can contain well-spaced trees. This species may also use barrens or regenerating forests following a disturbance, which are present across PHP's operating area. Currently, no special management practices exist for this species, but once developed PHP will implement them as applicable to forest management. If an active nest is located during regular operational activities, the activity will be stopped and the local DNRR Wildlife Biologist will be notified so appropriate measures can be implemented.</p>	
<p>2022 MONITORING UPDATE</p>	
<p>A COSEWIC recovery strategy was finalized for this species in 2018.</p> <p>There are no locations of Eastern Whip-poor-will in ACCDC's May 2022 dataset.</p> <p>- NSDNRR is currently renewing the provincial species at risk recovery programs by developing new recovery teams for listed species. As of April 2019, a new recovery team was established for all listed birds in the province, including Eastern Whip-poor-will. These teams will "set the goals and objectives to address data gaps or threats, monitor the success of recovery and provide recovery-related advice to government to resolve management questions" (<a href="http://www.novascotia.ca/news">www.novascotia.ca/news</a>).</p>	
<p>SUPPORTING DOCUMENTS/REFERENCES</p>	
<p>COSEWIC Website; ACCDC Data</p>	

## HCV – Eastern Wood Peewee Habitat



HCV ATTRIBUTE		Species at Risk – Habitat	
<b>OPERATIONAL MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Eastern Wood Peewee Habitat		
INDICATOR	Reserve stand structure in Eastern wood peewee habitat		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	Monitor COSEWIC and NSDNRR's websites for recovery strategies, actions plans, and/or special management practices developed for this species.
DATA SOURCES	ACCDC	COST AND DIFFICULTY	Low – PHP does not yet implement SMP's
<b>FOREST MANAGEMENT PRESCRIPTION</b>			
<p>The Eastern wood peewee can be impacted by forest management activities since this species of bird prefers mature and intermediate age stands of deciduous and mixed forests. However, PHP manages the forest management area by creating a range of age classes through forest modeling, long-term planning, and operational planning. Also, PHP manages deciduous and mixed forest stands with a variety of harvest treatments that can still maintain adequate forest structure and large mature trees for nesting and perching (e.g. single selection, group selection, partial cuts, shelterwoods, patch cuts).</p> <p>Forestry practices that maintain large tracts of intermediate aged forest with closed canopy and limited clear cuts (less than 10 ha) along with thinning to remove mature trees and large-diameter woody growth should provide adequate habitat for Eastern Wood-Peewees (Stauffer and Best 1980, Crawford et al. 1981).</p>			


Currently, no special management practices exist for this species, but once developed PHP will implement them as applicable to forest management. If an active nest is located during regular operational activities, the activity will be stopped and the local DNRR Wildlife Biologist will be notified so appropriate measures can be implemented.

#### 2022 MONITORING UPDATE

Currently, there is no recovery strategy, action plan and/or special management practices issued by either COSEWIC or NSDNRR.

On PHP's Crown license area, there are 55 locations in ACCDC's sensitive species dataset from May 2022. Observation dates range from 1986 to 2020 (see below map). Deferral of management activities during the breeding season of May to August may occur depending on data quality, year of original sighting, and input from regional NSDNRR biologists. Deferral of management activities in 2021 have occurred related to at-risk bird known locations.

- NSDNRR is currently renewing the provincial species at risk recovery programs by developing new recovery teams for listed species. As of April 2019, a new recovery team was established for all listed birds in the province, including Eastern Wood Pewee. These teams will "set the goals and objectives to address data gaps or threats, monitor the success of recovery and provide recovery-related advice to government to resolve management questions" ([www.novascotia.ca/news](http://www.novascotia.ca/news)).



SUPPORTING DOCUMENTS/REFERENCES

COSEWIC Website; ACCDC Data

## HCV – Canada Warbler Habitat



HCV ATTRIBUTE		Species at Risk – Habitat
<b>OPERATIONAL MONITORING PROGRAM</b>		
MANAGEMENT OBJECTIVE	Canada Warbler Habitat	
INDICATOR	Reserve stand structure in Canada warbler habitat	
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY	
Annual	Monitor COSEWIC and NSDNRR's websites for recovery strategies, actions plans, and/or special management practices developed for this species.	
DATA SOURCES	COST AND DIFFICULTY	
ACCDC	Low – PHP does not yet implement SMP's	
<b>FOREST MANAGEMENT PRESCRIPTION</b>		
<p>The habitat characteristics of Canada warbler are minimally impacted by forestry activities due to where they typically inhabit. PHP creates regenerating stand structures and forest edge through active management, which is preferred by this species, but also avoid steep slope areas, ravines, swamps, and bogs. The provision of stumps and coarse woody debris left by PHP is also believed to create understory conditions preferred by the Canada warbler. Furthermore, PHP does not contribute to habitat loss by converting swamp forests to agricultural land. Currently, no special management practices exist for this species, but once developed PHP will implement them as applicable to forest management. If an active nest is located during regular operational activities, the activity will be stopped and the local DNRR Wildlife Biologist will be notified so appropriate measures can be implemented.</p>		
<b>2022 MONITORING UPDATE</b>		
<p>A COSEWIC recovery strategy was finalized for Canada Warbler in March 2016.</p> <p>Regarding critical habitat for this species, the recovery strategy states:</p>		

*“The available information is not adequate to identify critical habitat at a landscape scale for the following reasons:*


- There is a lack of understanding and data to indicate the appropriate configuration of important landscape biophysical attributes.*
- Habitat requirements may vary across the range of the species. Management units (i.e., geographic units within which critical habitat would be managed) need to be identified in such a way to best reflect variation in habitat use and management patterns.*
- There is a lack of data related to Canada Warbler presence and abundance in large portions of its range. Without this information any model used to predict critical habitat with current data may have a limited ability to do so in these areas.*
- For Canada Warbler, it is unclear whether certain habitats with specific biophysical attributes may be functionally more important than others. For example, specific habitats may have greater densities of individuals or pairs and/or result in higher reproductive success.*
- The relationships between anthropogenic disturbance and habitat quality are poorly known. A better understanding of these relationships is needed to ensure sufficient suitable habitat is available for Canada Warbler and to identify at what scale and intensity activities would be likely to destroy the critical habitat.*

*A Schedule of Studies has been developed to provide the information necessary to identify the critical habitat that will be sufficient to meet the population and distribution objectives. The identification of critical habitat will be included in a revised recovery strategy or an action plan.*

On PHP's Crown license area, there are 159 locations in ACCDC's sensitive species dataset from May 2022. Observation dates range from 1987 to 2021 (see below map). Deferral of management activities during the breeding season of May to August may occur depending on data quality, year of original sighting, and input from regional NSDNRR biologists. Deferral of management activities in 2021 have occurred related to at-risk bird known locations.

- PHP is an active partner on a Habitat Stewardship Fund project titled 'Conserving Habitat for Canada Warbler, Olive-sided Flycatcher and Rusty Blackbird in Mainland Nova Scotia'. The main objectives of this project are to build partnerships with landowners and forest managers on breeding habitat conservation as well as develop and test beneficial management practices for the three species in the working forest of mainland Nova Scotia. Project work began in the fall of 2020 and will end in March 2023.

- NSDNRR is currently renewing the provincial species at risk recovery programs by developing new recovery teams for listed species. As of April 2019, a new recovery team was established for all listed birds in the province, including Canada Warbler. These teams will “set the goals and objectives to address data gaps or threats, monitor the success of recovery and provide recovery-related advice to government to resolve management questions” ([www.novascotia.ca/news](http://www.novascotia.ca/news)).



SUPPORTING DOCUMENTS/REFERENCES

COSEWIC Website; ACCDC Data

## HCV – CHIMNEY SWIFT Habitat



HCV ATTRIBUTE		Species at Risk – Habitat	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Chimney Swift Habitat		
INDICATOR	Reserve stand structure in Chimney swift habitat		
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY		
Annual	Monitor COSEWIC and NSDNRR's websites for recovery strategies, actions plans, and/or special management practices developed for this species.		
DATA SOURCES	COST AND DIFFICULTY		
ACCDC	Low – PHP does not yet implement SMP's		
FOREST MANAGEMENT PRESCRIPTION			
<p>The Chimney swift can be impacted by forest management activities since this species of bird may nest in wooded areas with large diameter trees. Current management practices such as group and single selection implemented by PHP in shade-tolerant hardwood stands provides large-diameter trees as part of the forest cover for roosting and nest sites. Additionally, the old-growth forest protocol implemented by NSDNRR is increasing the amount of protected old-growth mature and climax hardwood forests, which is beneficial for the Chimney Swift as well as many other bird species. Currently, there are no special management practices identified for forest managers regarding Chimney swift habitat. Additionally, feeding and nesting habitat relies heavily on urban and suburban areas where there is an abundance of chimneys for nesting, so PHP believes it currently has a low impact on Chimney swift populations.</p>			




## 2022 MONITORING UPDATE

Currently, there is no recovery strategy, action plan and/or special management practices issued by either COSEWIC or NSDNRR.

On PHP's Crown license area, there are 16 locations in ACCDC's sensitive species dataset from May 2022. These locations range in observation dates from 1986 to 2011 (see below map). Deferral of management activities during the breeding season of May to August may occur depending on data quality, year of original sighting, and input from regional NSDNRR biologists.

- NSDNRR is currently renewing the provincial species at risk recovery programs by developing new recovery teams for listed species. As of April 2019, a new recovery team was established for all listed birds in the province, including Chimney swift. These teams will "set the goals and objectives to address data gaps or threats, monitor the success of recovery and provide recovery-related advice to government to resolve management questions" ([www.novascotia.ca/news](http://www.novascotia.ca/news)).



## SUPPORTING DOCUMENTS/REFERENCES

COSEWIC Website; ACCDC Data

## HCV – COMMON NIGHTHAWK Habitat



HCV ATTRIBUTE		Species at Risk – Habitat	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Common Nighthawk Habitat		
INDICATOR	Reserve stand structure in Common nighthawk habitat		
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY		
Annual	Monitor COSEWIC and NSDNRR's websites for recovery strategies, actions plans, and/or special management practices developed for this species.		
DATA SOURCES	COST AND DIFFICULTY		
ACCDC	Low – PHP does not yet implement SMP's		
FOREST MANAGEMENT PRESCRIPTION			
<p>The Common Nighthawk prefers some habitats where PHP does not operate such as rocky areas, sandy areas, and wetlands. However, they do prefer open wooded areas, which PHP does create through its forest management (e.g. clearcuts, partial cuts, shelterwoods, selection cuts). The Common Nighthawk Recovery Strategy lists a variety of threats including changes in natural processes, climate and natural disasters, accidental mortality, pollution, exotic or invasive species, and habitat loss or degradation. Types of habitat loss include change in roof construction and materials, residential and commercial development, agriculture, and logging and wood harvesting. It is currently unknown if logging and wood harvesting causes a significant severity to populations with a low causal certainty that there is a high degree of evidence linked to the threat of logging. Regardless, PHP provides habitat conditions for the Common Nighthawk through its forest management and does not apply pesticides which can impact foraging areas near nest patches.</p>			

## 2022 MONITORING UPDATE

COSEWIC finalized a recovery strategy for the Common Nighthawk in March 2016.

Regarding critical habitat for this species, the recovery strategy states:

*“The available information is not adequate to enable the identification of critical habitat for the following reasons:*

- *There is a lack of understanding and data to indicate the appropriate biophysical attributes required by the species and their configuration at a landscape scale.*
- *Habitat requirements may vary across the range of the species. Management units (i.e., geographic units within which critical habitat would be managed) need to be identified in such a way to best reflect variation in habitat use and land planning processes.*
- *There is a lack of data related to presence, site usage where detected (e.g., foraging, roosting, defending a territory, nesting, transiting), and abundance in large portions of the species’ range and the northern limit of the species’ range is unknown. Without this information any model used to predict critical habitat with current data may have a limited ability to do so.*
- *For Common Nighthawk, it is unknown whether certain habitats with specific biophysical attributes may be functionally more important than others. For example, specific habitats may have greater densities of individuals or pairs and/or result in higher reproductive success.*
- *The relationships between anthropogenic disturbance and habitat quality are poorly known. A better understanding of these relationships is needed to ensure sufficient suitable habitat is currently available for Common Nighthawk and to identify at what scale and intensity activities would be likely to destroy critical habitat.*


*Locating nests is difficult and determining general nesting locations is problematic using typical point-count survey methodology. Common Nighthawks defend a large area and their foraging habitats can be separated from nest sites by many kilometers, so it is not possible to determine how an individual is using the habitat where it is detected (e.g., foraging, defending a territory, transiting). Furthermore, traditional point-count survey methodology in the morning is not appropriate for this crepuscular species (Government of Alberta 2013; Saskatchewan Ministry of Environment 2014).*

*A schedule of studies has been developed to provide the information necessary to identify the critical habitat that will be sufficient to meet the population and distribution objectives. The identification of critical habitat will be included either in a revised recovery strategy or an action plan.”*

On PHP’s Crown license area, there are 43 locations in ACCDC’s sensitive species dataset from May 2022. Observation dates range from 2006 to 2021 (see below map). The location documented in 2019 was not near any on-going or planned harvest, however a planting was undertaken in a stand nearby. Deferral of management activities during the breeding season of May to August may occur depending on data quality, year of original sighting, and input from regional NSDNRR biologists.

- NSDNRR is currently renewing the provincial species at risk recovery programs by developing new recovery teams for listed species. As of April 2019, a new recovery team was established for all listed birds in the province, including Common nighthawk. These teams

will “set the goals and objectives to address data gaps or threats, monitor the success of recovery and provide recovery-related advice to government to resolve management questions” ([www.novascotia.ca/news](http://www.novascotia.ca/news)).



#### SUPPORTING DOCUMENTS/REFERENCES

COSEWIC Website; ACCDC Data

## HCV – WOOD THRUSH Habitat



HCV ATTRIBUTE		Species at Risk – Habitat	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Wood Thrush Habitat		
INDICATOR	Reserve stand structure in Wood thrush habitat		
MONITORING/REPORTING FREQUENCY		MONITORING STRATEGY	
Annual		Monitor COSEWIC and NSDNRR's websites for recovery strategies, actions plans, and/or special management practices developed for this species.	
DATA SOURCES		COST AND DIFFICULTY	
ACCDC		Low – PHP does not yet implement SMP's	
FOREST MANAGEMENT PRESCRIPTION			
<p>Currently, there are no required management practices for Wood Thrush in Nova Scotia or Canada. Regardless, given the preferred breeding habitat of Wood Thrush in mature deciduous and mixed-wood forests, PHP believes its uneven-aged and mixed-wood forest management techniques in these forest types do not greatly impact the breeding requirements of the Wood Thrush. The COSEWIC 2012 report supports this notion by stating that “the species is relatively tolerant of forest management activities that are conducted on a small spatial scale (i.e. single-tree, group selection cuts, uneven-age forest management, selective removal of mature trees). The report further states that Sugar Maple and American Beech are preferred species for nesting. PHP manages tolerant Sugar Maple stands using only single or group selection depending on tree quality. American Beech is present throughout the forest management area and pure stands are not managed, but if found dispersed throughout a hardwood stand, it is managed as necessary to meet the forest management prescription. Additionally, PHP does not apply herbicides in its forest</p>			

management area, which allows for the continued natural growth of deciduous trees and shrubs in forest stands.

#### 2022 MONITORING UPDATE

Currently, there is no recovery strategy, action plan and/or special management practices issued by either COSEWIC or NSDNRR.

On PHP's Crown license area, there are 2 locations in ACCDC's sensitive species dataset from May 2022 (no change from 2019). These locations range in observation dates from 2008 to 2009 (see below map). Deferral of management activities during the breeding season of May to August may occur depending on data quality, year of original sighting, and input from regional NSDNRR biologists.



#### SUPPORTING DOCUMENTS/REFERENCES

COSEWIC Website; ACCDC Data

## HCV – EVENING GROSBEEK Habitat




HCV ATTRIBUTE		Species at Risk – Habitat	
<b>OPERATIONAL MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Evening Grosbeak Habitat		
INDICATOR	Reserve stand structure in Evening Grosbeak habitat		
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY		
Annual	Monitor COSEWIC and NSDNRR's websites for recovery strategies, actions plans, and/or special management practices developed for this species.		
DATA SOURCES	COST AND DIFFICULTY		
ACCDC	Low – PHP does not yet implement SMP's		
<b>FOREST MANAGEMENT PRESCRIPTION</b>			
<p>Currently, there are no required management practices for Evening Grosbeak in Nova Scotia or Canada. Regardless, given the preferred breeding habitat of Evening Grosbeak in large mature and old mixedwood forest stands, PHP may have an impact on the breeding success of Evening Grosbeak. Fluctuations of spruce budworm populations are likely a key factor in fluctuations of Evening Grosbeak populations. Other known threats include window strike mortality in winter, reduction of mature and old-growth mixedwood forests, and road collision mortality. On a large landscape scale, PHP manages the forest land-base to ensure a diversity of stand types and ages, which includes mature and old mixedwood stands. Currently, PHP is actively involved in assessing mature and old forest stands for old-growth protection under the provincial Old Forest Policy. Where stands meet the Old Forest Policy stand definition, measures are put in place to allocate the stand as protected under the Old Forest Policy which will benefit Evening Grosbeak habitat. In 2019, 14 stands that were assessed for old-growth characteristics were confirmed as old-growth and are now protected under the provincial Old Forest Policy. The COSEWIC Assessment and Status Report for Evening Grosbeak (2016) has identified the loss, alteration and fragmentation of breeding habitat due to commercial logging as a low threat overall.</p>			

**2022 MONITORING UPDATE**

Currently, there is no recovery strategy, action plan and/or special management practices issued by either COSEWIC or NSDNRR.

On PHP's Crown license area, there are 98 locations in ACCDC's sensitive species dataset from May 2022. These locations range in observation dates from 1986 to 2021 (see below map). Deferral of management activities during the breeding season of May to September may occur depending on data quality, year of original sighting, and input from regional NSDNRR biologists.

**SUPPORTING DOCUMENTS/REFERENCES**

COSEWIC Website; ACCDC Data



## HCV – BLACK-FOAM LICHEN Habitat



HCV ATTRIBUTE		Species at Risk – Habitat
<b>OPERATIONAL MONITORING PROGRAM</b>		
MANAGEMENT OBJECTIVE	Black-foam Lichen Habitat	
INDICATOR	Reserve stand structure in Black-foam lichen habitat	
MONITORING/REPORTING FREQUENCY	Annual	<b>MONITORING STRATEGY</b>  Confirm with NS Department of Natural Resources & Renewables, NS Environment, Atlantic Canada Conservation Data Centre, and Mersey Tobeatic Research Institute if any new locations of black-foam lichen have been discovered on PHP's Crown lease.
DATA SOURCES	ACCDC	<b>COST AND DIFFICULTY</b>  Low – PHP does not yet implement SMP's
<b>FOREST MANAGEMENT PRESCRIPTION</b>		
<p>In May 2018, the NSDNRR released new Special Management Practices for At-Risk Lichens for several lichen species including: boreal felt lichen, vole ears, Hibernia jellyskin lichen, powdered moon lichen, eastern waterfan, wrinkled shingle lichen, frosted glass-whiskers, black-foam lichen, blue felt lichen, and poor-man's shingles lichen. The SMP is triggered by areas of overlap between planned forest management activities and the modelled lichen habitat for boreal felt lichen. Where there is overlap, a lichen survey is conducted and if any of the listed species in the SMP is found, a protection zone is established around the site. The sizes of these protection zones vary depending on the species. There are also restrictions related to road construction and maintenance.</p> <p>For black-foam lichen, a 100 metre radius protection zone is established around each site. This zone does not permit harvesting or silviculture, mineral exploration drill sites, or new road or trail construction (unless in exceptional circumstances and under approval by DNRR). Existing road maintenance is permitted subject to review and conditions.</p>		

<b>2022 MONITORING UPDATE</b>
No new locations of black-foam lichen have been found on PHP's Crown lease. Currently, there is still only one known location of this lichen in the seven eastern counties where PHP operates and it is in the Cape Breton Highlands National Park. An expert lichenologist also verified that this lichen is still most commonly found in southwest Nova Scotia.
<b>SUPPORTING DOCUMENTS/REFERENCES</b>
NS Department of Natural Resources & Renewables, NS Environment, ACCDC, MTRI

### HCV – LITTLE BROWN MYOTIS Habitat



<b>HCV ATTRIBUTE</b> <b>Species at Risk – Habitat</b>	
<b>OPERATIONAL MONITORING PROGRAM</b>	
<b>MANAGEMENT OBJECTIVE</b>	Little Brown Myotis Habitat
<b>INDICATOR</b>	Reserve stand structure in Little brown myotis habitat
<b>MONITORING/REPORTING FREQUENCY</b>  Annual	<b>MONITORING STRATEGY</b>  Monitor Environment Canada's work on the development of beneficial management practices for the forest industry. Monitor Crown contractor audits to verify that unmerchantable trees, such as snags, wolf trees, and cavity trees, are being retained on site. Monitor ACCDC data for any known locations.
<b>DATA SOURCES</b>  ACCDC	<b>COST AND DIFFICULTY</b>  Low – PHP does not yet implement SMP's

**FOREST MANAGEMENT PRESCRIPTION**

Currently in Nova Scotia, there are no best forest management practices required for bats. Regarding wolf trees which are important for roosting, the NS Forest Wildlife Guidelines of 1988, which is now a Crown land policy, recommends that snags, wolf trees, and cavity trees be left on harvest sites as much as possible. Most often, wolf trees are so large and difficult to harvest because of many branches, and have low economic value, that PHP leaves on site. PHP is currently managing the forest in a variety of ways that benefit bat habitat needs, based on a 2006 report called "Forest Management & Bats" by Bat Conservation International which lists a variety of forest management activities that can support bat habitat needs. The most beneficial stand-level best management practices include leaving the best roost sites with woodpecker holes, rot holes, cracks and splits. Retention of snags and large mature trees may also increase the suitability of forested habitat for most bat species. Additionally, NSDNRR is aware of locations in Nova Scotia where overwintering habitat such as caves, abandoned mines, and wells are used by many bat species. Restrictions to forest management activities near these locations are provided by NSDNRR during the approval process for operational plans.

**2022 MONITORING UPDATE**

Currently, there are no beneficial management practices developed for the forest industry. A large colony of approximately 300 females was recently found (July 2016) in Nova Scotia. Due to the highly sensitive nature of bat populations, its location was not made publicly available. However, it was confirmed to PHP by a DNRR management executive that the colony was not found on PHP's Crown lease. The 2022 Crown operations audits show that unmerchantable trees were left on harvest sites, which could serve as roost sites. ACCDC does have two locations for this species in the May 2022 dataset and both are in the Plaster Bat Cave which is located in a protected nature reserve.

**SUPPORTING DOCUMENTS/REFERENCES**

NS Department of Natural Resources & Renewables, Environment Canada

## HCV – TRI-COLORED BAT Habitat



HCV ATTRIBUTE		Species at Risk – Habitat	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Tri-colored Bat Habitat		
INDICATOR	Reserve stand structure in Tri-colored bat habitat		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	Monitor Environment Canada’s work on the development of beneficial management practices for the forest industry. Monitor Crown contractor audits to verify that unmerchantable trees, such as snags, wolf trees, and cavity trees, are being retained on site. Monitor ACCDC data for any known locations.
DATA SOURCES	ACCDC	COST AND DIFFICULTY	Low – PHP does not yet implement SMP’s
FOREST MANAGEMENT PRESCRIPTION			
<p>Currently in Nova Scotia, there are no best forest management practices required for bats. Regarding wolf trees which are important for roosting, the NS Forest Wildlife Guidelines of 1988, which is now a Crown land policy, recommends that snags, wolf trees, and cavity trees be left on harvest sites as much as possible. Most often, wolf trees are so large and difficult to harvest because of many branches, and have low economic value, that PHP leaves on site. PHP is currently managing the forest in a variety of ways that benefit bat habitat needs, based on a 2006 report called “Forest Management &amp; Bats” by Bat Conservation International which lists a variety of forest management activities that can support bat habitat needs. The most beneficial stand-level best management practices include leaving the best roost sites with woodpecker holes, rot holes, cracks and splits. Retention of snags and large mature trees may also increase the suitability of forested habitat for most bat species. Additionally, NSDNRR is aware of locations in Nova Scotia where overwintering habitat such as caves,</p>			

abandoned mines, and wells are used by many bat species. Restrictions to forest management activities near these locations are provided by NSDNRR during the approval process for operational plans.

#### 2022 MONITORING UPDATE

Currently, there are no beneficial management practices developed for the forest industry. The 2022 Crown operations audits show that unmerchantable trees were left on harvest sites. There are no locations of this bat species in the ACCDC dataset available to PHP.

#### SUPPORTING DOCUMENTS/REFERENCES

NS Department of Natural Resources & Renewables, Environment Canada

### HCV – NORTHERN MYOTIS Habitat



HCV ATTRIBUTE		Species at Risk – Habitat
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Northern Myotis Habitat	
INDICATOR	Reserve stand structure in Northern myotis habitat	
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY
		Monitor Environment Canada's work on the development of beneficial management practices for the forest industry. Monitor Crown contractor audits to verify that unmerchantable trees, such as snags, wolf trees, and cavity trees, are being retained on site. Monitor ACCDC data for any known locations.


<p>DATA SOURCES</p> <p>ACCDC</p>	<p>COST AND DIFFICULTY</p> <p>Low – PHP does not yet implement SMP's</p>
<p>FOREST MANAGEMENT PRESCRIPTION</p>	
<p>Currently in Nova Scotia, there are no best forest management practices required for bats. Regarding wolf trees which are important for roosting, the NS Forest Wildlife Guidelines of 1988, which is now a Crown land policy, recommends that snags, wolf trees, and cavity trees be left on harvest sites as much as possible. Most often, wolf trees are so large and difficult to harvest because of many branches, and have low economic value, that PHP leaves on site. PHP is currently managing the forest in a variety of ways that benefit bat habitat needs, based on a 2006 report called "Forest Management &amp; Bats" by Bat Conservation International which lists a variety of forest management activities that can support bat habitat needs. The most beneficial stand-level best management practices include leaving the best roost sites with woodpecker holes, rot holes, cracks and splits. Retention of snags and large mature trees may also increase the suitability of forested habitat for most bat species. Additionally, NSDNRR is aware of locations in Nova Scotia where overwintering habitat such as caves, abandoned mines, and wells are used by many bat species. Restrictions to forest management activities near these locations are provided by NSDNRR during the approval process for operational plans.</p>	
<p>2022 MONITORING UPDATE</p>	
<p>Currently, there are no beneficial management practices developed for the forest industry. The 2022 Crown operations audits show that unmerchantable trees were left on harvest sites. There are no locations of this bat species in the ACCDC dataset available to PHP.</p>	
<p>SUPPORTING DOCUMENTS/REFERENCES</p>	
<p>NS Department of Natural Resources &amp; Renewables, Environment Canada</p>	

## HCV – New Jersey Rush Habitat



HCV ATTRIBUTE		Species at Risk – Habitat	
<b>OPERATIONAL MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Maintain New Jersey Rush Habitat		
INDICATOR	Administratively protect New Jersey Rush habitat identified in NSDNRR's Significant Habitat database and the Atlantic Coastal Plain Flora database		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	Monitor annual harvest operations to ensure New Jersey Rush habitat is administratively protected from all forest management activities.
DATA SOURCES	The Forest Manager (TFM)	COST AND DIFFICULTY	Low – PHP does not conduct forest management activities within New Jersey Rush habitat
<b>FOREST MANAGEMENT PRESCRIPTION</b>			
- PHP does not conduct forest management activities within New Jersey Rush habitat identified in NSDNRR's Significant Habitat database and/or the Atlantic Coastal Plain Flora database			
<b>2022 MONITORING UPDATE</b>			
Known locations of New Jersey Rush are protected through the Atlantic Coastal Plain Flora dataset which is provided by NSDNRR.			
<b>SUPPORTING DOCUMENTS/REFERENCES</b>			
Recovery Strategy and Management Plan for Multiple Species of Atlantic Coastal Plain Flora 2010; ACCDC Dataset			

**Protected Areas for Atlantic Coastal Plain Flora**



**HCV – Boreal Felt Lichen Occurrences**



<b>HCV ATTRIBUTE</b>		<b>Species at Risk – Habitat and Population</b>	
<b>OPERATIONAL MONITORING PROGRAM</b>			
<b>MANAGEMENT OBJECTIVE</b>	Protect identified locations of Boreal Felt Lichen		
<b>INDICATOR</b>	Administratively protect identified locations of Boreal Felt Lichen by establishing 100-meter buffer around site		
<b>MONITORING/REPORTING FREQUENCY</b>	<b>MONITORING STRATEGY</b>		



Annual	Monitor annual harvest operations to identify areas needing Boreal Felt Lichen presence/absence surveys prior to active operations. Locations of Boreal Felt Lichen are buffered by 100 meters and excluded from forest management activities.
DATA SOURCES	COST AND DIFFICULTY
The Forest Manager (TFM); Boreal Felt Lichen Potential Habitat Layer	Moderate – PHP financially contributes annually to Boreal Felt Lichen surveys. Surveys are conducted by an expert lichenologist.
<b>LONG-TERM STRATEGIC MONITORING PROGRAM</b>	
MANAGEMENT	Boreal Felt Lichen population recovery
OBJECTIVE	
INDICATOR	Population estimates
MONITORING STRATEGY	
DNRR is responsible for population inventory and studying habitat use.	
DATA SOURCES	COST AND DIFFICULTY
NSDNRR; NSDOE	Low to High - Dependent on PHP's required level of involvement
<b>FOREST MANAGEMENT PRESCRIPTION</b>	
<p>In May 2018, the NSDNRR released new Special Management Practices for At-Risk Lichens for several lichen species including: boreal felt lichen, vole ears, Hibernia jellyskin lichen, powdered moon lichen, eastern waterfan, wrinkled shingle lichen, frosted glass-whiskers, black-foam lichen, blue felt lichen, and poor-man's shingles lichen. The SMP is triggered by areas of overlap between planned forest management activities and the modelled lichen habitat for boreal felt lichen. Where there is overlap, a lichen survey is conducted and if any of the listed species in the SMP is found, a protection zone is established around the site. The sizes of these protection zones vary depending on the species. There are also restrictions related to road construction and maintenance.</p> <p>For boreal felt lichen, a 500 metre radius special management zone is established around each site. Within this zone is a protected zone (200 m out from occurrence) and a restricted zone (300 m from edge of protected zone). In the protected zone, harvesting or silviculture, mineral exploration drill sites, or new road or trail construction (unless in exceptional</p>	

circumstances and under approval by DNRR) are permitted. Existing road maintenance is permitted subject to review and conditions.


In the restricted zone, partial harvesting is favoured by using the 'restoration' pathway in DNRR's Forest Management Guides (FMG). If the FMG determines that a clearcut is the appropriate treatment, areas of clearcut shall not exceed 10 ha and the distance between clearcuts must not be less than 100 metres. Buffers are also established around forested wetlands and provisions are made based on the state of regenerating development within the zone.

#### 2022 MONITORING UPDATE

- In 2022, there were 36 planned harvest sites surveyed where Boreal Felt Lichen potential habitat was identified. BFL was not found during these surveys.
- Since 2008, PHP has worked with the Mersey Tobeatic Research Institute to conduct Boreal Felt Lichen surveys. Prior to these surveys, there were three known locations of Boreal Felt Lichen in Nova Scotia. Since MTRI's surveys began for PHP, the number of known locations has increased to 526 in PHP's forest management area.
- NSDNRR recently published a paper entitled "Forest Harvesting Impacts on Mortality of an Endangered Lichen at the Landscape and Stand Scales". This paper supports protection buffers around known Boreal Felt Lichen sites to maintain the micro-climate around the site. NSDNRR is also currently working on a habitat supply research paper. NSDNRR will also be working on improving the predicted habitat model for Boreal Felt Lichen. NSDNRR will also be monitoring how different buffer widths affect microclimate using iButtons (micro-climate data loggers). Currently, there are about 24 iButtons in the field which are being used to assess the variation between and within stands. This information will help determine the sample size needed for a future buffer width study.
- PHP is a supporting partner on a recently approved Dalhousie University PhD research project by Mitacs. The project titled "Disturbance thresholds and factors influencing community dynamics of epiphytic cyanolichens in Nova Scotia, with an emphasis on rare and at-risk species". The project began in January 2018 and is expected to last two years. The researcher is expected to spend at least 35% of his time on PHP Crown licensed lands each year for site selection and planning, field work, and advisory meetings. Preliminary results indicate a higher proportion of gastropod grazing on lichens near clearcut edges than in the forest interior. There also appears to be a greater mix of non-native slug species on sites that are closer to forest roads, indicating anthropogenic vectors of spread.

#### SUPPORTING DOCUMENTS/REFERENCES

Boreal Felt Lichen Recovery Strategy; Boreal Felt Lichen Recovery Team



### HCV – Vole Ears Lichen Occurrences



HCV ATTRIBUTE		Species at Risk – Habitat and Population	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Protect identified locations of Vole Ears Lichen		
INDICATOR	Administratively protect identified locations of Vole Ears Lichen according to SMP		
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY		
	Spatial data of known vole ears lichen has been provided to PHP by the NS Department of		

Annual	Environment. There are no known locations of vole ears lichen in the 7 eastern counties where PHP operates.
DATA SOURCES	COST AND DIFFICULTY
The Forest Manager (TFM); Boreal Felt Lichen Potential Habitat Layer	Low – There are no known locations of Vole Ears Lichen in PHP’s management area
FOREST MANAGEMENT PRESCRIPTION	
<p>In May 2018, the NSDNRR released new Special Management Practices for At-Risk Lichens for several lichen species including: boreal felt lichen, vole ears, Hibernia jellyskin lichen, powdered moon lichen, eastern waterfan, wrinkled shingle lichen, frosted glass-whiskers, black-foam lichen, blue felt lichen, and poor-man’s shingles lichen. The SMP is triggered by areas of overlap between planned forest management activities and the modelled lichen habitat for boreal felt lichen. Where there is overlap, a lichen survey is conducted and if any of the listed species in the SMP is found, a protection zone is established around the site. The sizes of these protection zones vary depending on the species. There are also restrictions related to road construction and maintenance.</p> <p>For vole ears lichen, a 200 metre radius protection zone is established around each site. In the protected zone, harvesting or silviculture, mineral exploration drill sites, or new road or trail construction (unless in exceptional circumstances and under approval by DNRR) are permitted. Existing road maintenance is permitted subject to review and conditions.</p>	
2022 MONITORING UPDATE	
There are two known locations (year 2010 and 2015) of vole ears lichen in PHP’s forest management area that are buffered with a 200 metre protection zone. Surveys completed in 2022 resulted in no new locations of vole ears lichen.	
SUPPORTING DOCUMENTS/REFERENCES	
COSEWIC Assessment and Status Report, 2009; NSDNRR; ACCDC 2019 Database	

## HCV – Blue Felt Lichen Occurrences



HCV ATTRIBUTE		Species at Risk – Habitat and Population	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Protect identified locations of Blue Felt Lichen		
INDICATOR	Administratively protect identified locations of Blue felt lichen according to SMP		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	Spatial data is collected annually by ACCDC for all rare species. Also, new locations are being found on PHP's Crown license during boreal felt lichen surveys.
DATA SOURCES	The Forest Manager (TFM); Boreal Felt Lichen Potential Habitat Layer	COST AND DIFFICULTY	Low – A survey was conducted for the two known locations of blue felt lichen in PHP's management area.
FOREST MANAGEMENT PRESCRIPTION			
<p>In May 2018, the NSDNRR released new Special Management Practices for At-Risk Lichens for several lichen species including: boreal felt lichen, vole ears, Hibernia jellyskin lichen, powdered moon lichen, eastern waterfan, wrinkled shingle lichen, frosted glass-whiskers, black-foam lichen, blue felt lichen, and poor-man's shingles lichen. The SMP is triggered by areas of overlap between planned forest management activities and the modelled lichen habitat for boreal felt lichen. Where there is overlap, a lichen survey is conducted and if any of the listed species in the SMP is found, a protection zone is established around the site. The sizes of these protection zones vary depending on the species. There are also restrictions related to road construction and maintenance.</p> <p>For blue felt lichen, a 100-metre radius protection zone is established around each site. In the protected zone, harvesting or silviculture, mineral exploration drill sites, or new road or</p>			

trail construction (unless in exceptional circumstances and under approval by DNRR) are permitted. Existing road maintenance is permitted subject to review and conditions.

#### 2022 MONITORING UPDATE

There were six new locations of blue felt lichen found in PHP's forest management area during lichen surveys conducted in 2022. A 100-meter no harvest buffer was applied to each location. In total, there are 429 locations of blue felt lichen on PHP's Crown license area. A proposed management plan for Blue Felt Lichen was released for public comment in late 2020 by Environment and Climate Change Canada. This plan references the management approach currently being implemented in Nova Scotia through the provincial At-Risk Lichens SMP.

#### SUPPORTING DOCUMENTS/REFERENCES

COSEWIC Assessment and Status Report, 2009; NSDNRR; ACCDC 2019 Database

### HCV – Eastern White Cedar



HCV ATTRIBUTE		Species at Risk – Habitat and Population
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Protect identified locations of Eastern White Cedar	
INDICATOR	Protection of all known locations of Eastern White Cedar	
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY Ensure all known locations of Eastern White Cedar in PHP's management area are protected from harvest activities.

<b>DATA SOURCES</b>	<b>COST AND DIFFICULTY</b>
The Forest Manager (TFM); NSDNRR, NSE, ACCDC databases	Low – PHP does not conduct include the harvest of Eastern White Cedar in its management
<b>FOREST MANAGEMENT PRESCRIPTION</b>	
- PHP does not include the harvest of Eastern White Cedar in its forest management. Queries of the NSDNRR forest inventory, as well as reviews of the rare species databases from NSDNRR, NSE, and ACDC, did not identify eastern white cedar stands for lands managed by PHP.	
<b>2022 MONITORING UPDATE</b>	
No known stands or individual trees of eastern white cedar have been found or are known to occur within PHP's area of operation.	
<b>SUPPORTING DOCUMENTS/REFERENCES</b>	
A Management Plan for Native Occurrences of Eastern White Cedar in Nova Scotia, 2010	

### HCV – Black Ash



<b>HCV ATTRIBUTE</b>	<b>Species at Risk – Habitat and Population</b>	
<b>OPERATIONAL MONITORING PROGRAM</b>		
<b>MANAGEMENT OBJECTIVE</b>	Protect identified locations of Black Ash	
<b>INDICATOR</b>	Protection of all known locations of Black Ash	
<b>MONITORING/REPORTING FREQUENCY</b>	<b>MONITORING STRATEGY</b>	

Annual	Ensure all known locations of Black Ash in PHP's management area are protected from harvest activities.
DATA SOURCES  The Forest Manager (TFM); NSDNRR, NSE, ACCDC databases	COST AND DIFFICULTY  Low – PHP does not conduct include the harvest of Black Ash in its management
<b>FOREST MANAGEMENT PRESCRIPTION</b>	
- PHP does not include the harvest of Black Ash in its forest management. Queries of the NSDNRR forest inventory, as well as reviews of the rare species databases from NSDNRR, NSE, and ACDC, did not identify black ash stands for lands managed by PHP.	
<b>2022 MONITORING UPDATE</b>	
<p>No new stands of black ash were identified in 2022.</p> <p>In the spring of 2021, a stand of black ash was discovered by a PHP Woodlands Supervisor during a routine survey of a 19.8 hectare area of forest in Cape Breton. This newly uncovered stand is significant because it is a cluster of trees, and there is evidence of seed-bearing trees, which indicates male and female trees in the area. Most known black ash populations in Cape Breton are reproducing by stump sprouts, not seed. Other trees easily out-compete black ash trees, but the large diameters of some trees indicate that they are mature and not suppressed. A formal survey of the property was completed to determine if more trees are present and to create a plan to protect the current stand.</p>	
<b>SUPPORTING DOCUMENTS/REFERENCES</b>	
NSDNRR, NSE, ACCDC databases	



## HCV – Frosted Glass Whiskers Habitat



HCV ATTRIBUTE		Species at Risk – Habitat	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Maintain Frosted Glass Whiskers Habitat		
INDICATOR	Administratively protect Frosted Glass Whiskers habitat identified in NSDNRR's Significant Habitat database and Atlantic Canada Conservation Data Centre database		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	All known locations of frosted glass whiskers are protected.
DATA SOURCES	The Forest Manager (TFM)	COST AND DIFFICULTY	Low – PHP does not conduct forest management activities within Frosted Glass Whiskers habitat
FOREST MANAGEMENT PRESCRIPTION			
<p>In May 2018, the NSDNRR released new Special Management Practices for At-Risk Lichens for several lichen species including: boreal felt lichen, vole ears, Hibernia jellyskin lichen, powdered moon lichen, eastern waterfan, wrinkled shingle lichen, frosted glass-whiskers, black-foam lichen, blue felt lichen, and poor-man's shingles lichen. The SMP is triggered by areas of overlap between planned forest management activities and the modelled lichen habitat for boreal felt lichen. Where there is overlap, a lichen survey is conducted and if any of the listed species in the SMP is found, a protection zone is established around the site. The sizes of these protection zones vary depending on the species. There are also restrictions related to road construction and maintenance.</p> <p>For frosted glass whiskers, a 100 metre radius protection zone is established around each site. In the protected zone, harvesting or silviculture, mineral exploration drill sites, or new road or</p>			

trail construction (unless in exceptional circumstances and under approval by DNRR) are permitted. Existing road maintenance is permitted subject to review and conditions.

#### 2022 MONITORING UPDATE

- One new location of frosted glass whiskers was found in 2022 during regular lichen surveys. This location was buffered with a 100-meter no harvest buffer.
- There are 12 known locations of frosted glass whiskers in eastern Nova Scotia. Five of these locations occur on privately owned land. The other seven locations are on PHP's Crown license area and are buffered with a protection zone.

#### SUPPORTING DOCUMENTS/REFERENCES

Management Plan for the Frosted Glass Whiskers, Nova Scotia Population, 2011; ACCDC 2019 Database; Boreal Felt Lichen Surveys, 2017-18

### HCV – Wrinkled Shingle Lichen Habitat



HCV ATTRIBUTE		Species at Risk – Habitat	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Maintain Wrinkled Shingle Lichen Habitat		
INDICATOR	Administratively protect Wrinkled Shingle Lichen habitat in identified locations by NSDNRR, ACCDC, or PHP lichen survey results.		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	All known locations of wrinkled shingle lichen are protected.
DATA SOURCES	The Forest Manager (TFM)	COST AND DIFFICULTY	Low – PHP does not conduct forest management activities within wrinkled shingle lichen habitat
FOREST MANAGEMENT PRESCRIPTION			

In May 2018, the NSDNRR released new Special Management Practices for At-Risk Lichens for several lichen species including: boreal felt lichen, vole ears, Hibernia jellyskin lichen, powdered moon lichen, eastern waterfan, wrinkled shingle lichen, frosted glass-whiskers, black-foam lichen, blue felt lichen, and poor-man's shingles lichen. The SMP is triggered by areas of overlap between planned forest management activities and the modelled lichen habitat for boreal felt lichen. Where there is overlap, a lichen survey is conducted and if any of the listed species in the SMP is found, a protection zone is established around the site. The sizes of these protection zones vary depending on the species. There are also restrictions related to road construction and maintenance.

For wrinkled shingle lichen, a 100 metre radius protection zone is established around each site. In the protected zone, harvesting or silviculture, mineral exploration drill sites, or new road or trail construction (unless in exceptional circumstances and under approval by DNRR) are permitted. Existing road maintenance is permitted subject to review and conditions.

#### 2022 MONITORING UPDATE

No new locations of wrinkled shingle lichen were found on PHP's Crown license area in 2022. There is one known location of wrinkled shingle lichen in eastern Nova Scotia, which was discovered in 2014. This location has a 100-meter no harvest buffer applied to it.

#### SUPPORTING DOCUMENTS/REFERENCES

COSEWIC Assessment and Status Report, 2017; ACCDC 2019 Database; Boreal Felt Lichen Surveys, 2017-18

## HCV – Cold Water Refugia Sub-watersheds



HCV ATTRIBUTE		Long-term hydrologic functions	
<b>OPERATIONAL MONITORING PROGRAM</b>			
MANAGEMENT OBJECTIVE	Maintenance of thermal cover for Atlantic Salmon and Brook Trout habitat		
INDICATOR	Maintain minimum 50% crown closure at the stand level in cold water refugia areas (total 30,015 hectares) with the exception of stands containing non-wind firm trees.		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	Monitor implementation of stand structure reserve using GIS overlay of completed harvest treatments with cold water refugia sub-watershed areas.
DATA SOURCES	The Forest Manager (TFM)	COST AND DIFFICULTY	Low – PHP monitors this internally with resources currently available.
<b>FOREST MANAGEMENT PRESCRIPTION</b>			
<ul style="list-style-type: none"> <li>- Cold water refugia areas are managed to maintain as much thermal cover as possible by leaving a minimum 50% crown closure wherever possible at the stand level following harvest treatments.</li> <li>- The only exception is in stands containing a high proportion of non-wind firm trees, such as balsam fir, black spruce, or white spruce that are vulnerable to blowdown.</li> <li>- No intensive forest management will occur in these HCVF's (i.e. establishing FSC defined plantations).</li> </ul>			

2022 MONITORING UPDATE

Below is a summary of treatments completed in cold water refugia areas in 2022. This is approximately 4% of the total area of cold water refugia (30,015 ha).

Treatment	Sum_Total_HA
Commercial Thinning	55.883995
Partial Overstory Removal Retain 1/5	46.318506
Variable Retention 10%	17.197749

SUPPORTING DOCUMENTS/REFERENCES

N/A

HCV – International Bird Areas



HCV ATTRIBUTE		Migratory birds habitat
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintain and/or enhance migratory bird habitat	
INDICATOR	Reserve stand structure in Important Bird Areas (IBA's)	
MONITORING/REPORTING FREQUENCY	Annual	<p><b>MONITORING STRATEGY</b></p> <p>Currently, all identified IBA's in PHP's operating area are not impacted by forest management activities due to their location (i.e. off shore islands or inaccessible forest areas). Specific protocols for monitoring birds at IBAs are in development and will leverage and adapt existing monitoring programs that are directly relevant to the IBA Program (IBA Canada website). Verify annually that spatial list of IBA's is up-to-date for PHP's operating area.</p>

<p>DATA SOURCES</p> <p>The Forest Manager (TFM); IBA Canada</p>	<p>COST AND DIFFICULTY</p> <p>Low – PHP does not conduct forest management activities in IBA’s, therefore, monitoring is not considered necessary.</p>
<p>FOREST MANAGEMENT PRESCRIPTION</p>	
<p>- The Scaterie Island IBA site is located within the boundaries of a legally-designated protected wilderness area. No harvesting is permitted to occur within this site.</p> <p>- Coastal IBA sites are not impacted by PHP’s forest management activities, therefore, no special management practices are required.</p> <p>- IBA sites Cape North and Central Cape Breton Highlands have been addressed in Category 1, Question 1 for Bicknell’s thrush. Additionally, the Cape North IBA site contains significant concentrations of Boreal owl. For this HCVF, no harvesting currently occurs and is not expected to occur in the future. Should harvest plans be developed, a management strategy for this HCVF will be developed.</p>	
<p>2022 MONITORING UPDATE</p>	
<p>PHP has not conducted any forest management activities in IBA’s identified within the forest management area.</p>	
<p>SUPPORTING DOCUMENTS/REFERENCES</p>	
<p>IBA Canada website <a href="http://www.ibacanada.ca/">http://www.ibacanada.ca/</a></p>	

## HCV – Red Spruce



HIGH CONSERVATION VALUE – RED SPRUCE	
HCV ATTRIBUTE	Natural Red Spruce Stands
OPERATIONAL MONITORING PROGRAM	
MANAGEMENT OBJECTIVE	Manage red spruce stands according to PHP Work Instruction for red spruce
INDICATOR	Management and maintenance of red spruce stands to improve the quality of uneven-aged conditions over time.
MONITORING/REPORTING FREQUENCY Annual	MONITORING STRATEGY Verify that annual harvest completions in natural red spruce stands were implemented using PHP's work instruction for red spruce management.
DATA SOURCES The Forest Manager (TFM)	COST AND DIFFICULTY Low – PHP has forest cover and historical data that shows natural red spruce stand locations. The PHP planner identifies these areas for management.
FOREST MANAGEMENT PRESCRIPTION	
<p><u>Objectives for Red Spruce Dominated Stands</u></p> <ul style="list-style-type: none"> <li>- Strive for two to three cohort stand structures.</li> <li>- Over time, we will strive to increase the area of multiple ages in many stands.</li> <li>- Promote natural red spruce regeneration</li> <li>- At harvest (other than tending), trees should be large and of high value. Management (spacings, thinnings) should be carried out to help meet this objective.</li> <li>- Retain some overstory structure, including snags; future snags; other tolerant species; and residual red spruce component – both individuals when windfirm and in clumps (structure and seed).</li> </ul>	

### Immature stands

Commercial thinning when windfirm.

An option for non-windfirm immature red spruce stands is to partially remove the overstory in 2-3+ stages separated by a period of 10 to 20 years. The trees retained should be windfirm and quality immature trees. This helps ensure increased value of residual stand and regeneration establishment, and subsequent regeneration release(s).

### Maturing stands

Shelterwood to promote regeneration.

As an option, a modified shelterwood treatment providing increased retention will be implemented, with a plan for two ages initially, with the intent of a third as the newly regenerated stand grows in to the existing overstory canopy. As possible considering tree ages and wind firmness:

- Step One: Initial shelterwood harvest is modified to include more patch retention, by doubling the present wildlife clump retention – move to 20 trees per hectare, with patches scattered throughout the treatment area. ie an irregular shelterwood
- Step Two once regeneration is 60 cm tall (5-10 yrs): Overstory harvest to release regeneration is needed (regeneration protection harvest techniques implemented). The retention includes both small patches of residuals, as well as individuals (as available, few isolated pines/hemlock/hardwoods, and snags with designated red spruce retention). Ten living trees per hectare are required.
- Shelterwood completed with adequate established regeneration.
- Step Three: 15-35 yrs The young and immature stand is tended as it grows (space – thin).
- Longer term: As trees grow into the upper canopy, some of the patches and individual trees will be harvested, excluding designated wildlife clumps and legacy trees.
- At this time (in the future), three cohorts are introduced into the stand with the intent of patterning an uneven-aged structure.

In some instances, trees in the forest stand planned for treatment are not wind-firm and excessive blowdown and significant wood losses would occur following implementing one of the treatments described above. If the stand is determined to be a high risk for blowdown, an alternative treatment may be implemented (over story removal and planting), or it should be left to grow until maturity then harvested.



## 2022 MONITORING UPDATE

Approximately 42 hectares of red spruce stands were treated in 2022.

Treatment	Sum_Total_HA
Commercial Thinning	47.503765
Variable Retention 10%	3.978492

## HCV – Protected Areas



HCV ATTRIBUTE		Protected Area
<b>OPERATIONAL MONITORING PROGRAM</b>		
MANAGEMENT OBJECTIVE	Establish protected areas (legal, pending, and/or administrative) in PHP's management area	
INDICATOR	Establishment of legal, pending, and/or administrative protected areas	
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY Continue to monitor provincial government's protected lands process for the establishment and legal protection of new wilderness areas and/or other decisions made regarding areas.
DATA SOURCES	The Forest Manager (TFM); NSDNRR; NSDOE	COST AND DIFFICULTY Low
<b>FOREST MANAGEMENT PRESCRIPTION</b>		
<ul style="list-style-type: none"> <li>- All identified forest lands for legal or pending protection by the provincial government have been delineated in TFM and are clearly marked as legal or pending protected areas.</li> <li>- All identified forest lands for administrative protection by PHP have been delineated in TFM and are clearly marked as administrative protected areas.</li> <li>- PHP staff is aware that no forest management activities are allowed to occur in these areas.</li> </ul>		

<b>Protected Area Category</b>	<b># of Sites</b>	<b>Total Hectares</b>
New Provincial Protected Area	89	98,184
Provincial Parks and Reserves	21	1,492
Provincial Nature Reserves	7	1,868
Provincial Wilderness Areas	19	106,526
National Migratory Bird Sanctuaries	1	392
National Parks	1	94,870
<b>TOTAL HECTARES</b>		<b>303,332</b>

<b>Administratively Protected Area Category</b>	<b># of Sites</b>	<b>Total Hectares</b>
Old Forest Areas	N/A	84,717
PHP Protected Area	8	6,147
IBP Sites & Sites of Ecological Significance	12	3,107
<b>TOTAL HECTARES</b>		<b>93,971</b>

**2022 MONITORING UPDATE**

A GIS overlay using completed harvest treatment data from 2022 shows that there have been no forest management activities conducted in legal or administrative protected areas.

**SUPPORTING DOCUMENTS/REFERENCES**

Our Parks and Protected Areas: A Plan for Nova Scotia, 2013; TFM Data; NSE Protected Areas

## HCV – Special Management Zone Adjacent to Protected Area Boundaries

HCV ATTRIBUTE	Limit Protected Area Access
<b>OPERATIONAL MONITORING PROGRAM</b>	
MANAGEMENT OBJECTIVE	Minimize road construction
INDICATOR	Minimize road construction to reduce access points into protected areas by implementing a 200-meter wide special management zone.
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY
Annual	Assess whether new roads have been built in the special management zone using GIS overlay.
DATA SOURCES	COST AND DIFFICULTY
The Forest Manager (TFM)	

	Low – PHP currently monitors the special management zone and road construction using TFM.
<b>FOREST MANAGEMENT PRESCRIPTION</b>	
- Minimize road construction to reduce access points into the protected area. If roads are needed, they are to build parallel to the protected area boundary to minimize access points.	
<b>2022 MONITORING UPDATE</b>	
No new roads were built in the special management zone adjacent to protected area boundaries in 2022. Also in 2021, NSDNRR implemented a 100-m buffer around all existing protected areas. These buffers are for implementing special management practices and/or additional protection measures for protected areas.	
<b>SUPPORTING DOCUMENTS/REFERENCES</b>	
N/A	

## HCVF Category 2 – Large Landscape Level Forests


### HCV – Intact Forest Landscapes

HCV ATTRIBUTE		Intact Forest Landscapes	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Adhere to FSC requirements for management of Intact Forest Landscapes		
INDICATOR	Maintain the integrity and intactness of intact forest landscapes.		
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY		
Annual	Assess whether more than 20% of the IFL has been impacted, and if the IFL has been reduced in size below 50,000 ha.		
DATA SOURCES	COST AND DIFFICULTY		
The Forest Manager (TFM)	Low – PHP currently monitors the IFL using TFM.		
FOREST MANAGEMENT PRESCRIPTION			
<p>There is one known Intact Forest Landscape as identified by Global Forest Watch Canada in PHP's forest management area. It is 103,849 ha and encompasses the Cape Breton Highlands National Park as well as other area outside the park. The total area of Crown land managed by PHP inside the IFL is 20,402 ha (20%). Of that 20%, approximately 10,000 ha has been identified as a pending new protected area by the provincial government with an existing additional 1,260 ha already established as a Crown Wilderness Area. These pending and existing protected areas are included in PHP's Crown license area. Approximately 9% of remaining area is potential operable forest area. When calculating the impact of PHP's operations in the IFL, the total area to be assessed will be 20,402 ha. PHP's total allowable harvest area in the portion of the IFL in the management unit is 4,080 ha.</p> <ul style="list-style-type: none"> <li>- Do not impact more than 20% of Intact Forest Landscapes within the Management Unit,</li> <li>- Do not reduce any IFLs below the 50,000 ha threshold in the landscape.</li> </ul>			
2022 MONITORING UPDATE			
Year	Total Harvest (hectares)	Total Silviculture (hectares)	Total Roads Built (km)
2017	0	0	0
2018	0	0	0

<b>2019</b>	0	0	0
<b>2020</b>	23.7	0	0
<b>2021</b>	0	0	0
<b>2022</b>	0	0	0

**SUPPORTING DOCUMENTS/REFERENCES**

Global Forest Watch; FSC Advice Note on Intact Forest Landscapes



## HCV – Large Landscape Level Forests

HCV ATTRIBUTE		Biodiversity and Intactness	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	To maintain biodiversity values and intactness in large landscape level forests		
INDICATOR	Manage large landscape level forests with special practices in protected areas, core roadless areas, and special management areas		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	
		Assess management activities within large landscape level forests to ensure practices comply with requirements outlined for protected areas, core roadless areas, and special management areas.	
DATA SOURCES	The Forest Manager (TFM)	COST AND DIFFICULTY	
		Low – PHP currently monitors large landscape level forests using TFM.	
FOREST MANAGEMENT PRESCRIPTION			
<ul style="list-style-type: none"> <li>- No new roads in Core Roadless areas</li> <li>- For HCV area outside core roadless, follow road design objectives as shown below. Road Index value at HCV level not to exceed 0.58 km/km<sup>2</sup>. If feasible and where necessary, block off access to reduce road travel.</li> </ul>			
<p>Maximum road density = 0.58 km/km<sup>2</sup></p> <p>✓ - build roads in from the edge where possible</p> <p>✓ - close access between resource use</p> <p>X - avoid roads that will fragment the patch</p> <p>X - avoid loop roads</p> <p>Roadless Core Area</p>			
<ul style="list-style-type: none"> <li>- Use the provincial Forest Ecosystem Classification Guide to identify ecosite level prescriptions that: <ul style="list-style-type: none"> <li>○ Promote ecosite patches by combining stands through treatment</li> <li>○ Employ 'extensive' management practices that support: <ul style="list-style-type: none"> <li>- natural regeneration</li> </ul> </li> </ul> </li> </ul>			

- longer rotations with consideration of natural disturbance processes
- tree species diversity consistent with the vegetation type, while promoting those that support long-term resilience (i.e. best options for future)
- No full-tree logging
- Reduce road length by increasing average forwarding distance targets by 20% (from 250 m to 300 m)
- Bridge construction may be temporary and removed as practical
- Retain minimum 60% area in non-clearcut condition (at the HCV level). Non-clearcut defined as forest stand greater than 10 years of age.
- No FSC plantations / Intensive management
- No planting of exotic species
- Acadian Forest Restoration (considering N.S. Forest Code; FSC)
- Management will align with natural disturbance regimes
- Application of Forest Ecosystem Classification to identify appropriate treatments
- Appropriate forest covertype management: Use of hardwood management keys
- Appropriate forest covertype management: Use of mixedwood management keys
- Natural regeneration where appropriate
- Appropriate use of PHP's 12 different harvest techniques (CC, PC, SW, ST, Single, Group, Patch, CT, OR, CTR, RS, SC)
- Species at Risk Recovery Strategy/SMP Implementation
- No herbicides
- Steep Slope Exclusion
- Leave patches (e.g. active eagle/hawk nest sites, inoperable areas, vernal pools, DNR requests during approval process)

#### 2022 MONITORING UPDATE

PHP uses the provincial Forest Ecosystem Classification manual for all forest management decisions. PHP does not conduct full-tree logging, plant exotic species, use herbicides, and has not implemented intensive management in these areas. All other management prescriptions mentioned above are implemented across the entire forest management area.

The below table summarizes the current status of each large landscape level forest with respect to road index and minimum non-clearcut condition. Non-clearcut condition is defined as anything greater than 10 years of age. No new roads were built in large landscape level forests in 2022, so the road index remains the same as in 2016.

		Road Index Maximum Allowed = 0.58 km/km <sup>2</sup>		Minimum Non-clearcut Condition = 60%
HCVF LLLF Name	Total HA	2022 Road Index	Future Road Index	Non-clearcut Condition in 2022
Barren Hill	1,318	0.08 km/km <sup>2</sup>	0.20 km/km <sup>2</sup>	99%
Boisdale Hills	5,630	0.40 km/km <sup>2</sup>	0.52 km/km <sup>2</sup>	100%
Bornish Hill (fully protected)	2,106	0 km/km <sup>2</sup>	0 km/km <sup>2</sup>	100%
Country Harbour	8,202	0.03 km/km <sup>2</sup>	0.03 km/km <sup>2</sup>	100%
East Bay Hills	1,865	0.23 km/km <sup>2</sup>	0.31 km/km <sup>2</sup>	98%
French River	25,226	0 km/km <sup>2</sup>	0 km/km <sup>2</sup>	100%
Hill Lake	877	0.55 km/km <sup>2</sup>	0.65 km/km <sup>2</sup>	100%
Ingonish River	15,210	0.01 km/km <sup>2</sup>	0.01 km/km <sup>2</sup>	100%
Isaacs Harbour River	6,157	0.25 km/km <sup>2</sup>	0.42 km/km <sup>2</sup>	98%
Jim Campbells Barren (fully protected)	4,586	0.21 km/km <sup>2</sup>	0.21 km/km <sup>2</sup>	100%
Masons Mountain (fully protected)	1,022	0.06 km/km <sup>2</sup>	0.06 km/km <sup>2</sup>	100%
North River	6,328	0.20 km/km <sup>2</sup>	0.20 km/km <sup>2</sup>	100%
Oban	1,618	0.57 km/km <sup>2</sup>	0.78 km/km <sup>2</sup>	89%



Petit Lake Ruiss Noir (fully protected)	1,612	0 km/km <sup>2</sup>	0 km/km <sup>2</sup>	100%
Salmon Gaspereaux	2,357	0.30 km/km <sup>2</sup>	0.61 km/km <sup>2</sup>	94%
Upper Liscomb River	7,398	0.07 km/km <sup>2</sup>	0.07 km/km <sup>2</sup>	100%
<b>TOTAL HECTARES</b>	<b>91,512</b>		Future index may exceed maximum 0.58 km/km <sup>2</sup> . Will need to manage road index to meet target.	

HCVF LLLF Name	Total HA	2021 Total Area Treated	Treatment Used
----------------	----------	-------------------------	----------------

Barren Hill	1,318	No area treated	
Boisdale Hills	5,630	No area treated	
Bornish Hill (fully protected)	2,106	No area treated	
Country Harbour	8,202	No area treated	
East Bay Hills	1,865	No area treated	
French River	25,226	No area treated	
Hill Lake	877	No area treated	
Ingonish River	15,210	No area treated	
Isaacs Harbour River	6,157	No area treated	

Jim Campbells Barren (fully protected)	4,586	No area treated	
Masons Mountain (fully protected)	1,022	No area treated	
North River	6,328	No area treated	
Oban	1,618	No area treated	
Petit Lake Ruiss Noir (fully protected)	1,612	No area treated	
Salmon Gaspereaux	2,357	No area treated	
Upper Liscomb River	7,398	No area treated	
<b>TOTAL HECTARES</b>	<b>91,512</b>	<b>No area treated</b>	

## HCVF Category 3 – Rare, Threatened or Endangered Ecosystems

### HCV – Significant Ecosites

HCV ATTRIBUTE		Rare, threatened or endangered ecosystems	
OPERATIONAL MONITORING PROGRAM			
MANAGEMENT OBJECTIVE	Maintain rare, threatened or endangered ecosystems		
INDICATOR	Rare, threatened or endangered ecosystems administratively protected from forest management activities		
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY	
		Monitor rare, threatened or endangered ecosystems to ensure they are administratively protected from forest management activities. Exception applies if the mapped ecosystem type does not match on-the-ground characteristics.	
DATA SOURCES	The Forest Manager (TFM); NSDOE	COST AND DIFFICULTY	
		Low	
FOREST MANAGEMENT PRESCRIPTION			
<p>- All significant ecosites are administratively protected from forest management activities with the following exceptions:</p> <ul style="list-style-type: none"> <li>- Karst conifer forest, karst hardwood forest, calcareous forest, and hemlock forest that have been previously managed will continue to be managed to maintain and restore mature climax conditions.</li> <li>- Significant ecosites are identified using the provincial forest inventory data and there has been limited field verification, so there is a certain amount of ambiguity within the dataset. Since there may be data inaccuracies between the digital information versus on-the-ground characteristics, stands that do not match the inventory data are exempt from special management activities as outlined here.</li> </ul>			
2022 MONITORING UPDATE			
A GIS overlay using completed harvest treatment data from 2022 and significant ecosite data shows that there was 2 hectares managed in a significant ecosite categorized as a coastal barren. Pre-treatment assessment data has verified that the areas managed did not meet the criteria for significant ecosite conditions.			
SUPPORTING DOCUMENTS/REFERENCES			

Significant Ecosite data layer, NSDOE

### HCV – Significant, Old or Unique Forests

HCV ATTRIBUTE	Rare, threatened or endangered ecosystems	
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintain rare, threatened or endangered ecosystems	
INDICATOR	Rare, threatened or endangered ecosystems administratively protected from forest management activities	
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY	
Annual	Monitor rare, threatened or endangered ecosystems to ensure they are administratively protected from forest management activities. Exception applies if the mapped ecosystem type does not match on-the-ground characteristics.	
DATA SOURCES	COST AND DIFFICULTY	
The Forest Manager (TFM); NSDOE	Low	
FOREST MANAGEMENT PRESCRIPTION		
<p>- All significant, old or unique forests (SOUF) are administratively protected from forest management activities that meet the following species composition (SOUF code in brackets). Exception applies if the mapped ecosystem type does not match on-the-ground characteristics.</p>		
<b>Species composition</b>		<b>Stand height</b>
70% or more spruce or red spruce (SPRU)		≥17m
50% or more eastern hemlock (HEML)		≥15m
50% or more white pine (WHPI)		≥18m
70% or more climax coniferous species with the most common species no more than 60% (MCCO)		≥17m
50% or more sugar maple (SUMA)		≥17m
70% or more climax deciduous species or tolerant hdwd (MCDE)		≥17m
70% or more climax coniferous or deciduous species with neither group exceeding 60% (MCCD)		≥17m

30% or more red pine (excl. plantations) (REPIO)	≥12m
50% or more Beech	≥18m
<b>2022 MONITORING UPDATE</b>	
A GIS overlay using completed harvest treatment data from 202 shows no SOUF stand was managed.	
<b>SUPPORTING DOCUMENTS/REFERENCES</b>	
Significant, old or unique data layer, NSDOE	

### HCV – Old Forest

HCV ATTRIBUTE		Old Forest Protected Area
<b>OPERATIONAL MONITORING PROGRAM</b>		
MANAGEMENT OBJECTIVE	Establish old forest protected areas on land-base	
INDICATOR	Establishment and legal protection of old forest protected areas	
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY	
Annual	Monitor old forest protected areas TFM to ensure no forest management activities are conducted.	
DATA SOURCES	COST AND DIFFICULTY	
The Forest Manager (TFM); NSDNRR; NSDOE	Low	
<b>FOREST MANAGEMENT PRESCRIPTION</b>		
<ul style="list-style-type: none"> <li>- All identified old forest areas are legally protected by the provincial government.</li> <li>- PHP staff is aware that no forest management activities are allowed to occur in these areas.</li> </ul>		
<b>2022 MONITORING UPDATE</b>		
A GIS overlay using completed harvest treatment data from 2022 shows that there have been no forest management activities conducted in the old forest areas identified by the provincial government. In 2018, the NS Department of Natural Resources & Renewables initiated a new old-growth forest protocol for the assessment of mature climax hardwood stands greater than 11 meters height based on forest inventory data. Since that time, an additional 3,235		

hectares of old-growth areas have been identified and protected on PHP's Crown license area.

#### SUPPORTING DOCUMENTS/REFERENCES

Old forest GIS layer, NSDNRR

## HCV – Poorly Represented Ecosystems

HCV ATTRIBUTE		Protection of Poorly Represented Ecosystems
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Establish protection of poorly represented ecosystems on land-base	
INDICATOR	Establishment and administrative protection of poorly represented ecosystems	
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY Monitor poorly represented ecosystems in TFM to ensure no forest management activities are conducted.
DATA SOURCES	The Forest Manager (TFM)	COST AND DIFFICULTY Low
FOREST MANAGEMENT PRESCRIPTION		
<ul style="list-style-type: none"> <li>- All identified poorly represented ecosystems are administratively protected by PHP.</li> <li>- PHP staff is aware that no forest management activities are allowed to occur in these areas.</li> </ul>		
<b>Poorly Represented Ecosystem</b>		<b>Total Hectares</b>
Masons Mountain		197
Jim Cambells Barren		2,844
Boisdale Hills		1,727
Country Harbour		829
North River		27
Oban		170
Hill Lake		113
Salmon Gaspereaux		240
<b>TOTAL HECTARES</b>		<b>6,147</b>

<b>2022 MONITORING UPDATE</b>
There have been no management activities in the above PHP administratively protected areas.
<b>SUPPORTING DOCUMENTS/REFERENCES</b>
PHP Gap Analysis Report

### HCV – Connectivity Management Zones

HCV ATTRIBUTE	<b>Continuous Canopy cover</b>	
<b>OPERATIONAL MONITORING PROGRAM</b>		
MANAGEMENT OBJECTIVE	Maintain continuous canopy cover between protected areas and old forest areas	
INDICATOR	Maintain a 100-meter wide continuous canopy cover (minimum 30%) corridor within the 500-meter wide Connectivity Management Zone (CMZ)	
MONITORING/REPORTING FREQUENCY	Annual	MONITORING STRATEGY
		Monitor 100 meters within the CMZ to ensure a continuous canopy cover and CMZ's are not severed across their width.
DATA SOURCES	The Forest Manager (TFM)	COST AND DIFFICULTY
		Low
<b>FOREST MANAGEMENT PRESCRIPTION</b>		
<p>- The Connectivity Management Zones will be managed to provide continuous canopy cover (minimum 30%) within the 500-meter wide corridors, which will include a solid 100-meter wide core zone. Although harvesting can occur within the CMZ's, these corridors will not be severed across their width.</p> <p>- The 500-meter wide CMZ's are static on the landscape, but the 100-meter wide core zone can 'move' within the CMZ.</p>		
<b>2022 MONITORING UPDATE</b>		
The Connectivity Management Zones continue to maintain a continuous canopy cover within the 100-meter wide core zone.		

## SUPPORTING DOCUMENTS/REFERENCES

The Forest Manager

## HCV – Margaree &amp; St. Mary's River Watershed

HIGH CONSERVATION VALUE – MARGAREE & ST. MARY'S RIVER WATERSHED	
HCV ATTRIBUTE	Non-clearcut Condition
OPERATIONAL MONITORING PROGRAM	
MANAGEMENT OBJECTIVE	To maintain a high level of non-clearcut condition in St. Mary's and Margaree Watersheds, and restoration management
INDICATOR	Each watershed shall have minimum 80% of its area (that is managed by PHP) in a non-clearcut condition, and 90% of each watershed shall be managed for restoration (i.e. no more than 10% of each watershed will be established as a FSC plantation).
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY
Annual	Monitor non-clearcut condition in each watershed to ensure target of minimum 80% is met.
DATA SOURCES	COST AND DIFFICULTY
The Forest Manager (TFM)	Low
FOREST MANAGEMENT PRESCRIPTION	
<ul style="list-style-type: none"> <li>- The St. Mary's and Margaree watersheds will be managed to maintain 80% or more of all lands managed by PHP in the watershed in a closed forest condition (&gt; 12 years of age).</li> <li>- Additionally, PHP will maintain at least 90% of the St. Mary's and Margaree watersheds in a natural condition for restoration, and will establish 200 m forest restoration zones (i.e. non-intensive management) along all main watercourses.</li> </ul>	
2022 MONITORING UPDATE	



PHP has been monitoring the non-clearcut condition in these watersheds for several years. See Indicator 3.2 on page 23 for current condition of St. Mary's and Margaree watersheds. Since 2008, these two watersheds have maintained minimum 80% non-clearcut condition.

#### SUPPORTING DOCUMENTS/REFERENCES

The Forest Manager

## HCVF Category 4 – Basic Services of Nature

### HCV – Legally Protected Municipal Water Supply Areas

HCV ATTRIBUTE	Water Health	
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Maintain water health for communities	
INDICATOR	Implement water protection measures in legally protected municipal water supply areas	
MONITORING/REPORTING FREQUENCY Annual	MONITORING STRATEGY Monitor implementation of water protection measures.	
DATA SOURCES The Forest Manager (TFM); NSDOE	COST AND DIFFICULTY Low	
FOREST MANAGEMENT PRESCRIPTION		
There is no land managed by PHP within the water supply areas unless requested or approved by the municipality through a watershed committee.		
2022 MONITORING UPDATE		
A GIS overlay using completed harvest treatment data from 2021 shows that there was no harvest inside designated municipal water supply areas.		
SUPPORTING DOCUMENTS/REFERENCES		
Nova Scotia Department of Environment		

## HCV – Water Supply Intake Areas

HCV ATTRIBUTE	Water Health	
<b>OPERATIONAL MONITORING PROGRAM</b>		
MANAGEMENT OBJECTIVE	Maintain water health for communities	
INDICATOR	Implement water protection measures around water supply intake areas.	
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY	
Annual	Monitor implementation of water protection measures.	
DATA SOURCES	COST AND DIFFICULTY	
The Forest Manager (TFM); NSDOE	Low	
<b>FOREST MANAGEMENT PRESCRIPTION</b>		
- Water protection measures include the Wildlife and Habitat Watercourse Protection Regulations, monitoring of % closed forest condition, steep slope management, HCVF aquatic watershed management, and rutting and ground disturbance guidelines.		
<b>2022 MONITORING UPDATE</b>		
A GIS overlay of completed harvest treatments and water supply intake areas shows no hectares were managed within the intake areas.		
<b>SUPPORTING DOCUMENTS/REFERENCES</b>		
Nova Scotia Department of Environment		

## HCV – Steep Slopes

HCV ATTRIBUTE	Soil Health; Community Health	
<b>OPERATIONAL MONITORING PROGRAM</b>		
MANAGEMENT OBJECTIVE	Maintain soil health and community health	

INDICATOR	No conventional harvesting in steep slope areas (30% average slope or greater)	
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY	
Annual	Monitor steep slope areas and conventional harvesting activities.	
DATA SOURCES	COST AND DIFFICULTY	
The Forest Manager (TFM)	Low	
<b>FOREST MANAGEMENT PRESCRIPTION</b>		
- Conventional harvesting is not permitted in areas with 30% average slope or greater. Non-conventional harvesting such as cable logging is permitted, however, PHP is currently not using this practice.		
<b>2022 MONITORING UPDATE</b>		
See page 22, indicator 3.1 for update.		
NOTE: This indicator is based on spatial data that identifies slopes > than 30% average using contour data. It is not based on the actual % slope for any given area as could be determined on-the-ground. Therefore, to calculate the results for the indicator, a GIS exercise is done which overlaps the steep slope data with completed harvest jobs to determine non-conformances. Most often, the areas showing as harvested are slivers due to inaccuracies in the data.		
<b>SUPPORTING DOCUMENTS/REFERENCES</b>		
The Forest Manager		

## HCVF Category 5 – Basic Needs of Local Communities

### HCV – Cattle Grazing on Cape Breton Highlands

HCV ATTRIBUTE		Local communities
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Support needs of local communities	
INDICATOR	Cattle grazing on the Cape Breton Highlands is allowed	
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY	
Annual	Monitor any issues arising from cattle grazing on Cape Breton Highlands	
DATA SOURCES	COST AND DIFFICULTY	
N/A	Low	
FOREST MANAGEMENT PRESCRIPTION		
<ul style="list-style-type: none"> <li>- Local farmers have let their cattle graze on the Cape Breton Highlands for several years during the summer/fall months.</li> <li>- PHP does not restrict this use.</li> </ul>		
2022 MONITORING UPDATE		
No issues have arisen in 2022 regarding cattle grazing in the Cape Breton Highlands.		
SUPPORTING DOCUMENTS/REFERENCES		
N/A		

### HCV – Viewshed Areas

HCV ATTRIBUTE		Local Communities
OPERATIONAL MONITORING PROGRAM		
MANAGEMENT OBJECTIVE	Minimize visual impacts to local communities from harvest activities	

INDICATOR	Implement work instruction 'Harvest View from Roadside'	
MONITORING/REPORTING FREQUENCY  Annual	MONITORING STRATEGY  Monitor any issues in identified viewshed areas arising from harvest activities.	
DATA SOURCES  The Forest Manager (TFM); Harvest View from Roadside Work Instruction	COST AND DIFFICULTY  Low	
FOREST MANAGEMENT PRESCRIPTION		
<p>District staff is responsible for determining the visibility rating using the TFM layer 'Viewsheds'. A harvest area determined to be within the <b>low category</b> will not require any specific landscape planning beyond regular housekeeping measures and removal of unsightly damaged residual trees.</p> <p>Cut blocks falling in <b>the medium category</b> on the visibility grid should be designed using the "Landscape Level" instructions in the Harvest View from Roadside Work Instruction.</p> <p>Blocks falling into the <b>high visibility category</b> will follow the "Landscape level", "Stand level" and "Road design" practices as applicable.</p>		
2022 MONITORING UPDATE		
<p>Forest stands treated in 2022 that overlapped with low, medium or high viewshed areas are below. These treatments include special management practices that meet PHP's procedure for minimizing impacts to medium and high viewscape areas.</p>		

Treatment	Visibility	Hectares
Variable Retention 30%	Low	8.891796
Variable Retention 20%	Low	4.173377
Variable Retention 10%	Low	1.810581
Commercial Thinning	Low	16.810886
Variable Retention 30%	Moderate	12.843432
Variable Retention 30%	Low	2.229244
Commercial Thinning	Moderate	2.608158
Variable Retention 10%	Low	3.377708
Variable Retention 10%	Moderate	5.111414
Variable Retention 20%	Moderate	5.598538
Variable Retention 10%	Low	25.310128
Variable Retention 30%	Moderate	8.323431
Variable Retention 20%	Low	8.435176
Commercial Thinning	Moderate	8.831779
Commercial Thinning	High	14.919628
Variable Retention 10%	Moderate	8.024412
Variable Retention 10%	Moderate	20.796022
Commercial Thinning	Low	53.082581

**SUPPORTING DOCUMENTS/REFERENCES**

Viewshed layer in TFM; Harvest View from Roadside Work Instruction

## HCVF Category 6 – Traditional Cultural Identity

### HCV – Forest Values and Uses

HCV ATTRIBUTE	First Nations Forest Values and Uses
<b>OPERATIONAL MONITORING PROGRAM</b>	
MANAGEMENT OBJECTIVE	Minimize impacts to First Nations Forest Values and Uses
INDICATOR	Implement work instruction 'Aboriginal Value – Suspending Operations'
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY
Annual	Monitor any issues identified during annual review of operations plans with First Nations communities. PHP also maintains a public inquiry database, which captures concerns or questions the general public may have regarding planned operations.
DATA SOURCES	COST AND DIFFICULTY


The Forest Manager (TFM); Public Inquiry Database	Low
<b>FOREST MANAGEMENT PRESCRIPTION</b>	
- If operations plans are known to affect First Nations forest values or uses through a review of annual operating plans or public inquiries, PHP will suspend all activities until a resolution is found.	
<b>2022 MONITORING UPDATE</b>	
- There were no public inquiries in 2022 related to PHP's operating plans that may affect First Nations. - There were no harvests in 2022 that occurred in identified medicinal plant areas.	
<b>SUPPORTING DOCUMENTS/REFERENCES</b>	
PHP Work Instruction 'Aboriginal Values – Suspending Operations'	

### HCV – Traditional Cultural Identity

HCV ATTRIBUTE	First Nations Traditional Cultural Identity	
<b>OPERATIONAL MONITORING PROGRAM</b>		
MANAGEMENT OBJECTIVE	Minimize impacts to First Nations Traditional Cultural Identity	
INDICATOR	Successful implementation of a Free Prior Informed Consent (FPIC) Agreement with Nova Scotia Mi'kmaq	
MONITORING/REPORTING FREQUENCY	MONITORING STRATEGY	
Annual	Monitor development and implementation of a FPIC Agreement with Nova Scotia Mi'kmaq	
DATA SOURCES	COST AND DIFFICULTY	
The Forest Manager (TFM); Public Inquiry Database	Moderate	
<b>BACKGROUND</b>		
Port Hawkesbury Paper initiated a Free Prior Informed Consent process with the Mi'kmaq Forestry Initiative Committee in 2020. The purpose of this process is to engage Mi'kmaq		

communities that may have legal and/or customary rights affected by PHP's forest management activities so that participation in forest management planning can be conducted to the extent necessary to protect their rights, resource lands and territories.

FPIC is a collective right held by Indigenous Peoples and recognized in international law and other agreements such as the United Declaration on Rights of Indigenous Peoples (UNDRIP 2007). Indigenous peoples have the right (legal and customary) to participate in decisions that could affect their rights, property, cultures and environment. In this broad context, Indigenous peoples have the right to make their own decisions to say 'yes' or 'no' whenever governments or corporations propose actions that could impact their lives and futures. FPIC includes the right to grant, modify, withhold or withdraw approval. It is not a one-time process but may occur at different levels of decision making and multiple times over life of a plan



#### 2022 MONITORING UPDATE

Work on the FPIC process began in the summer of 2020 with the Mi'kmaq Forestry Initiative Committee. Since that time, two working committees have been formed as well as a project plan. The Process Committee is tasked with identifying the appropriate process for conducting the FPIC process and the Technical Committee is responsible for sharing of data and information to assist the process. Currently, an Umbrella Framework Agreement is being developed to outline various collaborative initiatives between First Nations and PHP. These initiatives include FPIC as well as Indigenous Protected & Conservation Areas, Mi'kmaq Forestry Initiative lands, and other areas of mutual interest. An FPIC engagement plan is still to be developed to ensure culturally appropriate engagement is established with Mi'kmaq communities.

A request was received from an elder of Membertou First Nation to harvest approximately 80 pole-sized softwood trees for construction of two teepees for Mi'kmaq traditional gathering. The request was approved with no conditions applied.